


A New NMR Technique for Measuring the Ground State Populations of Formate Esters: Polar Aprotic Solvents Favor the *s*-cis (*E*)-Isomer of *t*-Butyl Formate

Michael E. Jung* and Jacquelyn Gervay¹

Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90024

Abstract: A new NMR technique for measuring the conformational population of *t*-butyl formate in a rapidly equilibrating mixture ($^3J_{\text{CH}}$ averaging) has been developed and used to show that polar aprotic solvents favor the *E*-isomer.


Recently we reported² a dramatic and synthetically quite useful solvent effect on the rate of the intramolecular Diels-Alder cycloaddition of substituted furfuryl methyl fumarates **1**. Polar solvents increased the rate of cyclization significantly, e.g., the ratio of the rates of cyclization of the dihydrido compound **1** ($R=R'=H$) in DMSO vs. toluene at 25° C is 3200. We attributed this effect to the fact that due to dipole overlap, the transition state **3** is more polar than the *s*-trans or *Z* ground state ester conformation **1** and thus polar solvents would be expected to lower the energy of activation and thereby increase the rate. We now report the use of a new NMR technique which shows that this polar



effect applies to ground state conformations as well, namely, that polar solvents stabilize the less stable but more polar *s*-cis or *E* ground state ester conformation such as **2**. By proper choice of polar solvent, as much as 33% of the less stable *s*-cis or *E* conformation can be present in equilibrium at 25° C.

Most esters exist almost completely in the *s*-trans or *Z* conformation with little or none of the *s*-cis or *E* conformation present at room temperature ($\Delta G^\circ > 4$ kcal/mole).³ The most common explanations for this preference for the *Z* conformation are dipole-dipole interactions,⁴ lone pair- σ^* interactions,⁵ and aromatic stabilization.⁶ Bulky alkyl formates, e.g., *t*-butyl formate **5**, generally have the highest percentage of the *E* isomer because the steric interaction of the *t*-butyl group with the carbonyl oxygen is greater than that with the formyl hydrogen and thus this steric interaction mitigates the preference for the *Z* isomer due to dipole interactions. For example, Ōki and Nakanishi

reported that the ^1H NMR spectrum of a 50% solution of **5** in dimethylformamide (DMF) at -94°C showed two singlets (at 8.0 and 8.9 ppm, respectively) due to the *Z* and *E* isomers in a ratio of 86:14.⁷ We reasoned that since the *E* isomer is more polar than the *Z* due to dipole overlap,⁷ this isomer should be favored in polar solvents and therefore its percentage should increase. This is indeed the case!



However, since the free energy of activation for the *Z* to *E* isomerization is only about 9-11 kcal/mole,⁸ a method had to be developed to study this equilibrium in certain polar solvents that could not be cooled to the very low temperatures (ca. -100°C) where the equilibrium is slow enough for the individual isomers to be seen. Drakenberg and Forsén⁹ had published a method based on a 5-bond coupling between the formyl and methyl hydrogens of **5** at 60 MHz, in which they made several assumptions in order to estimate the $^5J_{\text{HH}}$ for each isomer. Their indirectly estimated values were $^5J_{\text{trans}} = 0.42\text{ Hz}$ and $^5J_{\text{cis}} = 0.11\text{ Hz}$. We had trouble in obtaining well-split spectra at very low temperatures where the $^5J_{\text{HH}}$ could be measured experimentally for each isomer and therefore decided to develop a new technique to determine the percentages of both isomers at higher temperatures. We expected that the 3-bond coupling between the formyl hydrogen and the quaternary *t*-butyl carbon, $^3J_{\text{CH}}$ would be significantly different in each isomer and therefore could be used in a *J* averaging method to determine the isomeric composition.¹⁰ The 125 MHz ^{13}C NMR spectrum of a 10% solution of *t*-butyl formate **5** in several solvents at temperatures of -100°C to -122°C ¹¹ (with selective decoupling of the singlet due to the methyls of the *t*-butyl group) showed a large downfield doublet for the *Z* isomer in all solvents and a much smaller upfield signal for the *E* isomer. This upfield peak could be resolved into a clean doublet in only two solvent mixtures, dimethyl ether/10% acetone- d_6 at -111°C and acetaldehyde/10% acetone- d_6 at -121°C (the acetone- d_6 was added for a deuterium lock). From these spectra, the measured coupling constants were *Z* isomer, $J = 4.4\text{ Hz}$; *E* isomer, $J = 1.3\text{ Hz}$. Integration of the absorptions in the ^{13}C spectra gave the isomeric composition shown in Table 1. As expected, as the polarity of the solvent is increased from carbon disulfide and toluene to acetone and DMF, the percentage of the more polar *E* isomer increases (from ~ 0 to 17%). It is

Table 1: Isomeric Composition of *t*-Butyl Formate **5** at Low Temperature^a

Solvent	Temperature	% <i>Z</i>	% <i>E</i>
DMF- d_7	-100°C	83	17
Acetone- d_6	-95°C	86	14
Acetaldehyde + 10% Acetone- d_6	-121°C	87	13
Dimethyl ether + 10% Acetone- d_6	-111°C	95	5
Ethanol + 10% Acetone- d_6	-101°C	95	5
Tetrahydrofuran- d_8	-122°C	96	4
Toluene- d_8	-106°C	~ 100	~ 0
Carbon disulfide + 10% CD_2Cl_2	-100°C	~ 100	~ 0

a) A 10% solution of **5** in the indicated solvent was used in all cases except DMF where a 50% solution was used (in order to be able to cool to -100°C without freezing).

reassuring to note that the percentages derived from the ^{13}C integration for the sample in DMF- d_7 matches well those calculated by Ōki and Nakanishi⁷ by the ^1H integration (83:17 vs. 86:14). A variable temperature study was carried

out in several solvents and the $^3J_{\text{CH}}$ for the single peak observed for the ethereal carbon measured carefully in each. The percentage of the *E* isomer of **5**, calculated by the *J* averaging method, is given in Table 2 along with the dielectric constant ϵ and E_{T} parameter for each solvent (the solvents are listed in order of decreasing dielectric constant). Inspection of the results at 24° C shows that with the exception of the protic solvents, ethanol and *N*-methylformamide (NMF), the population of the *E* isomer increases linearly with dielectric constant (and with E_{T} except

Table 2: Percentage of *E* isomer of *t*-Butyl Formate **5** with Changes in Solvent^a

Solvent	-25°C	24°C	77°C	110°C	$\epsilon_{25^\circ\text{C}}^{\text{b}}$	E_{T}^{c}	E_{T}^{b}
NMF + 10% Acetone-d ₆	16.0	6.0	~ 0	~ 0	166.2	-----	52.9
DMSO-d ₆	-----	32.9	31.6	25.5	48.9	45.0	45.0
CD ₃ CN	30.0	27.4	28.7	-----	37.5	47.5	46.0
DMF-d ₇	31.9	26.4	26.4	19.4	36.7	43.7	43.8
Ethanol	6.5	16.1	-----	-----	24.3	51.2	51.9
Acetone-d ₆	23.5	21.0	-----	-----	20.5	42.0	42.2
CCl ₄ + 20% CDCl ₃	14.6	-----	-----	-----	2.72	37.8	33.8
Toluene-d ₈	11.6	12.3	11.0	8.4	2.38	35.7	33.9

- a) A 10% solution of **5** in the indicated solvent was used.
 b) The dielectric constants and reported E_{T} values for the pure solvents are from reference 12. The values for the mixtures were calculated by using a weighted average.
 c) The E_{T} values were obtained by dissolving Reichardt's dye in the actual sample used for the NMR study and recording the UV absorbance.

also for DMSO, which seems to be more polar than its E_{T} value would indicate).¹² Thus, as predicted, as the polarity of the solvent increases, the percentage of the more polar conformation increases. The ground state free energy difference at 24° C for the *Z* to *E* equilibration is given in Table 3. Examination of the numbers in Tables 1 and 2

Table 3: *Z* to *E* Equilibration of *t*-Butyl Formate **5** at 24° C

Solvent	% <i>E</i> Isomer	K_{eq}	ΔG°_{297} (kcal/mole)
DMSO-d ₆	32.9	0.49	0.42
CD ₃ CN	27.4	0.38	0.58
DMF-d ₇	26.4	0.36	0.61
Acetone-d ₆	21.0	0.27	0.78
Toluene-d ₈	12.3	0.14	1.17

for various solvents shows that, in general, the percentage of the minor *E* isomer first increases and then decreases with increasing temperature. The initial increase is readily explained since one would normally expect that the percentage of the minor isomer of an equilibrium would be increased at higher temperatures. However, here an additional effect is operative, namely that the polarity of solvents (especially very polar solvents) decreases with increasing temperature.¹³ Therefore at higher temperatures, the solvent is less polar, the stabilization of the polar *E* isomer is lessened and its percentage decreases. The crossover point for these two opposing effects differs depending on the solvent but in nearly all cases an initial increase and then a decrease in the percentage of the *E*-isomer is seen. For aprotic solvents, then, the effect is clear: higher polarity causes a higher percentage of the *E* isomer.

However, at this time, we have no explanation for the very low percentages of the *E* isomer in the very polar protic solvents ethanol and NMF. With these solvents, one would expect other factors such as hydrogen bonding to

be important.¹⁴ However, recent theoretical calculations¹⁵ (Monte Carlo simulations on ester stability in protic solvents) show clearly that the *E* isomer would be expected to be stabilized to a greater degree by hydrogen bonding than the *Z* isomer and thus one would predict more, not less, of the more polar *E* isomer in these solvents. Further studies, both experimental and theoretical, are underway to try to explain this discrepancy.

In summary, for the first time the $^3J_{\text{CH}}$ of the *Z* and *E* conformers of *t*-butyl formate **5** have been determined and used to calculate the isomeric composition of **5** in various solvents at different temperatures. Polar aprotic solvents cause an increase in the percentage of the minor more polar *E* isomer to a maximum of 33% in DMSO at 24° C.

Acknowledgement: We thank the National Institutes of Health (GM-31349) for financial support and Frank Anet, Ken Houk, Jane Strouse, Nazim Jaffer, Dan O'Leary, Jeff Evanseck, and Julie Bartley, all at UCLA, for helpful advice and discussions. We thank Prof. Jay Siegel of UC San Diego for pointing out an earlier use of $^3J_{\text{CH}}$ for the conformational analysis of acetaldehyde and its oxime *O*-methyl ether by Karabatsos and coworkers.¹⁶

References and Notes

1. Winstein Dissertation Awardee, UCLA 1989.
2. Jung, M. E.; Gervay, J. *J. Am. Chem. Soc.* **1989**, *111*, 5469.
3. For reviews, see: a) Exner, O. In *The Chemistry of Double-Bonded Functional Groups*; Patai, S., Ed.; Interscience: London, 1977; p 1. b) Jones, G. I. L.; Owen, N. L. *J. Mol. Struct.* **1973**, *18*, 1.
4. Wiberg, K. B.; Laidig, K. E. *J. Am. Chem. Soc.* **1987**, *109*, 5935.
5. Larson, J. R.; Epiotis, N. D.; Bernardi, F. *J. Am. Chem. Soc.* **1978**, *100*, 5713.
6. Epiotis, N. D.; Cherry, W. R.; Shaik, S.; Yates, R. L.; Bernardi, F. *Top. Curr. Chem.* **1977**, *70*, 1. b) Mark, H.; Baker, T.; Noe, E. A. *J. Am. Chem. Soc.* **1989**, *111*, 6551.
7. Ōki, M.; Nakanishi, H. *Bull. Chem. Soc. Jpn.* **1970**, *43*, 2558.
8. a) Grindley, T. B. *Tetrahedron Lett.* **1982**, *23*, 1757. b) Nakanishi, H.; Fujita, H.; Yamamoto, O. *Bull. Chem. Soc. Jpn.* **1978**, *51*, 214.
9. Drakenberg, T.; Forsén, S. *J. Phys. Chem.* **1972**, *76*, 3582.
10. We assume here that the coupling constant for both isomers is both solvent- and temperature-independent. Our measured values for J_{trans} in several solvents at -100 to -121° C was more sensitive to the shimming of the NMR instrument than anything else and varied less than 10% in all experiments, so that we think these assumptions are fair. The usual *J* averaging method was used: $J_{\text{obs}} = J_{\text{E}} \rho_{\text{E}} + J_{\text{Z}} \rho_{\text{Z}}$, where ρ_{x} = mole fraction of *x* isomer.
11. Special precautions had to be taken to reach these temperatures. It was necessary to rigorously exclude water, to use new, high quality NMR tubes to avoid complications from uneven spinning or scratches on the inside wall of the tube, and to very gradually reduce the temperatures below -70° C.
12. a) Reichardt, C. *Angew. Chem. Int. Ed. Eng.* **1965**, *4*, 29. b) Reichardt, C. *Solvents and Solvents Effects in Organic Chemistry*, 2nd Ed.; Verlag: Weinheim, 1988.
13. a) McClellan, A. L. *Tables of Experimental Dipole Moments*; Freeman: San Francisco, 1963. b) Carvajal, C.; Tölle, K. J.; Smid, J.; Szwarc, M. *J. Am. Chem. Soc.* **1965**, *87*, 5548.
14. One factor affecting this discrepancy is the fact that in a hydrogen bonding solvent the $^3J_{\text{CH}}$ may change significantly since the effective hybridization of the carbonyl carbon is altered. Therefore the application of the *J* averaging method to the results in NMF, for example, would have to be modified to utilize the actual coupling constants, which we cannot measure in this system due to the freezing point of NMF (-40° C).
15. Houk, K. N.; Evanseck, J. D. private communication.
16. Karabatsos, G. J.; Orzech, C. E., Jr.; Hsi, N. *J. Am. Chem. Soc.* **1966**, *88*, 1817.

(Received in USA 23 May 1990)