

Pelter A.	1657	Scola P. M.	1557
Pestellini V.	1637	Sharma H. A.	1593
Pohlmann K.	1597	Sharma M.	1641
Pokkuluri P. R.	1535	Shimizu R.	1619
Pramanik B. N.	1641	Shingu T.	1621
Quéguiner G.	1611	Solladié G.	1605
Reina M.	1661	Sonawane H. R.	1645
Rodier F.	1609	Spanton S.	1573
Rodriguez M. L.	1661	Takahashi C.	1621
Rookhuizen R. B.	1633	Tanaka J.	1593
Sakai M.	1651	Teng K.	1539
Sako Y.	1651	Theobald F.	1609
Sasaki O.	1651	Trotter J.	1535
Scheffer J. R.	1535	Tsunoda T.	1651
Schmidt R. R.	1585	Usami Y.	1621
Schmitt G.	1609	Williams D. E.	1561
Scola P. M.	1549	Yoon S. K.	1553
Scola P. M.	1553	Yoshida Z.	1619

CORRIGENDUM

G. I. Georg, P. M. Mashava, E. Akgün and M. Milstead, Asymmetric synthesis of β -lactams and *N*-benzoyl-3-phenylisoserines via the Staudinger reaction, *Tetrahedron Letters*. **1991**, 32, 3151-3154

A reevaluation of our preliminary results on the Staudinger reaction of the benzaldimine derived from 2,3,4,6-tetra-*O*-acetyl- β -D-galactosamine with *p*-methoxyphenoxyacetyl chloride revealed that we had inadvertently overlooked the presence of the other *cis*-isomer. The reaction typically produces a 60:40 ratio of the two possible *cis*-isomers in 75% yield, with isomer **7** as the minor product. A more detailed account of this work is in press in *Tetrahedron Lett.*