

Seto M.	345	Takaishi Y.	339
Shankar R.	231	Theil F.	305
Shibuya M.	339	Thierry J.	323
Shirahama H.	331	Toske S. G.	207
Shishido K.	339	Truong V.	227
Shiue J.-S.	335	Turnbull K.	239
Siegel J. S.	273	Turner W. R.	281
Simpkins N. S.	363	Wan W.-Q.	261
Sims J. J.	223	Watson P. S.	199
Singaram B.	257	Wegmann S.	307
Singh J.	211	Weidner J.	305
Smith M. J.	223	Werblood H. M.	203
Sprules T. J.	247	White J. D.	207
Stangeland E. L.	257	Wild H.	285
Stayton M. M.	223	Würthwein E.-U.	307
Stille J. R.	215	Xu D.	227
Suto M. J.	281	Yue C.	323
Suzuki K.	341		

## CORRIGENDUM

M<sup>a</sup> I. Matheu, R. Echarri and S. Castillón, Stereoselective synthesis of nucleosides by metallocene-promoted activation of glycosyl fluorides, *Tetrahedron Letters*, **1992**, *33*, 1093.

On page 1094 line 7 should read:

using  $\text{Cp}_2\text{HfCl}_2\text{-AgOTf}$  in benzene as the solvent, giving the  $\alpha$  isomer with a high stereoselectivity, which