

## A Convenient Synthesis of Dibenzo[a,c]cyclooctene Based on Thioenol Ether Reduction


Pierre-André Lottaz, Timothy R.G. Edwards, Yves G. Mentha<sup>1</sup> and Ulrich Burger\*

Département de chimie organique, Université de Genève,  
CH- 1211 Genève-4, Switzerland

**Key Words:** Intramolecular reductive coupling; desulfurization of thioenol ethers; axial chirality; carbene addition


**Abstract:** 9,10,11,12-Tetrahydrodibenzo[a,c]cyclooctene-9,12-diol (**6**) is found to be reluctant to double bond forming elimination. It gives a bridged ether (**7**) instead. The corresponding diketone (**5**) reacts with benzyl mercaptan to provide the bis-thioenol ether (**8-a**). The latter, upon treatment with Raney nickel, gives the desired title diene **1**. An alternative approach to **1**, based on repeated carbene addition to phenanthrene, is also presented.

The efficient synthesis and the properties of *o-o'*-bridged biphenyls continue to be of interest to many areas of chemistry<sup>2,3</sup>. The seemingly simple title hydrocarbon **1**, although known for nearly three decades<sup>4</sup>, has only recently been resolved into its atropisomers (R)-**1** and (S)-**1**<sup>5</sup>. The chiral features of these antipodes, which are protected against racemization by an inversion barrier of  $\Delta G^\ddagger = 30$  kcal/mol, have not yet been exploited. In connection with our continuing studies of highly strained bicyclobutanes<sup>6</sup> we have demonstrated that the diene **1** undergoes a photo-Diels-Alder reaction with 4-methyl-3H-1,2,4-triazol-3,5(4H)-dione (MTAD). Thermally, **1** is inert towards the dienophile. Further studies on these lines are greatly hampered by the inefficiency of preparations of **1** according to literature procedures<sup>4a,7b</sup>.



One of the first syntheses reported for **1** starts from the photoadduct of maleic anhydride and phenanthrene<sup>4a</sup>. It implies an oxidative bis-decarboxylation and a disallowed thermal ring opening. The method has been extended to derivatives of both maleic anhydride and phenanthrene<sup>2a,8</sup>, but global yields are invariably low (1-6%). A somewhat better method was described by Wong and Sondheimer<sup>7b</sup>. It starts from 2,2'-bis(bromomethyl)-1,1'-biphenyl and avoids the disallowed ring opening. This method also has been extended to derivatives of the parent system<sup>2a</sup>. Global yields are in the range of 5-9%. In the present communication, we wish to report on a five-step synthesis of **1** which starts from the commercially available 2-aminoacetophenone **2**. The eight-membered ring is set up by intramolecular reductive coupling<sup>7a</sup>. The target diene **1** is obtained by this synthesis in 21% overall yield including the separation of the enantiomers (+)-**1** and (-)-**1**. Brief mention will be made at the end of this communication of an alternative approach towards **1**, which is based on ring expansion of phenanthrene *via* repeated carbene addition. It gives **1** in 14-16% yield.


For the initial biaryl formation (2 → 3) we have chosen reductive coupling of the diazonium salt derived from 2. The method had been shown to be very efficient in the analogous synthesis of diphenic acid<sup>9</sup> and proceeded in higher yield (96%) than the classical Ullmann reaction performed with the corresponding iodide (59%)<sup>10</sup>. The subsequent double bromination with phenyltrimethylammonium perbromide (PTAB) (3 → 4) as well as the zinc-copper promoted reductive cyclization providing the diketone 5, are known reactions<sup>10</sup>. The ring closure, with its modest 48% yield, is the weakest step in our synthesis. Our efforts to convert the diketone 3 directly into 5 by intramolecular oxidative coupling following Kobayashi's procedure<sup>11</sup> met with limited success. This procedure gave 5 in 26% yield only, and therefore does not present any advantage over the path *via* the dibromide 4.




Initially, we had planned to reduce the diketone 5 into its diol 6 (mixture of stereomers) and to reach the target diene 1 by standard elimination procedures. As expected, the reduction step (5 → 6) did not provide any problems. The difficulties started when the final elimination (6 → 1) was attempted. Most of the classical methods (*e.g.*  $\text{TsOH/benzene}$ ) when applied to the diol 6, gave the bridged ether 7 as the main product<sup>12,13</sup>.

We could not find methods transforming 6 or 7 in acceptable yield into the desired diene 1. In order to overcome these difficulties, we resorted to a desulfurization method which had been used successfully in steroid chemistry<sup>14</sup>. The diketone 5 was found to react with two equivalents of benzyl mercaptan in the presence of  $\text{BF}_3$  as a catalyst, to give the double thioenol ether 8-a in 88% yield<sup>12,15</sup>. Thiophenol reacts in

analogous fashion to give **8-b** (86% yield)<sup>12</sup>. Compounds **8-a** or **8-b** were readily desulfurized by treatment with Raney nickel in acetone to give the desired diene **1** in yields of 60% and 58% respectively<sup>16</sup>. This last step gives as a by-product the known 9,10-dihydro-dibenzo[*a,c*]cyclooctene **9**<sup>5</sup> in 15-20% yield. Careful tlc monitoring of the disappearance of starting material is required during the desulfurization in order to minimize this undesired overreduction. Compound **1** was separated into its antipodes by medium pressure chromatography on swollen microcrystalline triacetylcellulose (TAC)<sup>5</sup> with methanol/water (95/5) as eluent. This final chromatographic step results also in the separation from **9**. The compounds are eluted in the order (+)-**9**, (+)-**1**, (-)-**9**, and (-)-**1**. It must be mentioned however that compounds **1** and **9** have very similar retention times on ordinary silicagel columns. If one aims only at the isolation of racemic **1**, it is advantageous to first reflux the crude **1/9** mixture in *p*-xylene for 5h with selenium dioxide. This leaves **1** unchanged but converts **9** into the ketone **10**<sup>12</sup>. Final medium-pressure chromatography on silica gel with hexane/ether (95/5) as eluent gives pure **1**.



In addition to the synthesis described above, we have examined an entirely different approach towards **1** and wish to report briefly on that finding. Several laboratories<sup>2a,17</sup> had shown that phenanthrene can be readily expanded into 5H-dibenzo[*a,c*]cycloheptene **11**. The three step process consists of the addition of a dihalocarbene to phenanthrene with subsequent cyclopropyl-allyl rearrangement and reduction. Reported yields for **11** range from 45-60%. We found that the dibromocarbene adduct of **11** undergoes selective reduction with LiAlH<sub>4</sub> (Et<sub>2</sub>O/0°C/48h) to give the *endo*-bromocyclopropane derivative **12**<sup>12,18</sup> in 47% yield. It is accompanied by the corresponding fully reduced cyclopropane derivative (14%). Heating of **12** in dry pyridine in a sealed Pyrex tube (185°C, 15h) results in the formation of **1** in 70% yield based on **12**. Note that a pyridinium salt resulting from a substitution path is obtained if this reaction is run at too low temperature (<175°C). This carbene route towards **1** is clearly a viable alternative to the other approaches.



a) CHBr<sub>2</sub>/t-BuOK/t-BuOH/pentane/-10°C/5h.  
 b) LiAlH<sub>4</sub>/Et<sub>2</sub>O/0°C/48h. c) Pyridine/185°C/14h.

**Acknowledgments.** The authors wish to express their gratitude to Mrs. P. Millasson for technical help. Financial support was provided by the Swiss National Science Foundation (grant no. 20-32645.91).

## References and notes

1. New address: Firmenich S.A. Genève
  2. a) Müllen,K.; Heinz,W.; Klärner,F.G.; Roth,W.R.; Kindermann,I.; Adamczak,O.; Wette,M.; Lex,J. *Chem.Ber.* **1990**, *123*, 2349. b) Rashidi-Ranjbar,P.; Sandström,J.; Wong,H.N.C.; Wang,X.C. *J.Chem.Soc. Perkin Trans. 2*, **1992**, 1625.
  3. a) Bringmann,G.; Walter,R.; Weirich,R. *Angew.Chem.* **1990**, *102*, 1006; *Angew.Chem. Int.Ed. Engl.* **1990**, *29*, 977. b) Warshawsky,A.M.; Meyers,A.I. *J.Am.Chem.Soc.* **1990**, *112*, 8090. c) Pelter,A.; Ward,R.S.; Jones,D.M.; Maddocks,P. *Tetrahedron Asymm.* **1992**, *3*, 239. d) Becker,D.; Hughes,L.R.; Raphael,R.A. *J.Chem.Soc. Perkin Trans. 1*, **1977**, 1674. e) Mervic,M.; Ben-David,Y.; Ghera,E. *Tetrahedron Lett.* **1981**, *22*, 5091.
  4. a) Vogel,E.; Frass,W.; Wolpers,J. *Angew. Chem.* **1963**, *75*, 979; *Angew.Chem. Int. Ed. Engl.* **1963**, *2*, 625. b) X-ray structure: Huang,N.Z.; Mak,T.C.W.; *J.Mol. Structure* **1983**, *101*,135. c) detailed <sup>1</sup>H-NMR analysis: Günther,H.; Günther,M.E.; Mondeshka,D.; Schmickler,H. *Liebigs Ann. Chem.* **1978**, 165. d) dianion: Müllen,K. *Helv.Chim.Acta* **1978**, *61*, 1296.
  5. Rashidi-Ranjbar,P.; Sandström,J. *Tetrahedron Lett.* **1987**, *28*, 1537.
  6. Burger,U.; Mentha,Y.G.; Millasson,P.; Lottaz,P.A.; Mareda,J. *Helv.Chim.Acta* **1989**, *72*, 1722.
- 
8. a) Willner,I.; Rabinovitz,M. *J. Org. Chem.* **1980**, *45*, 1628 b) Willner,I.; Rabinovitz,M. *Tetrahedron Lett.* **1976**, 1223.
  9. Tietze,L.F.; Eicher,Th. *Reaktionen und Synthesen*, Thieme Verlag Stuttgart, **1981**, 197; English translation: *Reactions and Syntheses*, University Science Books, Mill Valley, California, **1989**, 208..
  10. Ghera,E.; Gaoni,Y.; Shoua,S. *J.Am.Chem.Soc.* **1976**, *98*, 3627.
  11. Kobayashi,Y.; Taguchi,T.; Morikawa,T.; Tokuno,E.; Sekiguchi,S. *Chem. Pharm. Bull.* **1980**, *28*, 262.
  12. All new compounds gave correct elemental analyses and mass, IR, <sup>1</sup>H-NMR, and <sup>13</sup>C-NMR spectra consistent with the structures indicated.
  13. Compd 7: colourless oil, <sup>1</sup>H-NMR(200MHz,CDCl<sub>3</sub>):1.97-2.30(m,4H);5.20-5.26(m,2H);7.11-7.81(m,8H).
  14. Boar,R.B.; Hawkins,D.W.; McGhie,J.F.; Barton,D.H.R. *J.Chem. Soc. Perkin Trans. 1*, **1973**, 654.
  15. Compd 8-a: colourless crystals. m.p.97-98°C. <sup>1</sup>H-NMR(400MHz,CDCl<sub>3</sub>): 3.61(s,4H); 5.97(s,2H); 7.0-7.5(m,18H).
  16. A slurry of Raney nickel (ca. 1g; Fluka 83440) was prewashed 3x with acetone to remove water. The slurry with 30 ml acetone was deactivated by boiling under reflux for 2h. 8-a (269mg, 0.6 mmole) in 15 ml acetone was added to the cold mixture and brought to reflux for 16h. The mixture was filtered over Celite and the solvent withdrawn under vacuum. The remaining mixture 1/9 was separated on TAC as indicated in the text.
  17. a) Christl,M. University of Würzburg, Germany, private communication. b) Nusser,R. Dissertation, University of Würzburg, Germany, 1990. c) cf. Wan,P.; Budac,D.; Earle,M.; Shukla,D. *J.Amer.Chem.Soc.* **1990**, *112*, 8048.
  18. Compd. 12: colourless crystals, m.p. 144-146°C. <sup>1</sup>H-NMR (200 MHz, CDCl<sub>3</sub>): 1.88(dddd, J=10.8, 9.6, 7.3, 6.0, 1H); 2.21(dd, J=9.6, 7.8, 1H); 2.72(dd, J=13.5, 10.8, 1H); 2.84(dd, J=13.5, 6.0, 1H); 3.39(dd, J=7.8, 7.3, 1H); 7.2-7.6(m, 8H<sub>arom.</sub>).