

## Villagorgin A and B. New Type of Indole Alkaloids with Acetylcholine Antagonist Activity from the Gorgonian *Villagorgia rubra*

Alfonso Espada,<sup>1</sup> Carlos Jiménez,<sup>2</sup> Cecile Debitus,<sup>3</sup> and Ricardo Riguera<sup>1\*</sup>

<sup>1</sup>Depart. de Química Orgánica, Fac. de Química, Universidad de Santiago, Santiago de Compostela. 15706. Spain;

<sup>2</sup>Depart. de Química Fundamental e Industrial, Fac. de Ciencias, Univ. da Coruña, A Coruña. 15071, Spain ;

<sup>3</sup>ORSTOM, Centre de Noumea, B. P. A5, Noumea, New Caledonia.

**Abstract:** Two new indoloquinolizidine alkaloids incorporating an imidazol ring, villagorgin A and villagorgin B, along with caffeine, tryptamine, *Nb*-methyltryptamine, and 1, 2, 3, 4 tetrahydro- $\beta$ -carboline, were isolated from the gorgonian *Villagorgia rubra*, and determined by spectral data. Villagorgin A has a calmodulin-related antagonist activity.

Gorgonians have been shown to contain a fair variety of terpene metabolites<sup>1</sup> mainly diterpenes (cembranoids), sesquiterpenes and steroids. *Paramuricea chamaeleon* represents the only case of a gorgonian rich in indole alkaloids<sup>2</sup>. In connection with our investigations of biologically active marine metabolites, we focused our attention on the gorgonian *Villagorgia rubra*, selected for study because the methanol extract was very active in the guinea-pig ileum contraction test. In this paper we report that *V. rubra* is a rich source of nitrogenous metabolites that were isolated from the butanolic fraction and identified as caffeine (1), the simple indoles, tryptamine (2), *Nb*-methyltryptamine (3), 1, 2, 3, 4 tetrahydro- $\beta$ -carboline (4), and two new complex indole alkaloids named as villagorgin A (5) and villagorgin B (6). The structure and absolute configuration of villagorgins were deduced by extensive use of 2D-NMR, FABMS and HREIMS, and CD data.

The gorgonian *Villagorgia rubra*, a genus which has never been studied before, was collected in New Caledonia. The lyophilized material gave a methanol extract which was then partitioned into n-hexane, CH<sub>2</sub>Cl<sub>2</sub> and n-BuOH. The n-butanol was passed through amberlite XAD-2 and the methanolic eluates (1 g) fractionated on a Sephadex LH-20 column eluted with methanol to produce eight fractions. Fractions 7-8 contained compounds 5 and 6 that were isolated by reversed phase HPLC (MeOH-OH<sub>2</sub> 60:40; r.t = 14 and 24 min., respectively). Fractions 4-6 contained known compounds 1-4 isolated by HPLC as before.

Villagorgin A (5, 12 mg), was obtained as a yellow powder [ $\alpha$ ]<sub>D</sub><sup>20</sup> = +7.8°. The molecular formula C<sub>16</sub>H<sub>16</sub>N<sub>4</sub>, determined by HREIMS ([M<sup>+</sup>] = 264.1378,  $\Delta$  0.3 mmu of calcd) inferred 11 degrees of unsaturation. The presence of six quaternary, six methine and four methylene carbon signals was deduced from the <sup>13</sup>C and DEPT NMR spectra. The HMQC experiment allowed us to assign and correlate all the protons to the corresponding carbons.

The UV spectra of 5 ( $\lambda$  (nm) 224 ( $\epsilon$  25600) and 280 ( $\epsilon$  52409))<sup>3</sup> and <sup>1</sup>H NMR signals in the 7.5-6.9 ppm region revealed the presence of an indole moiety in the molecule.

The proton and carbon chemical shifts for the C-5 and C-6 methylenes and the C-3 methine indicated that the indole system constitutes a part of a tetrahydro  $\beta$ -carboline moiety<sup>4a</sup>. The  $m/z$  169 (100%) ion in the EIMS of **5** is characteristic of this type of structure.<sup>4b</sup>

The <sup>1</sup>H-<sup>1</sup>H COSY NMR spectrum showed a coupling between the highly deshielded H-3 methine proton at  $\delta$  3.83 ppm and the H-14 methylene protons at  $\delta$  3.36 and 2.72 ppm. Relay-COSY displayed five bond correlations between H-14 and both H-20 methylene protons at  $\delta$  3.99 and 3.62 ppm. Furthermore, HMBC experiments showed that the olefinic quaternary carbons C-15 and C-19 are correlated with H-14 and H-20. All these data indicate that **5** has the indoloquinolizidine system characteristic of well known plant alkaloids like yohimbine and corynantheine.<sup>5</sup>

The remaining two unsaturations and the CHN<sub>2</sub> atoms were assumed to comprise an imidazol<sup>6</sup> system on the basis of the carbon signal at  $\delta$  135.2 (C-17, d,  $J_{CH} = 207.5$  Hz) bonded to the proton at  $\delta$  7.57 ppm. HMBC correlations and particularly those between C-19 and H-17 corroborated the D/E ring connections and the structure of villagorgin A as **5**.<sup>7</sup>

The trans quinolizidine stereochemistry was determined by the strong ROESY correlation between H-3 and the axial proton H-20. Furthermore, the circular dichroism curve of **5** presents a negative Cotton effect at 277 nm, ( $\Delta\epsilon$  -0.19) and this allowed us to determine the 3 $\beta$ -H absolute stereochemistry<sup>8</sup> for villagorgin A (**5**).

Villagorgin B (**6**, 2mg), has molecular formula C<sub>16</sub>H<sub>13</sub>N<sub>4</sub>. The positive ion FABMS showed the molecular ion M<sup>+</sup> at  $m/z$  261. Its uv spectrum showed absorbances due to an indole system and an additional chromophore.<sup>9</sup>



The <sup>13</sup>C and <sup>1</sup>H-NMR data of compound **6** indicate the presence of the indole system. The 5, 6 dihydro  $\beta$ -carboline system with a positively charged nitrogen atom was deduced from the chemical shifts of the C-6 and C-5 ethylene group<sup>10</sup>. In accordance with that, the positive ion FABMS of villagorgin B gives a prominent ion  $m/z$  171 (100 %).

In addition, the <sup>13</sup>C and <sup>1</sup>H-NMR spectra of **6** displayed signals for three isolated aromatic methines (C-14, C-17 and C-20). Five bond correlations were found among those protons by RELAY COSY experiments. HMBC spectra displayed the correlations showed in Table 1. Particularly relevant are those of C-17 to H-14, C-20 to H-17, and C-3 to H-20. All these data suggest an imidazo [4, 5-c] pyridine system for the D/E rings. The ions at  $m/z$  259 (M<sup>+</sup>-2, fully aromatic structure),  $m/z$  235 (M<sup>+</sup> -CN), and 207 (M<sup>+</sup> -2 HCN) in the (+) FABMS confirms the presence of the imidazol group and structure **6** for villagorgin B. The small amount available and its fast decomposition impeded the realization of further experiments.

The known alkaloids caffeine (**1**, 4 mg), tryptamine (**2**, 5 mg), *Nb*-methyltryptamine (**3**, 4 mg), and 1, 2, 3, 4 tetrahydro- $\beta$ -carboline (**4**, 2 mg) were also isolated from *V. rubra* and identified by comparison with published data and authentic samples.

Villagorgin A (**5**) was shown to produce strong inhibition on the acetylcholine induced contraction of guinea-pig ileum and a dose-dependent inhibitory effect against human platelet aggregation induced by thrombin and calcium ionophore A23187. Since this biological process is a calcium-calmodulin mediated event, the antiaggregatory activity of **5** could be due to the inhibition of that enzyme. It should be mentioned that villagorgin A and B are structurally related to the  $\beta$ -carboline marine alkaloid eudistomidin-A, isolated from a tunicate, and a strong calmodulin antagonist.<sup>11</sup>

Indole alkaloids have been isolated in a fair number from marine sources and some of them present outstanding pharmacological activity, but this is the first report of complex indoles in gorgonians.



Pos	$\delta$ H, m, J (Hz) (500 MHz)	Relay-COSY	$\delta$ C, m (60 MHz)	HMBC (JC-H = 7 Hz)
NH	10.98 s <sup>a</sup>			
2			128.4 s	H-6, H-14
3			142.4 s	H-14, H-20
5	4.86 t (7.0)		57.3 t	H-6
6	3.35 t (7.0)		21.2 t	
14	8.05 s	H-17, H-20	107.1 d	
15			164.4 s	H-14, H-20
17	8.93 s	H-14, H-20	135.2 d	H-14
19			155.6 s	H-20, H-17
20	8.38 s	H-14, H-17	163.5 d	H-17

Pos	$\delta$ H, m, J (Hz) (500 MHz) <sup>a</sup>	Relay-COSY	$\delta$ C, m (60 MHz)	HMBC (JC-H = 7 Hz)
NH	10.86 s <sup>a</sup>			
2			135.1 s	H-6 <sup>b</sup>
3	3.83 dt, (3.7, 11.0)	H-14, H-14'	58.4 d	H-14 <sup>b</sup> , H-14', H-5', 5'', H-20'
5	3.33 m	H-5', H-6, H-6'	54.0 t	H-6, 6', H-20'
5'	2.87 add (4.0, 12.0, 22.5)	H-5, H-6, H-6'		
6	3.00 m	H-5, H-5', H-6'	22.3 t	H-5, 5 <sup>b</sup>
6'	2.79 dt, (2.0, 16.0)	H-5, H-5', H-6		
14	3.36 m	H-3, H-14', H-20'	29.2 t	
14'	2.72 m	H-3, H-14, H-20		
15			129.7 s	H-14, H-20, 20'
17	7.57 s		135.2 d	
19			127.3 s	H-17 <sup>b</sup> , H-20, 20'
20	3.99 d, (13.0)	H-14', H-20'	53.4 t	H-3
20'	3.62 dt, (13.0)	H-14, H-14', H-20		

Table 1. Selected NMR data of villagorgin A (5) and villagorgin B (6) in CD<sub>3</sub>OD; a) In DMSO-d<sub>6</sub> b) HMBC with a J<sub>C-H</sub> = 3.5 Hz.

Indole alkaloids containing an imidazol ring are known from either marine or terrestrial origin, but compounds **5** and **6** are the first examples with the imidazol ring attached to an indoloquinolizidine and constitute a new skeleton among this type of alkaloids. A biogenetic route to **5** and **6** can be formulated from tryptophane and histidine as starting aminoacids, as in some *Penicillium* alkaloids such as oxaline and roquefortine<sup>12</sup>, but requires an additional carbon unit (C-20). The isolation of caffeine (**1**) from *Villagorgia rubra*, is noteworthy because it has been considered to be a terrestrial metabolite. This is, as far as we know, the first report on its presence in a marine organism.

**Acknowledgments:** This work was supported by Grants SAF 1023/92 (CICYT), XUGA-20906B91 (Xunta de Galicia) and 60902.25060 (Univ. da Coruña) . A.E. acknowledges fellowship from the Programa de Cooperación con Iberoamérica. We are grateful to Profs. Jesus Trujillo for the CD spectra, M. Feliz for some NMR spectra and Ernesto Cano for the pharmacological assays.

#### REFERENCES AND NOTES

1. a) Faulkner, D. J. *Tetrahedron* **1977**, *33*, 1421. b) Faulkner, D. J. *Nat. Prod. Rep.*, **1992**, *9*, 323.
2. Cimino, G. and De Stefano, S. *Comp. Biochem. Physiol.* **1978** *C61*, 361.
3. Verpoorte, R. *J. Nat. Prod.* **1986**, *49*, 1.
4. a) Debitus, C.; Laurent, D.; País, M. *J. Nat. Prod.* **1988**, *51*, 799. b) Besselièvre, R.; Cosson, J. P.; Das, B. C. and Husson, H. P. *Tetrahedron Lett.* **1980**, *21*, 63.
5. a) Jokela, R. *Heterocycles* **1992**, *30*, 1949. b) Broadbent, T.A. and Paul, E. G. *Heterocycles* **1983**, *20*, 863.
6. Grimmet M. R. : Imidazoles and the Benzo Derivatives: (i) Structure. In *Comprehensive Heterocyclic Chemistry*, Katritzky, A. R and Rees C. W. Eds., Pergamon Press. Oxford. 1984; Vol. 5, pp.345-372.
7. **5**: Red amorphous solid. IR  $\nu_{\max}$ : 3880, 2923, 1650, 1541  $\text{cm}^{-1}$ . <sup>1</sup>H-NMR (CD<sub>3</sub>OD)  $\delta$  (m, J in Hz): 7.41 (d, 7.5, H-9), 7.30 (d, 7.5, H-12), 7.06 (ddd, 1.0, 7.5, 16.0, H-11), 6.98 (ddd, 0.5, 7.5, 14.5, H-10) and Table 1. <sup>13</sup>C-NMR (CD<sub>3</sub>OD): 138.5 (C-13), 128.1 (C-8), 122.3 (C-11), 118.8 (C-9), 111.9 (C-10), 112.0 (C-12), 108.5 (C-7) and Table 1. HREIMS: C<sub>16</sub>H<sub>16</sub>N<sub>4</sub> 264.1375: calc. 264.1378. EIMS m/z (%): 264 (55), 234 (10), 220 (6), 207 (4), 169 (100), 154 (6), 142 (9), 129 (5), 115 (11), 94 (8). (-) FABMS, m/z, (%): 263 ([M-H]<sup>+</sup>, 100).
8. Barlett, L.; Dastoor, N. J.; Hrbek, J.; Klyne, W.; Schmid, H.; Snatzke, G. *Helv. Chim. Acta* **1971**, *54*, 1238.
9. **6**: Red amorphous solid. U.V. (MeOH)  $\lambda_{\max}$ : 228, 244, 280, 338 nm. UV (MeOH, HCl 1N)  $\lambda_{\max}$ : 222, 242, 332 nm. IR  $\nu_{\max}$ : 3480, 1648, 1632  $\text{cm}^{-1}$ . <sup>1</sup>H-NMR (CD<sub>3</sub>OD)  $\delta$  (m, J in Hz): 7.61 (d, 8, H-9), 7.46 (d, 8.5, H-12), 7.26 (ddd, 1.5, 7.5, 15, H-10), 7.12 ddd (1, 8, 15, H-10) and Table 1. <sup>13</sup>C-NMR (CD<sub>3</sub>OD): 139.9 (C-13), 126.9 (C-8), 125.4 (C-11), 121.4 (C-10), 120.3 (C-9), 113.8 (C-7), 112.5 (C-12) and Table 1. (+) FABMS (Glycerol matrix), m/z (%): 261 (M<sup>+</sup>, C<sub>16</sub>H<sub>16</sub>N<sub>4</sub>, 50), 205 (60), 171 (100), 135 (80); (+) FABMS (Magic +NaCl matrix), m/z, (%): 261 (M<sup>+</sup>, C<sub>16</sub>H<sub>16</sub>N<sub>4</sub>, 20), 235 (M<sup>+</sup>-CN, 22), 215 (37), 207 (M<sup>+</sup>-2HCN, 15), 177 (53), 115 (100).
10. Giri, V. S., Maiti, B. C., Pakrashi, S. C. *Heterocycles* **1984**, *22*, 233.
11. Kobayashi, J.; Nakamura, H.; Ohizumi, Y. *Tetrahedron Lett.* **1986**, *27*, 1191.
12. Vlegaar, R.; Wessels, P. L. *J. Chem. Soc. Chem. Comm.* **1980**, 160.