

S0040-4039(96)00172-4


Enantioselective Synthesis of the C1-C9 Segment of Bryostatin by Kinetic Resolution of Racemic β -Keto Esters

Markus Kalesse* and Marcus Eh

Institut für Organische Chemie der Universität Hannover, Schneiderberg 1b, 30167 Hannover, Germany

Abstract: The enantioselective synthesis of the C1-C9 segment of bryostatins is described. Racemic β -keto ester **2** was subjected to kinetic resolution. Reduction with baker's yeast establishes two of the three chiral centers. Chemical transformation of the terminal acetylene moiety generates aldehyde **5** which is transformed diastereoselectively to the corresponding alcohol **6** via Sakurai reaction.

The bryostatins are a family of 17 macrolides antibiotics isolated from the marine bryozoan *Bengula neritina* Linnaeus, which exhibit exceptional antineoplastic activity against lymphocytic leukemia and ovarian carcinoma.¹ The first total synthesis was reported by Masamune² in 1990 and since then various groups have made contributions to the synthesis of bryostatins.³ We would like to thank Prof. H. M. R. Hoffmann and J. Weiß for bringing this problem to our attention.⁴




bryostatins

R=H; OAc;
 O₂C(CH)₄(CH₂)₂Me

Scheme 1

Here we report on the synthesis of the C1-C9 segment of bryostatin utilizing the kinetic resolution of racemic β -keto ester **2** with baker's yeast. Analysis of the backbone of the C1-C9 segment identifies the product derived from baker's yeast kinetic resolution to be identical with the desired stereochemistry of the natural product (Scheme 1). Baker's yeast reduction of **2** already establishes two asymmetric centers of the C1-C9 segment and allows functionalization of the terminal acetylene moiety. The overall strategy is to generate an


aldehyde at C7 which enables a chelation-controlled Sakurai reaction. This reaction establishes the third asymmetric center and additionally the geminal dimethyl group at C8.



a) Ethyl acetoacetate, 1 eq. NaH, 0°C, 1 eq. LDA, -78°C, THF, then 1, 83 %; b) C₆H₅CH₂OC(=NH)CCl₃, CH₂Cl₂, 77 %; c) KF, 18-crown-6, DMF, H₂O, 99 %; d) baker's yeast, H₂O, sucrose, 3 d, 38 %.


Scheme 2

Racemic β-keto ester **2** was generated by addition of ethyl acetoacetate to propargylic aldehyde **1**. Protection with benzyl trichloroacetimidate and desilylation established compound **2** suitable for baker's yeast reduction. Kinetic resolution of **2** yields **3** in very good enantiomeric and diastereomeric excess (de = 82 %; ee = 84 %) (Scheme 2).⁵ Even though the kinetic resolution of **2** did not yield the excellent ee values observed for the kinetic resolution of **9** (Scheme 3),⁶ we decided to use **3** as the precursor due to the greater stability of the benzyl group under Sakurai conditions.




Scheme 3

The diastereomeric ratio of **3** was determined by chiral GC analysis.⁷ The enantiomeric purity was established by NMR-shift experiments with Eu(hfc)₃⁸ as described by Meyer and Oetting.⁹ However, hydroboration of the triple bond did not yield the desired aldehyde but methyl ketone **8** instead (Scheme 4).



Scheme 4

We therefore hydrogenated the acetylene moiety to the corresponding double bond with Lindlar catalyst (70 %) and reduced the ester group with LiAlH₄. Double protection with TBDMS triflate¹⁰ generated the protected triol **4** in 43 yield (three steps). Hydroboration and successive oxidation with Dess-Martin periodinane¹¹ gave aldehyde **5**. The Sakurai reaction was performed under standard conditions¹² at -78°C in CH₂Cl₂ with TiCl₄ as the Lewis acid and gave tetraol **6**¹³ as a 6:1 mixture of two diastereomers with the desired trans diol as the major isomer (Scheme 5).



a) Lindlar catalyst, H₂, EtOH, 70 %; b) LiAlH₄, Et₂O, 67 %; c) 2 eq. TBDMS-triflate, THF, 92%; d) BH₃xTHF, THF, 41 %; e) Dess-Martin periodinane, 88 %; f) TiCl₄, CH₂Cl₂, 1-Trimethylsilyl-3-methyl-2-butene, 82 %.

Scheme 5

Acknowledgment: M.E. gratefully acknowledges a stipend by the Graduierten Förderung.

References

1. (a) Pettit, G. R.; Day, J. F.; Wood, H. B. *Nature* **1970**, *227*, 962. (b) Pettit, G. R. *The Chemist* **1989**, 11.
2. (a) Masamune, S. *Chimica*, **1988**, *42*, 210. (b) Masamune, S. *Pure Appl. Chem.* **1988**, *60*, 1587. (c) Blanchette, M. A.; Malamas, M. S.; Nantz, M. H.; Roberts, J. C.; Somfai, P.; Whritenour, D. C.; Masamune, S.; Kageyama, M.; Tamura, T.; *J. Org. Chem.* **1989**, *54*, 2817. (d) Kageyama, M.; Tamura, T.; Nantz, M. H.; Roberts, J. C.; Somfai, P.; Whritenour, D. C.; Masamune, S.; *J. Am. Chem. Soc.* **1990**, *112*, 7407.
3. Review on bioactive marine macrolides including progress on the total synthesis of bryostatins: R. D. Norcross and I. Paterson; *Chem. Rev.* **1995**, *95*, 2041-2114.
4. Diplomarbeit J. Weiß, 1994, Universität Hannover; Diplomarbeit M. Schinner, 1995, Universität Hannover.
5. Sering, B.; Seebach, D. *Helv. Chim. Acta* **1977**, *60*, 1175. Baker's yeast reduction of compound 2: Baker's yeast (10 g) was suspended in 100 ml warm water (30°C) and sucrose (20 g) was added. From the time the yeast started fermentation, the mixture was stirred for an additional 45 min. Then β-keto ester 2 (0.52 g, 3.36 mmol) was added and the reaction mixture was stirred for 24 h. Another 10 g of baker's yeast were suspended in warm water containing sucrose (20 g) and added to the reaction mixture. β-Keto ester 2 (0.52 g, 3.36 mmol) was added to this slurry. The reaction was stirred gently for additional 2 days and then filtered through a plug of Celite (12 g). The aqueous layer was extracted with Et₂O (3 x 200 ml), the combined organic layers were dried over MgSO₄, concentrated, and purified by flash chromatography (hexane/EtOAc 3:1) to yield diol 3 (385 mg, 38%) and unchanged β-keto ester 2 (1.4 mmol, 440 mg, 43%).

- 3:** $[\alpha]_D^{20} = +36.7$ (c 1.0 in CHCl_3); $^1\text{H-NMR}$ (200 MHz, CDCl_3 , 20°C, TMS): δ 7.33 (s, 5H; CH); 4.82 (d, $^2\text{J}(\text{H,H}) = 11$ Hz, 1H; CH_2); 4.49 (d, $^2\text{J}(\text{H,H}) = 11$ Hz, 1H; CH_2); 4.47 - 4.28 (m, 2H; CH); 4.15 (q, $^3\text{J}(\text{H,H}) = 7$ Hz, 2H; CH_2); 3.30 (s, br., 1H; OH); 2.52 (d, $^4\text{J}(\text{H,H}) = 2$ Hz, 1H; CH); 2.57 - 2.40 (m, 2H; CH_2); 1.92 (dd, $^3\text{J}(\text{H,H}) = 7$ Hz, $^3\text{J}(\text{H,H}) = 6$ Hz, 2H; CH_2); 1.25 (tr, $^3\text{J}(\text{H,H}) = 7$ Hz, 3H; CH_3); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3 , 20°C, TMS): 172.34 (+), 137.41 (+), 128.08 (-), 127.78 (-), 82.21 (+), 74.46 (+), 70.95 (+), 65.86 (-), 64.67 (-), 60.67 (+), 42.01 (+), 41.39 (+), 14.16 (-); IR (CHCl_3): $\nu = 3520$ (w), 3304 (m), 1720 (s), 1496 (m), 1452 (m), 1256 (m).
6. Eh, M.; Kalesse, M. *Synlett*, **1995**, 837-838.
7. The diastomeric excess was determined by GC chromatography on a chiral GC cyclodextrin column at 175°C. (Macherey-Nagel Lipodex column: [Oktakis-(2,6-di-o-pentyl-3-O-butyl)- γ -cyclo-dextrin]; retention time 7.5-8.2 min; 25 m). We refer to the diastereomeric excess as the ratio of *trans*-diol/*cis*-diol.
8. Eu(hfc)₃: tris[3-(heptafluoropropyl)hydroxymethylene-(+)-camphorato] europium (III).
9. Oetting, J.; Ph.D. thesis, **1994**, Universität Hannover, Hannover.
10. Wetter, H.; Oertle, K. *Tetrahedron Lett.* **1985**, 26, 5515-5518.
11. a) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, 48, 4155; b) Dess, D. B.; Martin, J. C.; *J. Am. Chem. Soc.* **1991**, 113, 7277.
12. Reetz, M. T.; Kessler, K.; Jung, A.; *Tetrahedron Lett.* **1984**, 25, 729.
13. **6:** $[\alpha]_D^{20} = -18.0$ (c 0.05 in CHCl_3); $^1\text{H-NMR}$ (400 MHz, CDCl_3 , 20°C, TMS): δ 7.32 (s, 5H; CH); 5.87 (dd, $^3\text{J}(\text{H,H}) = 17$ Hz, $^3\text{J}(\text{H,H}) = 11$ Hz, 1H; CH); 5.07-4.98 (m, 2H; CH_2); 4.58 (d, $^2\text{J}(\text{H,H}) = 11$ Hz, 1H; CH_2); 4.50 (d, $^2\text{J}(\text{H,H}) = 11$ Hz, 1H; CH_2); 4.22 (t, $^3\text{J}(\text{H,H}) = 6$ Hz, 1H; CH); 4.00-3.93 (m, 1H; CH); 3.91-3.82 (m, 1H; CH); 3.67 (t, $^3\text{J}(\text{H,H}) = 6\text{Hz}$, 1H; CH); 1.75-1.58 (m, 6H; CH_2); 1.20 (s, 3H; CH_3); 1.05 (s, 3H; CH_3); 0.88 (s, 9H; CH_3); 0.87 (s, 9H; CH_3); 0.06 (s, 6H; CH_3); 0.05 (s, 6H; CH_3); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3 , 20°C, TMS): 145.49 (-); 138.45 (+); 128-.79 (-); 128.45 (-); 127.75 (-); 112.51 (+); 75.37 (-); 74.43 (-); 1.09 (+); 68.16 (+); 67.12 (-); 59.44 (+); 42.11 (+); 41.53 (+); 40.97 (+); 25.94 (-); 22.80 (-); 22.65 (-); -4.20 (-); -5.32 (-); IR (CHCl_3): $\nu = 3467$ (w); 1720 (m); 1260 (s).

(Received in Germany 4 December 1995; revised 23 January 1996; accepted 24 January 1996)