



S0040-4039(96)00338-3


Is Chorismate Mutase a Prototypic Entropy Trap? – Activation Parameters for the *Bacillus subtilis* Enzyme

Peter Kast, M. Asif-Ullah and Donald Hilvert*

The Scripps Research Institute, Departments of Chemistry and Molecular Biology,
10666 North Torrey Pines Road, La Jolla, CA 92037 USA

Abstract: Chorismate mutase is thought to accelerate the chorismate-to-prephenate rearrangement in part by significantly lowering the entropy barrier for the reaction. We have determined the activation parameters for the well-characterized *Bacillus subtilis* chorismate mutase and find that ΔS^\ddagger (-9.1 ± 1.2 eu) is nearly as unfavorable as the activation entropy for the uncatalyzed process. Our results suggest that chorismate mutase catalysts show greater mechanistic versatility than commonly believed.
Copyright © 1996 Elsevier Science Ltd

The conversion of (–)-chorismate (**1**) to prephenate (**3**) by chorismate mutase constitutes a key step in the biosynthesis of the aromatic amino acids phenylalanine and tyrosine in bacteria, fungi and higher plants.¹ The reaction is formally a Claisen rearrangement and a rare example of an enzyme-catalyzed pericyclic process. The enzymatic reaction,^{2,3} like the uncatalyzed thermal rearrangement,⁴ proceeds through the chair-like transition state **2**, but the origins of the 10^6 to 10^7 -fold rate acceleration remain poorly understood.⁵ Comparison of the activation parameters for the uncatalyzed reaction and for chorismate mutases from *Klebsiella pneumoniae* and *Streptomyces aureofaciens* (Table 1) suggests that lowering the entropy barrier of the reaction by restricting the conformational degrees of freedom of the flexible chorismate molecule is a major factor in catalysis.⁶ This conclusion has been generally assumed to apply to all chorismate mutases,⁵ although the two enzymes on which it is based are relatively poorly characterized. Here we report activation parameters for a kinetically and structurally well-studied enzyme from *Bacillus subtilis* and show that ΔS^\ddagger in this case is unfavorable and comparable to that for the uncatalyzed reaction. This surprising observation contravenes conventional wisdom and suggests that chorismate mutases may exploit diverse strategies to achieve their otherwise similar catalytic effects.



The monofunctional chorismate mutase from *B. subtilis* displays simple kinetic characteristics. Its gene (*aroH*) has been cloned and expressed at high protein levels in the bacterium *Escherichia coli*.⁷ Furthermore, the three-dimensional structure of the protein trimer unliganded or complexed with a transition state analog or **3** has been determined.^{8,9} Using a modification¹⁰ of published protocols,^{7,11} we have purified *B. subtilis* chorismate mutase from the *E. coli* host strain KA12 transformed with plasmid pBSCM27 carrying *aroH*. Since our specifically engineered strain KA12¹² lacks both endogenous chorismate mutase genes, any contamination of the protein preparation with *E. coli* chorismate mutase can be excluded *a priori*. The activation parameters for the *B. subtilis* enzyme were obtained by determining the temperature dependence of k_{cat} (Figure 1). The disappearance of chorismate¹³ was followed spectrophotometrically at 274 nm ($\epsilon_{274 \text{ nm}} = 2.63 \text{ mM}^{-1} \text{ cm}^{-1}$) in a buffer containing 50 mM potassium phosphate, pH 7.5, at five chorismate concentrations ranging from 20 to 500 μM . All initial velocities were corrected for the corresponding temperature-specific background reaction. At each temperature, k_{cat} was determined as described⁶ assuming Michaelis-Menten kinetics.¹⁴ Thermodynamic activation parameters were obtained from least-squares analysis of data collected from 5 °C to 45 °C (Figure 1) using the equation $k_{\text{cat}} = (kT/h)e^{-[(\Delta H^\ddagger/RT) - (\Delta S^\ddagger/R)]}$.¹⁵




Figure 1. Temperature dependence of k_{cat} , the observed reaction constant for the rate-limiting step of *B. subtilis* chorismate mutase.

Table 1. Comparison of Thermodynamic Parameters for Catalyzed and Uncatalyzed Chorismate-to-Prephenate Rearrangements

| Source of chorismate mutase activity | ΔG^\ddagger [kcal mol ⁻¹] | ΔH^\ddagger [kcal mol ⁻¹] | ΔS^\ddagger [cal mol ⁻¹ K ⁻¹ (eu)] |
|---|--|--|---|
| Uncatalyzed reaction ¹⁶ | 24.5 | 20.7 ± 0.4 | -12.9 ± 0.4 |
| <i>Bacillus subtilis</i> ^b | 15.4 | 12.7 ± 0.4 | -9.1 ± 1.2 |
| <i>Klebsiella pneumoniae</i> ^c | 16.2 | 15.9 ± 0.4 | -1.1 ± 1.2 ^d |
| <i>Streptomyces aureofaciens</i> ⁶ | 15.0 | 14.5 ± 0.4 | -1.6 ± 1.1 ^d |
| Catalytic antibody 11F1-2E11 ¹⁷ | 18.7 | 18.3 | -1.2 |
| Catalytic antibody 1F7 ¹⁸ | 21.6 | 15 ± 2 | -22 ± 6 |

^aCalculated for 25 °C. ^bThis work. ^cFormerly known as *Aerobacter aerogenes*.⁶ ^dAssuming two (*K. pneumoniae*) and three (*S. aureofaciens*) active sites per enzyme molecule.⁶

At 30 °C, we found a k_{cat} of $46 (\pm 3) \text{ s}^{-1}$ per active site and a Michaelis constant K_{m} of $67 (\pm 5) \mu\text{M}$, which are similar to values obtained under comparable conditions by Knowles and coworkers.⁷ As shown in Table 1, the entropy of activation for the *B. subtilis* chorismate mutase ($\Delta S^{\ddagger} = -9.1 \text{ eu}$) is nearly as unfavorable as ΔS^{\ddagger} for the uncatalyzed reaction.¹⁶ Consequently, the rate acceleration achieved by the enzyme results in large part from decreasing the activation enthalpy by 8.0 kcal/mol; the $-T\Delta\Delta S^{\ddagger}$ term contributes only -1.1 kcal/mol , or 12%, to the gain in free energy of activation at 25 °C. This finding contrasts with that obtained for the enzymes from *K. pneumoniae* and *S. aureofaciens* and for the catalytic antibody 11F1-2E11¹⁷ where the entropic contribution to $\Delta\Delta G^{\ddagger}$ ranges from 35% to 60%. It has generally been assumed that the activation parameters in the latter three cases actually reflect the chemical transformation and are typical for the enzyme-catalyzed Claisen rearrangement.⁵ In contrast, product dissociation may be rate-limiting at substrate saturation for the *B. subtilis* enzyme,¹⁹ although contradictory evidence has been presented.^{11,19} Meaningful interpretation of the unfavorable ΔS^{\ddagger} for this enzyme must therefore await further experimental data, for instance from kinetic isotope effect studies. Nevertheless, given the similarity in k_{cat} values of the natural enzymes,^{6,7} the differences in the activation parameters are striking and imply different detailed catalytic mechanisms.

Despite an unfavorable ΔS^{\ddagger} , *B. subtilis* chorismate mutase is likely to exert considerable conformational control over the flexible chorismate molecule. The active site is highly complementary to the reactive pseudo-diaxial conformation of the substrate, with multiple hydrogen-bonding and electrostatic interactions poised to lock the enolpyruvyl side chain in place over the cyclohexadienyl ring.^{8,9} Similarly, the catalytic antibody 1F7,¹⁸ which has the most unfavorable ΔS^{\ddagger} of all the chorismate mutase catalysts examined (Table 1), appears to bind this high energy species preferentially.^{20,21} Nevertheless, the energetic contribution of such conformational control in the sense of a classical entropy trap²² is experimentally inaccessible if steps other than the chemical rearrangement are rate-limiting. Stabilization of the putative dipolar transition state through electrostatic and hydrogen-bonding interactions,²³⁻²⁷ also inferable from the crystal structures^{9,21,28,29} and experimentally examined for the *B. subtilis* enzyme,^{30,31} could well provide the major catalytic effect.⁵ In this respect, 1F7 is clearly less optimized than its natural counterparts.²¹ While much emphasis has been put on the favorable ΔS^{\ddagger} of several chorismate mutases in the past,^{5,6,17} our results with the *B. subtilis* enzyme caution against overgeneralization of these activation parameters and suggest that natural chorismate mutases may show greater mechanistic versatility than commonly believed.

Acknowledgment. We thank R. Pulido for advice and help in the preparation of chorismate. We are also grateful to J. R. Knowles for the gift of plasmid pBSCM2 and to K. Backman for providing strain KB224. This work was supported by a postdoctoral fellowship from the Schweizerischer Nationalfonds to P. K. and by grants from the Office of Naval Research and the National Institutes of Health (GM38273).

REFERENCES AND NOTES

1. Haslam, E. *Shikimic Acid: Metabolism and Metabolites*; John Wiley & Sons: New York, 1993.
2. Andrews, P. R.; Cain, E. N.; Rizzardo, E.; Smith, G. D. *Biochemistry* **1977**, *16*, 4848-4852.
3. Sogo, S. G.; Widlanski, T. S.; Hoare, J. H.; Grimshaw, C. E.; Berchtold, G. A.; Knowles, J. R. *J. Am. Chem. Soc.* **1984**, *106*, 2701-2703.

4. Copley, S. D.; Knowles, J. R. *J. Am. Chem. Soc.* **1985**, *107*, 5306-5308.
5. Lee, A. Y.; Stewart, J. D.; Clardy, J.; Ganem, B. *Chem. Biol.* **1995**, *2*, 195-203.
6. Görisch, H. *Biochemistry* **1978**, *17*, 3700-3705.
7. Gray, J. V.; Golinelli-Pimpaneau, B.; Knowles, J. R. *Biochemistry* **1990**, *29*, 376-383.
8. Chook, Y. M.; Ke, H.; Lipscomb, W. N. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 8600-8603.
9. Chook, Y. M.; Gray, J. V.; Ke, H.; Lipscomb, W. N. *J. Mol. Biol.* **1994**, *240*, 476-500.
10. We noticed that chorismate mutase is easily released from *E. coli* cells when treated with high osmolarity buffer [20% (w/v) sucrose/30 mM Tris-HCl, pH 8.0/1 mM EDTA]. The highly enriched chorismate mutase fraction was chromatographed on DEAE DE-52 cellulose and then Sephadex G-75. A detailed protocol will be published elsewhere. Electrospray mass spectrometric analysis gave a subunit mass of 14487 ± 2 Da (calculated mass = 14489 Da). A prominent signal at 14357 Da (≈ 25% of 14487 peak) corresponds to a processed protein with cleaved N-terminal methionine. We believe that this truncation (seen in independent purifications and in samples from outside sources) occurs *in vivo* during protein synthesis and does not affect our results.
11. Rajagopalan, J. S.; Taylor, K. M.; Jaffe, E. K. *Biochemistry* **1993**, *32*, 3965-3972.
12. KA12 was constructed by generalized P1 transduction to delete the *recA* gene from strain KB224 (Balakrishnan, R.; Backman, K. *Gene* **1988**, *67*, 97-103.) according to a procedure described earlier: Kast, P.; Hennecke, H. *J. Mol. Biol.* **1991**, *222*, 99-124.
13. Chorismate was isolated from *Klebsiella pneumoniae* strain 62-1 culture supernatants and purified by preparative HPLC essentially as described in the literature: Connelly, J. A.; Siehl, D. L. *Methods Enzymol.* **1987**, *142*, 422-431. The final preparation was checked by ¹H-NMR, analytical HPLC and, before each experiment, by quantitative conversion by chorismate mutase.
14. Chorismate mutase concentration was determined using the Pierce BCA Protein Assay Reagent (Pierce, Rockford, IL) with bovine serum albumin as the calibration standard.
15. Connors, K. A. *Chemical Kinetics. The Study of Reaction Rates in Solution*; VCH Publishers: New York, 1990; pp. 187-243.
16. Andrews, P. R.; Smith, G. D.; Young, I. G. *Biochemistry* **1973**, *12*, 3492-3498.
17. Jackson, D. Y.; Liang, M. N.; Bartlett, P. A.; Schultz, P. G. *Angew. Chem. Int. Ed. Engl.* **1992**, *31*, 182-183.
18. Hilvert, D.; Carpenter, S. H.; Nared, K. D.; Auditor, M.-T. M. *Proc. Natl. Acad. Sci. USA* **1988**, *85*, 4953-4955.
19. Gray, J. V.; Eren, D.; Knowles, J. R. *Biochemistry* **1990**, *29*, 8872-8878.
20. Campbell, A. P.; Tarasow, T. M.; Masefski, W.; Wright, P. E.; Hilvert, D. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 8663-8667.
21. Haynes, M. R.; Stura, E. A.; Hilvert, D.; Wilson, I. A. *Science* **1994**, *263*, 646-652.
22. Westheimer, F. H. *Adv. Enzymol. Relat. Areas Mol. Biol.* **1962**, *24*, 441-482.
23. Addadi, L.; Jaffe, E. K.; Knowles, J. R. *Biochemistry* **1983**, *22*, 4494-4501.
24. Copley, S. D.; Knowles, J. R. *J. Am. Chem. Soc.* **1987**, *109*, 5008-5013.
25. Gajewski, J. J.; Jurayj, J.; Kimbrough, D. R.; Gande, M. E.; Ganem, B.; Carpenter, B. K. *J. Am. Chem. Soc.* **1987**, *109*, 1170-1186.
26. Severance, D. L.; Jorgensen, W. L. *J. Am. Chem. Soc.* **1992**, *114*, 10966-10968.
27. Wiest, O.; Houk, K. N. *J. Org. Chem.* **1994**, *59*, 7582-7584.
28. Lee, A. Y.; Karplus, P. A.; Ganem, B.; Clardy, J. *J. Am. Chem. Soc.* **1995**, *117*, 3627-3628.
29. Xue, Y.; Lipscomb, W. N. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 10595-10598.
30. Kast, P.; Hartgerink, J. D.; Asif-Ullah, M.; Hilvert, D. *J. Am. Chem. Soc.* **1996**, *118*, *in press*.
31. Kast, P.; Asif-Ullah, M.; Jiang, N.; Hilvert, D. *Proc. Natl. Acad. Sci. USA* **1996**, *in press*.

(Received in USA 11 December 1995; revised 19 February 1996; accepted 21 February 1996)