



S0040-4039(96)00416-9


2,2-Di(ethoxy)vinyl lithium : A Synthetic Equivalent of the Ethyl Acetate Anion

Heng-xu Wei and Manfred Schlosser *


Institut de Chimie organique de l'Université
Bâtiment de Chimie (BCh), CH-1015 Lausanne-Dorigny, Switzerland

Abstract : 2,2-Diethoxyvinyl lithium can be readily generated from 2-bromo-1,1-diethoxyethylene by treatment with butyllithium. When dissolved in tetrahydrofuran, it can be stored at -25 °C for hours, but decomposes rapidly at 0 °C. Copyright © 1996 Elsevier Science Ltd


In the same way as 2-ethoxyvinyl lithium ^{1, 2} can be conceived as a synthetical equivalent of the acetaldehyde derived enolate, 2,2-diethoxyvinyl lithium **1** may be used as a masked ethyl acetate (ethoxycarbonylmethyl) anion. The organometallic reagent is instantaneously formed when butyllithium is added to an ethereal solution of 2-bromo-1,1-diethoxyethylene. Interception with electrophiles such as iodine or iodonium chloride, chlorotrimethylsilane, dimethyl sulfate, benzylbromide, butanal and pivalaldehyde gives the ketene acetals **2** (64%), **3** (31%), **4** (61%), **5** (64%), **6** (65% [after treatment of alcoholate with chlorotrimethylsilane : MO = LiO → (H₃C)₃SiO]) and **7** (67%).³



The ketene acetals may be immediately submitted to acid catalyzed hydrolysis to afford the corresponding ethyl esters. When aldehydes had been employed as the electrophiles, simultaneous dehydration of the intermediate β -hydroxy esters occurs producing α,β -unsaturated ester such as ethyl 4,4-dimethylpent-2-enoate (**8**, 65%), ethyl cinnamate (**9**, 94%) and ethyl *p*-chlorocinnamate (**10**, 78%).



In the same way as 2,2-diethoxyvinyl lithium, 2,2-dimethoxyvinyl lithium and 2,2-(1,3-dioxolan-2,2-ylidene)-vinyl lithium have been generated from the corresponding bromides **11** ⁴ and **13** ⁴ by halogen/metal interconversion. Reaction with chlorotrimethylsilane at -75 °C afforded (2,2-dimethoxyvinyl)trimethylsilane (**12**, 67%) and [(2,2-dioxolan-2,2-ylidene)methyl]trimethylsilane (**14**, 72%), respectively.



Solutions of all these 2,2-dialkoxyvinyl lithium reagents in tetrahydrofuran can be stored for hours at -25 °C. Thus, their thermal stability equals or even surpasses that of (*Z*)-2-ethoxyvinyl lithium ¹ whereas the corresponding (*E*)-isomer was reported to decompose already at temperatures below -80 °C ¹. Comparing these structures, one is at first sight inclined to attribute a stabilizing, dipolar or coordinative effect to the alkoxy group occupying the *cis* position with respect to the metal. However, this conclusion might be fallacious since it does not take into account the differences in n- π conjugation energies ⁵ between ethoxyalkenes (enethers) and alkoxyacetylenes, the precursors and products of the elimination reactions under inspection. In other words, a more profound investigation deems warranted.

Typical working procedure : Precooled (-75 °C) tetrahydrofuran (40 mL) and 2-bromo-1,1-diethoxyethylene ⁶ (**1**; 3.7 mL, 4.9 g, 25 mmol) were consecutively added to butyllithium (25 mmol) from which the commercial solvent (hexane) had been stripped off. After 30 min at -75 °C, chlorotrimethylsilane (3.1 mL, 2.7 g, 25 mmol) was added. When the mixture had reached 25 °C, the solvent was evaporated and the residue distilled to afford (2,2-diethoxyvinyl)trimethylsilane as a colorless liquid; bp 59 - 60 °C/9 mmHg; n_D^{20} 1.4308; 64%.

Acknowledgment: This work was supported by the Schweizerische Nationalfonds zur Förderung der wissenschaftlichen Forschung, Bern (grant 20-41'887-94).

References

- 1 Generation by bromine/lithium interconversion : Lau, K.S.Y.; Schlosser, M.; *J. Org. Chem.* **1978**, *43*, 1595 - 1598.
- 2 Generation by tributyltin/lithium interconversion : Ficini, J.; Falou, S.; Touzin, A.M.; d'Angelo, J.; *Tetrahedron Lett.* **1977**, *40*, 3589 - 3592; Wollenberg, R.H.; Albizati, K.F.; Peries, R.; *J. Am. Chem. Soc.* **1977**, *99*, 7365 - 7367.
- 3 The identity and purity of all new compounds is corroborated by correct elementary analyses and spectral data (to be published in a full-length article).
- 4 McElvain, S.M.; Curry, M.J.; *J. Am. Chem. Soc.* **1948**, *70*, 3782 - 3786.
- 5 Greenberg, A.; Stevenson, T.A.; *J. Am. Chem. Soc.* **1985**, *107*, 3488 - 3494.
- 6 Beyerstedt, F.; McElvain, S.M.; *J. Am. Chem. Soc.* **1937**, *59*, 2266 - 2269.

(Received in France 5 February 1996; accepted 28 February 1996)