



Conformationally Constrained Bis(oxazoline) Derived Chiral Catalyst : A Highly Effective Enantioselective Diels-Alder Reaction

Arun K. Ghosh,* Packiarajan Mathivanan and John Cappiello

Department of Chemistry, University of Illinois at Chicago,
845 West Taylor Street, Chicago, Illinois 60607.

Abstract: The reaction of cyclopentadiene with various bidentate dienophiles in the presence of 4-10 mol% of copper(II)-bis(oxazoline) complexes afforded excellent endo/exo selectivity as well as endo enantioselectivity (95-99% ee) and isolated yields. On the other hand, Diels-Alder reaction with Mg(II)-complexes afforded a modest (61% ee) reversal in enantioselectivity.

Copyright © 1996 Elsevier Science Ltd

The enantioselective Diels-Alder reaction is of paramount interest in organic synthesis.¹ As a consequence, a number of effective enantioselective processes have been developed over the years.² Recent discoveries of various catalytic processes have added a new dimension to the enantioselective Diels-Alder reaction.^{3,4} Of particular interest, C₂-symmetric chiral bis(oxazoline)-metal complexes have shown exceptional promise as chiral catalysts for enantioselective Diels-Alder reactions.⁵ As part of our interests in various ligand assisted asymmetric syntheses,⁶ we have synthesized conformationally constrained chiral bis(oxazoline) ligands and investigated their effectiveness as chiral catalysts in asymmetric Diels-Alder reactions. Recent reports by Merck group⁷ on the synthesis and application of (1S, 2R)-1-amino-2-indanol derived other bis(oxazoline) ligands in catalytic Diels-Alder reaction, prompted us to disclose our results in this area. Herein, we report that various metal-ligand complexes of conformationally constrained bis(oxazoline) ligands derived from phenylglycinol surrogates, *cis*-1-amino-2-indanols⁸ are indeed highly effective catalysts for the enantioselective Diels-Alder reaction. Among various metal complexes examined, bis(oxazoline)-copper(II) complex is particularly effective. Reaction of cyclopentadiene with various bidentate dienophiles **6** in the presence of 4-10 mol% of bis(oxazoline)-copper(II) complexes afforded excellent endo/exo selectivity as well as endo enantioselectivity and excellent isolated yields. Since either enantiomer of constrained bis(oxazoline) is readily prepared from commercially available⁹ optically active *cis*-1-amino-2-indanols, the current methodology provides a convenient access to either enantiomer of various cyclopentadiene based Diels-Alder adducts in a stereopredictable fashion.

Conformationally constrained ligand **4** was readily prepared on a multi gram scale according to the procedure of Lehn.¹⁰ Treatment of malononitrile and ethanol in dioxane in the presence of anhydrous HCl afforded the amide enol ether hydrochloride salt **1** (Scheme I). Reaction of **1** with (1R, 2S)-1-amino-2-indanol **2** in DMF at 23°C provided the bis(oxazoline) **4** as a white solid (m.p. 206-208°C; $\alpha_D^{23} + 352$, c, 3.7, CHCl₃) after recrystallization from a mixture (1:1) of chloroform and methanol (60 % yield). Similarly, (1S, 2R)-1-amino-2-indanol **3** was converted to constrained bis(oxazoline) **5** (m.p. 207-209°C; $\alpha_D^{23} - 352.7$, c, 3.0, CHCl₃). The Diels-Alder reactions of bidentate dienophiles **6a-d** and cyclopentadiene in the presence of various chiral catalysts derived from the above ligands and metal triflates or halides, were investigated. For the Cu(II)-catalyzed reaction, the metal-ligand complex was prepared^{5c} by reaction of an equimolar mixture of Cu(II) triflate and bis(oxazoline) **4** in dry CH₂Cl₂ at 23°C under nitrogen atmosphere.

The resulting dark blue solution of the catalyst (4-50 mol%) was stirred for 1 h at 23°C and then cooled to the specified temperature and 1 equiv of dienophile (**6 a-d**) followed by 4-6 equiv of freshly cracked cyclopentadiene were added. After workup with saturated aqueous NH₄Cl solution and chromatography over silica gel, the resulting Diels-Alder cycloadducts were analyzed by ¹H-NMR (400MHz) and chiral HPLC analysis (Daicel Chiracel OD column,¹¹ 10% isopropanol/hexane as the eluent). The results of various metal catalyzed Diels-Alder reaction with a range of bidentate dienophiles are summarized in Table I. As shown, Cu(II)-catalyzed reaction with acryloyl-N-oxazolidinone **6a** proceeded with virtually complete endo selectivity (endo/exo ratio >99 :1) as well as excellent endo enantioselectivity (97-99% ee) and isolated yields. Such selectivity can even be observed with 4 mol% catalyst (entry 4). The endo adduct **7a** was formed exclusively with 99% enantiomeric excess ($\alpha_D^{23^\circ} +172$, *c* 6.1, CHCl₃; lit.¹²; $\alpha_D^{23^\circ} +171.05^\circ$, *c*, 1.04, CHCl₃) and 90% yield after silica gel chromatography. The identities of endo and exo isomers as well as endo enantioselectivities of the cycloadducts **7a** and **8a** were determined by chiral HPLC analysis. The absolute configuration of the cycloadducts **7a** (2*R*-configuration) and **8a** (2*S*-configuration) was assigned based upon the comparison of optical rotations with the literature values.¹² The Diels-Alder reaction of fumaroyl-N-oxazolidinone **6d**¹³ with 10 mol% Cu(II)-catalyst (entry 12) proceeded well at -45°C for 8 h providing a 93:7 endo/exo mixture ratio and 94% endo enantiomeric excess. The enantiomeric excess of endo adduct was determined after conversion of the cycloadducts to the corresponding iodolactone **9** ($\alpha_D^{23^\circ} -39.2$, *c* 4.65, CHCl₃; lit.^{5c}; $\alpha_D^{23^\circ} +40^\circ$, 96% ee, *c*, 1.78, CHCl₃).^{14, 15}

Scheme I




Table 1. Enantioselective Diels-Alder Reaction of Imides **6** with cyclopentadiene

Entry	Dienophile	Metal (mol %)	temp (time,h)	% Yield ^a	endo/exo ^d	endo ee (config.) ^e
1.	6a (R=H)	Cu (50)	-78°C (6)	78	>99 : 1	97 (2R)
2.	6a (R=H)	Cu (8)	-78°C (8)	94	>99 : 1	98 (2R)
3.	6a (R=H)	Cu (8) ^b	-78°C (8)	98	>99 : 1	94 (2S) ^b
4.	6a (R=H)	Cu (4)	-78°C (8)	90	>99 : 1	99 (2R)
5.	6a (R=H)	Mg (100)	-78°C (7)	81	98 : 2	61 (2S)
6.	6a (R=H)	Mg (10)	-78°C (7)	76	95 : 5	34 (2S)
7.	6a (R=H)	Fe (100)	-78°C (7)	79	94 : 6	65 (2R)
8.	6b (R=Me)	Cu (40)	0°C (26) ^f	84	92 : 8	94 (2R)
9.	6b (R=Me)	Cu (10)	0°C (48) ^f	77	90 : 10	84 (2R)
10.	6b (R=Me)	Mg (10)	0°C (48) ^f	76 ^c	92 : 8	55 (2S)
11.	6c (R=Ph)	Cu (10)	23°C (72) ^f	78	80 : 20	35 (2S)
12.	6d (R=CO ₂ Et)	Cu (10)	-45°C (8)	75	93 : 7	94 (2S)

^a Isolated yield after silica gel chromatography. ^b With 1S, 2R-bisoxazoline ligand. ^c Based on recovered starting material. ^d Endo/exo ratios were determined by ¹H-NMR and HPLC.

^e Enantiomeric excess was determined by HPLC and comparison of known optical rotation.

^f At -78°C, no reaction or very little conversion was obtained.


The high degree of enantioface selection in the current Cu(II)-bis(oxazoline) catalyzed Diels-Alder reaction can be rationalized based on transition state models proposed by Corey and Evans in which Cu(II) assumes a square planar complex with the bis(oxazoline) ligand **4** and bidentate dienophile while the dienophile assumes a *s-cis* conformation.⁵ Based on such a model, the endo-*Si*-face attack of cyclopentadiene is favored and this leads to the observed 2R-enantioselectivity (for **6a-b**) with ligand **4**. The reversal of enantioselectivity with the Mg(II)-bis(oxazoline) catalyzed reaction (for **6a-b**) could be explained by a Corey-Isihara type transition state^{5b} model in which the Mg(II)-complex (with **4**) assumes a tetrahedral geometry rather than a square planar geometry most probable for Cu(II). Now, based on such molecular arrangements, it appears that the endo-*Re*-face attack of the diene is favored providing 2S-enantioselectivity.¹⁶

In summary, the present conformationally constrained bis(oxazoline) based metal-ligand complexes particularly Cu(II)-complexes represent a highly effective catalytic system for enantioselective Diels-Alder reactions. Since both enantiomers of the bis(oxazoline) ligand are readily prepared from the commercially available optically active *cis*-1-amino-2-indanols, the current catalytic Diels-Alder technology provides a convenient access to either enantiomer of the cycloadducts in a stereopredictable fashion with high optical purity and isolated yields. Application of these chiral catalysts in synthesis is currently on going in our laboratory.

Acknowledgment: Financial support for this work was provided by the University of Illinois at Chicago. The authors thank Dr. Kurt Ritter of BASF, Germany for a gift of optically active *cis*-1-amino-2-indanol and Professor Robert Moriarty for helpful discussions.

References and Notes:

- (a) Carruthers, W. *Cycloaddition Reactions in Organic Synthesis*, 1990, Pergamon press; (b) Giuliano, R. M. *Cycloaddition Reactions in Carbohydrate Chemistry*; American Chemical Society: Washington, DC, 1992.
- (a) Oppolzer, W. *Comprehensive organic syntheses :Selectivity, Strategy and Efficiency in Modern Organic Chemistry*; Trost, B. M., Ed.; Pergamon Press: Oxford, 1991; vol. 5, p 315; (b) Corey, E. J. *Pure Appl. Chem.* 1990, 62, 1209; (c) Waldmann, H.; Braun, M. *Gazz. Chim. Ital.* 1991, 121, 277; (d) Waldmann, H. *Synthesis* 1994, 535 and references cited therein.
- (a) Corey, E. J.; Sarshar, S.; Lee, D.-H. *J. Am. Chem. Soc.* 1994, 116, 12089; (b) Corey, E. J.; Lee, D.-H. *Tetrahedron Asymmetry* 1995, 6, 3; (c) Corey, E. J.; Sarshar, S.; Bordner, J. *J. Am. Chem. Soc.* 1992, 114, 7938; (d) Corey, E. J.; Imwinkelreid, R.; Pikul, S.; Xiang, Y. B. *J. Am. Chem. Soc.* 1989, 111, 5493; (e) Corey, E. J.; Matsumura, Y.; *Tetrahedron Lett.* 1991, 32, 6289; (f) Narasaka, K.; Iwasawa, N.; Inoue, M.; Yamada, T.; Nakashima, M.; Sugimori, J. *J. Am. Chem. Soc.* 1989, 111, 5340; (g) Iwasawa, N.; Sugimori, J.; Kawase, Y.; Narasaka, K. *Chem. Lett.* 1989, 1581 and references cited therein.
- (a) Furuta, K.; Shimizu, S.; Miwa, Y.; Yamamoto, H. *J. Org. Chem.* 1989, 54, 1481; (b) Takemura, H.; Komeshima, N.; Takahashi, I.; Hashimoto, S-I.; Ikota, N.; Tomioka, K.; Koga, K. *Tetrahedron Lett.* 1987, 28, 5687; (c) Takasu, M.; Yamamoto, H.; *Synlett* 1990, 194; (d) Sartor, D.; Saffrich, J.; Helmchen, G. *Synlett* 1990, 197; (e) Kagan, H. B.; Riant, O. *Chem. Rev.* 1992, 92, 1007; (f) Pindur, U.; Lutz, G.; Otto, C. *Chem. Rev.* 1993, 93, 741; (g) Deloux, L.; Srebnik, M. *Chem. Rev.* 1993, 93, 763; (h) Togni, A.; Venanzi, L. M. *Angew. Chem. Int. Ed. Engl.* 1994, 33, 497; (i) Oh, T.; Reilly, M. *Org. Prep. Proceed. Int.* 1994, 26, 129; (j) Seebach, D.; Dahinden, R.; Marti, R. E.; Beck, A. K.; Plattner, D. A.; Kuhnle, N. M. *J. Org. Chem.* 1995, 60, 1788.
- (a) Corey, E. J.; Imai, N.; Zhang, H-Y. *J. Am. Chem. Soc.* 1991, 113, 728; (b) Corey, E. J.; Isihara, K. *Tetrahedron Lett.* 1992, 33, 6807; (c) Evans, D. A.; Miller, S. J.; Lectka, T. *J. Am. Chem. Soc.* 1993, 115, 6460; (d) Evans, D. A.; Murry, J. A.; Mat, P. V.; Norcross, R. D.; Miller, S. J. *Angew. Chem. Int. Ed. Engl.* 1995, 34, 798.
- (a) Ghosh, A. K.; Duong, T. T.; McKee, S. P. *J. Chem. Soc., Chem. Commun.* 1992, 1673; (b) Ghosh, A. K.; Chen, Y. *Tetrahedron Lett.* 1995, 36, 6811; (c) Ghosh, A. K.; Liu, W. *J. Org. Chem.* 1995, 60, 6198; (d) Ghosh, A. K.; Onishi, M. *J. Am. Chem. Soc.* 1996, 118, 2527; (e) Ghosh, A. K.; Mathivanan, P. *Tetrahedron: Asymmetry* 1996, 7, 375.
- (a) Davies, I. W.; Senanayake, C. H.; Larsen, R. D.; Verhoeven, T. R.; Reider, P. J. *Tetrahedron Lett.* 1996, 37, 813; (b) Davies, I. W.; Senanayake, C. H.; Larsen, R. D.; Verhoeven, T. R.; Reider, P. J. *ibid.* 1996, 37, 1725.
- Ghosh, A. K.; McKee, S. P.; Sanders, W. M. *Tetrahedron Lett.* 1991, 32, 711.
- Available from Aldrich Chemical Co., Milwaukee, WI 53201 and Sepracor Inc., Marlborough, MA 01752.
- Hall, J.; Lehn, J-M.; DeCian, A.; Fischer, J. *Helv. Chim. Acta.* 1991, 74, 1.
- Daicel OD column was purchased from Chiral Technologies, 730 Springdale Dr, Exton, PA.
- Narasaka, K.; Inoue, M.; Okada, N. *Chem. Lett.* 1986, 1109; and see refernece 3f.
- Kool, J.; Feringa, B. L. *Synlett* 1995, 1025.
- The iodolactone **9** was conveniently prepared by exposure of the adducts to iodine, KI and NaHCO₃ in aqueous CH₂Cl₂ at 23°C for 16 h (72% yield).



- All new compounds gave satisfactory spectroscopic and analytical results.
- The scope of the current constrained ligand in the Diels-Alder reactions with other dienes and counter ions has not been investigated.

(Received in USA 29 March 1996; revised 15 April 1996; accepted 16 April 1996)