


Photodecarbonylation of 2-Phenyl-4-alkylidene-5(4H)-oxazolones

Bokyung Jung, Hongbum Kim and Bong Ser Park*,

Department of Chemistry, Dongguk University, Seoul 100-715, Korea

Abstract: The title compounds were prepared by the Erlenmeyer synthesis and irradiated with or without a Pyrex filter using 450 W medium pressure mercury arc lamp. Cis-trans isomerization was the major reaction under pyrex filtered condition, whereas irradiation without a Pyrex filter resulted in decarbonylation, followed by trapping of the ketenimine intermediate by protic solvents.
 Copyright © 1996 Elsevier Science Ltd

For decades, several papers describing photochemistry of 2-phenyl-4-alkylidene-5(4H)-oxazolones have been appeared in the literature¹. Even though some of these compounds showed various photoreactivities depending upon their substituents and reaction media, most of them involved photoinduced cis-trans isomerization of the alkylidene moieties. Ullman and coworkers have studied the wavelength dependence on photochemistry of 2-phenyl-4-benzylidene-5(4H)-oxazolone, **1a**.² They reported that irradiation of **1a** at 365 nm resulted in cis-trans isomerization, while irradiation at 254 nm resulted in hydrogen atom abstraction from a solvent, 2-propanol. This report led us to study photobehavior of compounds **1b** and **1c** shown below, which we expected those E isomers to react by an intramolecular H-atom abstraction. Contrary to this expectation, these compounds showed different photoreactivities. Here we would like to report what we have found so far on this system.




a: R = Ph
 b: = Me
 c: = Et


The title compounds were prepared by Erlenmeyer synthesis³, which gave mixtures of Z- and E-isomers with a ratio of 3 to 1 for both **1b** and **1c**.⁴ The amount of E-isomers could be increased by using known methods⁵, but they were slowly isomerized to the more stable Z-isomers in solution at room temperature. Especially, the E-isomer of **1c** was not stable enough to be used for photolysis in pure form.

Irradiation was performed using 450 W Hanovia medium pressure mercury arc lamp with or without a Pyrex filter sleeve. Several different solvents were used for the photolysis; ethanol, 2-propanol, *tert*-butanol, benzene, THF and acetonitrile. With the Pyrex filter, the only reaction observed was *cis*-*trans* isomerization in all the solvents used. The E to Z ratios in photostationary mixtures were 3 to 1 and 1 to 1 for **1b** and **1c**, respectively.


The photolysis of **1b** and **1c** in alcoholic solvents without a Pyrex filter resulted in formation of a sole product.⁶ The product was identified as **2**, whose chemical yield was over 90%.⁷ We could not detect any products of hydrogen abstractions from solvents, which Ullman and coworkers had observed for **1a**.²



When **1b** and **1c** were irradiated in acetonitrile or THF containing a few percents of water, one product was formed predominantly, which was identified to be **3**.⁸ Even if well dried solvents were used for photolysis, the same product was obtained, which we believe some moisture got involved during the workup. When D₂O was added in the reaction mixture, **3-d₂** was obtained as a sole product after photolysis.⁹ When **1b** was irradiated in a degassed quartz tube using CD₃CN as a solvent and transferred to an NMR tube, a new product was observed besides **3b**.¹⁰ This product was tentatively assigned as **4b** based on its ¹H NMR spectroscopic data.¹¹ Further characterization of this compound could not be done due to its instability.



A plausible explanation based on our results is that **1b** and **1c** are decarbonylated and form ketenimine **4** on photolysis without a Pyrex filter. The ketenimine is rapidly trapped by protic solvents. Photodecarbonylation has been suggested as a reaction mechanism of *N*-(methylidene)acetamide formation from 2,4-dimethyl-4-phenyl- Δ^2 -oxazoline-5-one in which an alkylidene moiety is absent at 4 position.¹² To our knowledge, however, our result is the first direct observation of photodecarbonylation from 4-alkylidene-2-phenyl-5(4H)-oxazolones. Our results are unique in that the photodecarbonylation forms reactive ketenimines, which can be further elaborated toward synthetic applications. Also it can be a good candidate for photoaffinity labeling.¹³



Ullman has discussed the wavelength dependent photoreactivity of **1a** in terms of involvement of different excited states. His theoretical views predicted that the upper excited state $T_2(n, \pi^*)$ was responsible for hydrogen abstraction reaction and the lower excited state $T_1(\pi, \pi^*)$ was the origin of cis-trans isomerization. In our experiments with **1b** and **1c**, decarbonylation is observed unlike hydrogen abstraction for **1a** on photolysis at the shorter wavelength.¹⁴ The ultraviolet absorption spectra of **1a** and **1b** look similar, but the absorption maxima of **1a** is at the longer wavelength than those of **1b**; 259, 360 nm for **1a** and 236, 298 nm for **1b**. We think the excited state $T_2(n, \pi^*)$ is also responsible for decarbonylation reaction and the change of the reaction pathway may have resulted from the higher energy of **1b**.

In summary, irradiation of 2-phenyl-4-ethylidene(or propylidene)-5(4H)-oxazolone at short wavelength resulted in an efficient photodecarbonylation followed by trapping of the ketenimine intermediate by protic solvents, which was unprecedented for photochemistry of 2-phenyl-4-alkylidene-5(4H)-oxazolones so far. The detailed mechanistic picture of this reaction will be described in a full paper in due time.

Acknowledgement This work was generously supported by the Basic Science Research Institute Program, Ministry of Education (BSRI-95-3406 for B.S. P. and BSRI-95-3417 for H. K.) and in part by the Korea Science and Engineering Foundation (93-0500-09-01-3).

REFERENCES AND NOTES

1. (a) Rao, Y. S. *Synthesis* **1975**, 749-764, and refs therein. (b) Dixit, V. M.; Bhat, V.; Trozzolo, A. M.; George, M. V. *J. Org. Chem.* **1979**, *44*, 4169-4173. (c) Lawrenz, D.; Mohr, S.; Wendrander, B. *J. Chem. Soc. Chem. Commun.* **1984**, 863-865. (d) Walter, R.; Purcell, T. C.; Zimmer, H. *J. Heterocycl. Chem.* **1966**, 235-236.
2. (a) Baumann, N.; Song, M.; Ullman, E. F. *J. Am. Chem. Soc.* **1968**, *90*, 4157-4158. (b) Ullman, E. F.; Maumann, N. *J. Am. Chem. Soc.* **1968**, *90*, 4158-4160. (c) Ullman, E. F.; Maumann, N. *J. Am. Chem. Soc.* **1970**, *92*, 5892-5899.
3. Carter, H. E. *Org. React.* **1974**, *3*, 199-239.

4. The structural assignment of each isomer is based on relative stability of each isomer and on Sidhu's general observations on the NMR spectra of oxazolones; Sidhu, G. S.; Venkataratnam, V.; Prasad, K. K.; Iyengar, D. S. *Ind. J. Chem.* **1972**, *10*, 448-451. **Z-1b**: ^1H NMR (CDCl_3 , 200 MHz); δ 8.10 (d, 2H, $J = 8.4$ Hz), 7.60-7.51 (m, 3H), 6.76 (q, 1H, $J = 7.4$ Hz), 2.26 (d, 3H, $J = 7.4$ Hz); **E-1b**: ^1H NMR (CDCl_3 , 200 MHz); δ 8.05 (d, 2H, $J = 7.6$ Hz), 7.59-7.45 (m, 3H), 6.91 (q, 1H, $J = 7.9$ Hz), 2.37 (d, 3H, $J = 7.9$ Hz); **Z-1c**: ^1H NMR (CDCl_3 , 200 MHz); δ 8.10 (d, 2H, $J = 7.4$ Hz), 7.60-7.47 (m, 3H), 6.70 (t, 1H, $J = 7.9$ Hz), 2.71 (dq, 2H, $J = 7.9, 7.7$ Hz), 1.20 (t, 3H, $J = 7.7$ Hz); **E-1c**: ^1H NMR (CDCl_3 , 200 MHz); δ 8.06 (d, 2H, $J = 7.4$ Hz), 7.60-7.45 (m, 3H), 6.83 (t, 1H, $J = 7.9$ Hz), 2.86 (dq, 2H, $J = 7.9, 7.7$ Hz), 1.18 (t, 3H, $J = 7.7$ Hz). Other physical properties of **1b** were the same as the reported values.(ref 3 and refs therein)
5. Rao, Y. S. *J. Org. Chem.* **1976**, *41*, 722-725.
6. Irradiation of one isomer only or the mixture gave the same result.
7. Shown here are two representative cases where R' is an isopropyl group; **2b**: ^1H NMR (CDCl_3 , 200 MHz); δ 8.01 (d, 2H, $J = 7.4$ Hz), 7.60-7.40 (m, 3H), 5.19 (septet, 1H, $J = 7.0$ Hz), 2.38 (q, 2H, $J = 7.4$ Hz), 1.38 (d, 6H, $J = 7.0$ Hz), 1.12 (t, 3H, $J = 7.4$ Hz), IR(CCl_4) 1667, 1549, 1268 cm^{-1} , EI MS 219(M^+ , 3), 105(100), 77(48), 59(34). **2c**: ^1H NMR (CDCl_3 , 200 MHz); δ 8.00 (d, 2H, $J = 7.4$ Hz), 7.60-7.40 (m, 3H), 5.20 (septet, 1H, $J = 7.0$ Hz), 2.32 (t, 2H, $J = 7.4$ Hz), 1.59 (tq, 2H, $J = 7.4, 7.5$ Hz), 1.35 (d, 6H, $J = 7.0$ Hz), 0.88 (t, 3H, $J = 7.5$ Hz), IR(CCl_4) 1664, 1539, 1265 cm^{-1} , EI MS 233(M^+ , 2), 105(100), 77(48). The stereochemistry of the product could not be determined by any methods that we had tried so far. It is possible that this compound exists as stereolabile E, Z isomers that interconvert with a low free energy of activation.(See Guerra, A.; Lunazi, L. *J. Org. Chem.* **1995**, *60*, 7959-7965.) We, however, failed to observe such a phenomenon by low temperature NMR spectroscopy due to solubility problems.
8. **3b**: ^1H NMR (CDCl_3 , 200 MHz); δ 9.01 (br s, 1H), 7.89 (d, 2H, $J = 7.4$ Hz), 7.55-7.45 (m, 3H), 3.01 (q, 2H, $J = 7.4$ Hz), 1.24 (t, 3H, $J = 7.4$ Hz). IR(CCl_4) 3297, 1712, 1682 cm^{-1} , EI MS 177(M^+ , 13), 105(100), 77(47), 57(20), 51(21). **3c**: ^1H NMR (CDCl_3 , 200 MHz); δ 8.43 (br s, 1H), 7.84 (d, 2H, $J = 7.4$ Hz), 7.60-7.45 (m, 3H), 2.99 (t, 2H, $J = 7.4$ Hz), 1.77 (tq, 2H, $J = 7.4$ Hz), 1.03 (t, 3H, $J = 7.4$ Hz). IR(CCl_4) 3306, 1718, 1685 cm^{-1} , EI MS 191(M^+ , 8), 105(100), 77(56).
9. **3b-d₂**: ^1H NMR (CDCl_3 , 200 MHz) δ 7.89 (d, 2H, $J = 7.4$ Hz), 7.55-7.45 (m, 3H), 3.01 (br, 1H), 1.24 (br d, 3H, $J = 7.3$ Hz).
10. The amounts of products varied from trace to ten percents depending upon the reaction time under this reaction condition. The reaction mixture became complicated at high conversion.
11. **4b**: ^1H NMR (CD_3CN , 200 MHz) δ 8.00 (d, 2H, $J = 8.0$ Hz), 7.71-7.45 (m, 3H), 4.70 (q, 1H, $J = 8.0$ Hz), 1.59 (d, 3H, $J = 8.0$ Hz).
12. Johnson, M. R.; Sousa, L. R. *J. Org. Chem.* **1977**, *42*, 2439-2443.
13. Fleming, S. A. *Tetrahedron* **1995**, *51*, 12479-12520.
14. When **1a** was irradiated under our reaction condition, no decarbonylation products were detected. The hydrogen abstraction product that Ullman had reported was observed together with some other unidentifiable products.

(Received in Japan 22 February 1996; revised 15 April 1996; accepted 17 April 1996)