

Palladium Catalysed Cascade Hydrostannylation-Bis-Cyclisation-Intramolecular Anion Capture. Routes to Bridged- and Spiro-Cyclic Small and Macrocyclic Heterocycles.

Adele Casaschi,^a Ronald Grigg,^{a*} José M Sansano,^a David Wilson,^a and James Redpath.^b

a. School of Chemistry, Leeds University, Leeds LS2 9JT

b. Organon Laboratories, Newhouse, Lanarkshire ML1 5SH


Abstract: Palladium catalysed cascade hydrostannylation of terminal alkynes in α,ω -enynes at 0-25°C followed by bis-cyclisation initiated by insertion of Pd(0) into an aryl iodide at 100-110°C and terminating in an intramolecular Stille coupling provides a wide range of exo-methylene 5/6 and 5/12-17 membered spiro- and bridged-ring heterocycles. Copyright © 1996 Elsevier Science Ltd

A growing number of imaginative applications of the intramolecular Stille reaction for the construction of macrocyclic natural products have been reported following the pioneering work of Stille, Hegedus and Hirama.¹⁻³ These applications have mainly employed sp^2 - sp^2 Stille coupling apart from Hirama's and Danishefsky's (sp - sp^2) work.^{2,3} Macrocyclic formation via sp^2 - sp^3 Stille coupling has not been reported. We have shown that organotin(IV) reagents are excellent anion capture agents in our intermolecular palladium catalysed cascade cyclisation-anion capture processes⁴ and that anion capture is invariably slower than cyclisation when 3-7 membered rings are being constructed. Moreover, these cascade processes offer the opportunity to explore sp^2 - sp^3 Stille coupling as part of a polycyclisation strategy which incorporates a macrocyclisation. The successful implementation of such processes via palladium catalysed cascade cyclisation-intramolecular anion capture is reported herein.

The ready Pd(0) catalysed hydrostannylation of alkynes offered the possibility of a Pd(0) catalysed cascade process in which cyclisation-anion capture occurs intramolecularly. Good to excellent regioselective hydrostannylation of terminal alkynes can be achieved by the incorporation of a proximate (β - or γ -) heteroatom⁵. Bis-cyclisation processes involving creation of two small rings were explored as a prelude to macrocyclisation studies. In developing a cascade cyclisation-intramolecular anion capture process for small ring polycycles it is essential to generate the α -vinylstannane (1) since the β -vinylstannane (2) presents a sterically impossible barrier to anion capture.


A series of enynes (3a-c) (Scheme 1) was allowed to react with Bu_3SnH (1eq) in toluene at 0-25°C over 1h in the presence of 10mol% $Pd(OAc)_2$ and 20mol% PPh_3 at which time hydrostannylation was judged complete by ¹H n.m.r. monitoring. In the case of (3a) and (3b) only the α -vinylstannanes (4a) and (4b) were detected whilst (3c) gave a 3:1 mixture of (4c) and another stannane presumed to be the (E)- β -vinylstannane.

Heating the reaction mixtures to 100°C induced cyclisation-intramolecular anion capture furnishing (5a), (5b) and (5c) in 67, 79 and 70% overall yield respectively from (3a-c) based on α -vinylstannane (Scheme 1).




Scheme 1

An analogous cascade, using the same conditions and catalyst system, was carried out on (6) which afforded (7) in 56% yield.



The foregoing results encouraged us to explore macrocyclisation cascades. A series of enynes (8a-c) was prepared as outlined in Scheme 2. Hydrostannylation of (8a-c) (0.05 molar solutions) was carried out as described previously and gave the α -vinylstannanes (9a-c) as the only detectable (^1H n.m.r.) products. Heating the diluted reaction mixture (5×10^{-3} molar solutions) at 100°C (toluene) gave the macrocyclic spirocycles (10a-c) in 59, 44 and 53% overall yields from (8a-c) respectively (Scheme 2). The yield of (10c) was raised from 28% to 59% by syringe pump addition of the solution of (9c) to a solution of an additional charge of catalyst [10mol% $\text{Pd}(\text{OAc})_2$, 20mol% PPh_3] in toluene at 100°C over 20h.

A second series of macrocyclic spirocycles (12) has been prepared from (11). In this case the preferred catalyst comprised 5mol% Pd₂(dba)₃ and 20mol% tri(2-furyl)phosphine. Hydrostannylation at 0°C was less selective than previously and afforded 2:1 mixtures of the α- and β-vinylstannanes. Bis-cyclisation proceeded smoothly at 100°C (5x10⁻³ molar solutions) over 12h to afford (12) in 50-53% yield (Table 1).



Scheme 2. (i) HO(CH₂)_nOH, NaH, DMF, 0°C. (ii) HC≡CCH₂NHSO₂Ph, ADDP, PBu₃, Toluene, 25°C





Table 1. Biscyclisation-intramolecular anion capture of (11) to (12).

n	macrocyclic ring	Yield(5) ^a
3	12	53 (71)
4	13	52 (70)
5	14	52 (70)
6	15	53 (71)
7	16	50 (67)
8	17	53 (71)

a. Isolated yields. Yields in brackets are corrected for the α-/β-vinylstanne ratio.

A series of bridged-ring forming macrocyclisations employing (13a-c) afforded the desired products (14a-c) in moderate yield. In these cases the preferred catalyst system comprised 10mol% Pd₂(dba)₃, 80mol% tri(2-furyl)phosphine and LiCl(1-3eq). The bis-cyclisations employed 5x10⁻³ molar solutions and were complete in 2-4h at 110°C. Monitoring the hydrostannylation reactions (0.05 molar solutions) by ¹H n.m.r. indicated only the desired α-vinylstannane was being formed.



The reactions described in this paper demonstrate the synthetic potential of hydrostannylation-polycyclisation-intramolecular anion capture cascades and research is continuing on these and related processes.

We thank the EPSR, the Spanish Government (MEC), the E.U. Human Capital and Mobility Scheme, Leeds University and Organon Laboratories for support.

References

1. Stille, J.K.; Tanaka, M.J.; *J. Am. Chem. Soc.*; 1987, **109**, 3785-3786; Kalivretenos, A.; Stille, J.K.; Hegedus, L.S.; *J. Org. Chem.*, 1991, **56**, 2883-2894.
2. Hiram, M.; Fujiwara, K.; Shigematu, K.; Fukazawa, Y.; *J. Am. Chem. Soc.*, 1989, **111**, 4120-4122.
3. Baldwin, J.E.; Adlington, R.M.; Ramcharitar, S.H.; *Tetrahedron*, 1992, **48**, 2957-2976; Pattenden, G.; Thom, S.M.; *Synlett.*, 1993, 215-216; Nicolaou, K.C.; Chakraborty, T.K.; Piscopio, A.D.; Minowa, N.; Bertino, P.; *J. Am. Chem. Soc.*, 1993, **115**, 4419-4420; Shair, M.D.; Yoon, T.; Danishefsky, S.J.; *J. Org. Chem.*, 1994, **59**, 3755-3757; Barrett, A.G.M.; Boys, M.L.; Boehm, T.L.; *J. Chem. Soc., Chem. Commun.*, 1994, 1881-1882.
4. Grigg, R.; Redpath, J.; Sridharan, V.; Wilson, D.; *Tetrahedron Lett.*, 1994, **35**, 4429-4432 and 7661-7764; Grigg, R.; Putnikovic, B.; Urch, C.J.; *ibid.*, 1996, **37**, 695-698; for reviews see: Grigg, R.; *J. Heterocyclic Chem.*, 1994, **31**, 631-639; Grigg, R.; Sridharan, V.; *Comprehensive Organometallic Chemistry*, edit. Abel, E.; Wilkinson, G.; Stone, F.G.A.; Pergamon Press, 2nd Edition, 1995, vol. 12, chp. 3.6.
5. Guibe, F.; Balavoine, G.; Zhang, H.X.; *J. Org. Chem.*, 1990, **55**, 1857-1867; Ito, Y.; Inouye, M.; Yokata, H.; Murakami, M.; *J. Org. Chem.*, 1990, **55**, 2567-2568.

(Received in UK 1 March 1996; revised 1 May 1996; accepted 3 May 1996)