

## An Intermolecular Michael Addition of Benzene


Tai-Chi Wang,<sup>a,b</sup> Yeh-Long Chen,<sup>a</sup> Kuan-Han Lee,<sup>b</sup> and Cherng-Chyi Tzeng<sup>\*a</sup>

<sup>a</sup>School of Chemistry, Kaohsiung Medical College, Kaohsiung City 807, Taiwan, R. O. C.


<sup>b</sup>Department of Pharmacy, Tajen Junior College of Pharmacy, Pingtung, Taiwan, R. O. C.

**Abstract:** The first intermolecular Michael addition of benzene leading to the formation of 3,3-diphenylpropionanilide is described. 2-Methoxyaniline was reacted with cinnamoyl chloride to give 2-methoxycinnamanilide (1) which was treated with aluminum chloride in benzene at 80°C to afford 2'-hydroxy-3,3-diphenylpropionanilide (4) in an 85% overall yield. Accordingly, 4'-hydroxy-2'-methyl-3,3-diphenylpropionanilide (6) was prepared from 4-methoxy-2-methylcinnamanilide (5) in 76% yield.  
 Copyright © 1996 Elsevier Science Ltd

A number of 2(1*H*)-quinolinone derivatives have been synthesized and evaluated for their cardiovascular activities.<sup>1-5</sup> Recently, Fujioka *et al.* described the preparation of 6-hydroxy-8-methyl-2(1*H*)-quinolinone, a key precursor for many novel positive inotropic agents, from 4-methoxy-2-methylaniline *via* Schotten-Baumann reaction and an intramolecular Friedel-Crafts cyclization.<sup>2</sup> Although the cyclization mechanism which included an unusual dearylation, was previously proposed by Manimaran *et al.*,<sup>6</sup> we believe that dearylation occur *via* the enolate 2. We have been interested in the preparation of hydroxycoumarin derivatives and examining their antiplatelet activity.<sup>7,8</sup> In an effort to expand these studies, *i.e.*, to synthesize their bioisosteric isomers, hydroxy-2(1*H*)-quinolinones, Fujioka's procedures were followed.<sup>2</sup> 2-Methoxyaniline was reacted with cinnamoyl chloride to give 2-methoxycinnamanilide (1) in 98% yield. Cyclization of 1 with aluminum chloride in chlorobenzene at 120°C afforded the desired 8-hydroxy-2(1*H*)-quinolinone (3) in 76% yield (Scheme 1).



To optimize the cyclization reaction, chlorobenzene was replaced with benzene as the reaction solvent to provide a relatively mild condition (refluxed at 80°C). The  $^1\text{H}$  NMR spectrum of the sole product isolated in this reaction showed a doublet at  $\delta$  3.22 ppm, a triplet at  $\delta$  4.56 ppm, and a multiplet at  $\delta$  6.69–7.63 ppm corresponding to  $\text{CH}_2$ ,  $\text{CH}$ , and aromatic protons respectively. The  $^{13}\text{C}$  NMR spectrum supported the  $^1\text{H}$  NMR spectrum in confirming the presence of a methylene carbon resonance appeared at  $\delta$  41.51 ppm and a tertiary carbon resonance at  $\delta$  46.87 ppm. The intermolecular Michael addition of **1** with benzene to give 2'-hydroxy-3,3-diphenylpropionanilide (**4**) seems to be a reasonable deduction. However, we were reluctant to make this critical structural assignment founded only on this evidence and therefore we sought a more definitive answer; an X-ray crystallographic analysis. A view of a single molecule of the crystal revealed that an intermolecular Michael addition occurred in which the benzene functions as a Michael donor and **1** as a Michael acceptor leading to the formation of **4** instead of the expected **3**. In order to establish and to further confirm this novel addition, 4-methoxy-2-methylaniline was converted into 4-methoxy-2-methylcinnamanilide (**5**) which was treated with aluminium chloride in refluxed benzene. 4'-Hydroxy-2'-methyl-3,3-diphenylpropionanilide (**6**) was obtained in 76% yield (Scheme 2). The structure of **6** was also established by  $^1\text{H}$  NMR spectrum [  $\delta$  1.78 (s, 3H,  $\text{CH}_3$ ), 3.03 (d, 2H,  $\text{CH}_2$ ), 4.54 (t, 1H,  $\text{CH}$ ), 6.44–7.36 (m, 13H, Ar-H), 9.10, 9.15 (2H, NH & OH)],  $^{13}\text{C}$  NMR spectrum [  $\delta$  17.55, 41.43, 47.06, 112.37, 116.39, 126.14, 126.94, 127.49, 127.60, 128.31, 134.06, 144.18, 154.85, 168.99] and elemental analyses.



Scheme 2

**Acknowledgement:** We thank the National Science Council of the Republic of China for the support of this study and Professor Jack E. Baldwin, F.R.S. of the Dyson Perrins Laboratory for valuable discussions.

#### References and Notes

- Uno, T.; Ozeki, Y.; Koga, Y.; Chu, G.-N.; Okada, M.; Tamura, K.; Igawa, T.; Unemi, F.; Kido, M.; Nishi, T. *Chem. Pharm. Bull.* **1995**, *43*, 1724-1733.
- Fujioka, T.; Teramoto, S.; Mori, T.; Hosokawa, T.; Sumida, T.; Tominaga, M.; Yabuuchi, Y. *J. Med. Chem.* **1992**, *35*, 3607-3612.
- Tominaga, M.; Yo, E.; Ogawa, H.; Yamashita, S.; Yabuuchi, Y.; Nakagawa, K. *Chem. Pharm. Bull.* **1984**, *32*, 2100-2110.
- Nishi, T.; Tabusa, F.; Tanaka, T.; Shimizu, T.; Nakagawa, K. *Chem. Pharm. Bull.* **1985**, *33*, 1140-1147.
- Tomigawa, M.; Ogawa, H.; Yo, E.; Yamashita, S.; Yabuuchi, Y.; Nakagawa, K. *Chem. Pharm. Bull.* **1988**, *35*, 3699-3704.
- Manimaran, T.; Thiruvengadam, T.K.; Ramakrishnan, V.T. *Synthesis* **1975**, 739-741.
- Chen, Y.-L.; Wang, T.-C.; Lee, K.-H.; Chang, Y.-L.; Teng, C.-M.; Tzeng, C.-C. *Helv. Chim. Acta* in press.
- Chen, Y.-L.; Wang, T.-C.; Liang, S.-C.; Teng, C.-M.; Tzeng, C.-C. *Chem. Pharm. Bull.* in press.