



A General and Convenient Synthetic Method of Geometrically Pure (Z)-1-Bromo-1-alkenes

Jun'ichi Uenishi,* Reiko Kawahama, Yasuhiko Shiga, and Osamu Yonemitsu


Department of Chemistry, Faculty of Science, Okayama University of Science
Ridaicho, Okayama 700, Japan

Jiro Tsuji

Department of Chemical Technology, Kurashiki University of Science and the Arts
Tsurajima-cho, Kurashiki, Okayama 712, Japan



























Abstracts: Palladium catalyzed hydrogenolysis of 1,1-dibromo-1-alkenes by tributyltin hydride proceeds smoothly to give (Z)-1-bromo-1-alkenes with excellent stereoselectivity in good yields. Dibromomethylenation of aldehydes by a combination of CBr_4 and Ph_3P in methylene chloride and the successive hydrogenolysis affords (Z)-1-bromo-1-alkenes in one-pot.
Copyright © 1996 Elsevier Science Ltd

(Z)-1-Halo-1-alkene is an important building block for stereospecific synthesis of organic compounds possessing a Z-olefin unit in the molecules.¹ To date, (Z)-1-halo-1-alkenes have been prepared using several methods, including *cis* hydrogenation of 1-halo-1-alkynes with diimine,² haloalkenylation of aldehydes with Wittig type reagents,³ stereospecific substitution of Z- or E-metalloalkenes with halogen-groups,⁴ and others.⁵ Sometimes, however, they are unsatisfactory, having low reactivity, poor stereoselectivity, or undesired side reactions including over-reduction. Herein, we report a very simple and reliable method for preparing geometrically pure (Z)-1-bromo-1-alkenes from 1,1-dibromo-1-alkenes. The method can be expressed in the following general scheme.



When β,β -dibromostyrene (**1a**) was treated with tributyltin hydride (1.05~1.15 eq) in the presence of tetrakis(triphenylphosphine)palladium (4 mol%) in benzene⁶ at room temperature, (Z)- β -bromostyrene (**2a**) was obtained in 76% yield. The reaction was completed in 15 min and none of the E-isomer was detected in the crude product.⁷ The starting 1,1-dibromoalkenes, **1a-1m**, were prepared from the corresponding aldehydes in 70 to 98% yields by a standard method.⁸ Results of the hydrogenolysis for **1a-1m**, are summarized in Table 1.^{9,10} *Ortho* and *para* methyl substituted styrenes, **1b** and **1c**, gave the corresponding (Z)-bromostyrenes, **2b** and **2c**, in both 79% yields. An acetoxy group substituting on the aromatic ring did not influence the reaction, thus, 4-acetoxy-(Z)- β -bromostyrene (**2d**) was obtained in 83% yield. Even in the presence of a phenol group,

Table 1. Palladium Catalyzed Hydrogenolysis of 1,1-Dibromoalkenes with Bu₃SnH

Entry	1,1-Dibromoalkene		Z-Bromoalkene ^a	Yield ^b (%)
1		1a		2a 76
2		1b		2b 79
3		1c		2c 79
4		1d		2d^c 83
5		1e		2e 82
6		1f		2f^d 90
7		1g		2g 61
8		1h		2h 79
9		1i		2i 84
10		1j		2j 70
11		1k		2k 77
12		1l		2l 56
13		1m		2m 80

^a Geometric purities were determined to be greater than 98% by ¹H NMR. ^b Isolated yields.^c mp 44–46°C. ^d mp 49–50°C.

In conclusion, we have succeeded in the stereoselective hydrogenolysis of 1,1-dibromoalkenes leading to *Z*-bromo-1-alkenes exclusively. The resulting (*Z*)-bromoalkenes would be valuable compounds for stereospecific synthesis of polyenes or other systems possessing a *Z*-olefin unit in the molecules.

Acknowledgment

We appreciate the financial support of a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan.

References and Notes

- 1, (a) Tsuji, J. *Palladium Reagents and Catalysts*; John Wiley & Sons Ltd.: Chichester, **1995**. (b) Knight, D. W. In *Comprehensive Organic Synthesis*; Trost, B. M. and Fleming, I., Eds.: Pergamon Press: New York, **1991**, Vol. 3, pp 481-520.
- 2, Kluge, A. F.; Untch, K. G.; Fried, J. H. *J. Am. Chem. Soc.* **1972**, *94*, 9256-9258.
- 3, (a) Bestmann, H. J.; Rippel, H. C.; Dostalek, R. *Tetrahedron Lett.* **1989**, *30*, 5261-5262. (b) Matsumoto, M.; Kuroda, K. *Tetrahedron Lett.* **1980**, *21*, 4021-4024.
- 4, (a) Miller, R. B.; Reichenbach, T. *Tetrahedron Lett.* **1974**, 543-546. (b) Brown, H. C.; Hamaoka, T.; Ravindran, N. *J. Am. Chem. Soc.* **1973**, *95*, 6456-6457.
- 5, (a) Alexakis, A.; Duffault, J. M. *Tetrahedron Lett.* **1988**, *29*, 6243-6246. (b) Brown, H. C.; Blue, C. D.; Nelson, D. J.; Bhat, N. G. *J. Org. Chem.* **1989**, *54*, 6064-6067. (c) Harada, T.; Katsuhira, T.; Hara, D.; Kotani, Y.; Maejima, K.; Kaji, R.; Oku, A. *J. Org. Chem.* **1993**, *58*, 4897-4907.
- 6, Other solvents, such as THF, ether, toluene and CH₂Cl₂ are also suitable for the hydrogenolysis.
- 7, For (*E*)- and (*Z*)- β -bromostyrenes; see Miller, R. B.; McGarvey, G. *J. Org. Chem.* **1978**, *43*, 4424-4431.
- 8, (a) Corey, E. J.; Fuchs, P. L. *Tetrahedron Lett.* **1972**, 3769-3772. (b) Ramirez, F.; Desai, N. B.; McKelvie, N. *J. Am. Chem. Soc.* **1962**, *84*, 1745-1747.
- 9, A typical experimental procedure is as follows: To a stirred solution of dibromoalkene, **1** (1 mmol) and Pd(PPh₃)₄ (0.04 mmol) in anhydrous benzene (7 mL) was added Bu₃SnH (1.1 mmol) in anhydrous benzene (3 mL) under Ar atmosphere. After the mixture was stirred for 15 to 60 min at room temperature, an extractive work up with hexane or benzene and a purification by alumina column chromatography gave **2**.
- 10, The stereochemistries of all the products were confirmed by comparison with previous data reported in the literature, or determined by a coupling constants of the olefinic protons in proton nmr spectra.
- 11, Roush, W. R.; Moriarty, K. J.; Brown, B. B. *Tetrahedron Lett.* **1990**, *31*, 6509-6512.
- 12, Minato, A.; Suzuki, K.; Tamao, K. *J. Am. Chem. Soc.* **1987**, *109*, 1257-1258.
- 13, For example, the reaction of **1a** and **1i** gave a 1 : 1 mixture of (*E*)- and (*Z*)-bromoalkenes along with a mixture of other complex products.
- 14, The desired alkenes were obtained in 35~37% yields, although these authors succeeded in stereospecific hydrogenolysis of 1-iodo-1-alkenes with tributyltin hydride using a slow addition technique. See, Taniguchi, M.; Takeyama, Y.; Fugami, K.; Oshima, K.; Utimoto, K. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 2593-2595.
- 15, Theoretically, five equivalents of tributyltin hydride are totally required in this case; three for the reaction with Ph₃PBr₂ (1.5 eq), one for the reaction with the remaining Ph₃P=CBr₂ (0.5 eq), and an additional equivalent for the hydrogenolysis.

(Received in Japan 28 June 1996; revised 18 July 1996; accepted 23 July 1996)