



## Chemoselective Reduction of Highly-functionalized Azidopyridazines to Corresponding Aminopyridazines Using Fe/NH<sub>4</sub>Cl in Organic Solvent-Water Two-phase Solution

Su-Dong Cho, Woo-Yong Choi, Sang-Gyeong Lee, Yong-Jin Yoon\*, and Sung Chul Shin\*


Department of Chemistry, Gyeongsang National University, Chinju, 660 - 701, Korea

**Abstract;** Highly functionalized azidopyridazines can be reduced chemoselectively to the corresponding amines in excellent yields.

Copyright © 1996 Elsevier Science Ltd

The reduction of azides to amines is a synthetically important process and its value has been well established.<sup>1</sup> Of the convenient methods available for the conversion of azides to amines, the most popular involve catalytic hydrogenation,<sup>2</sup> Staudinger condition,<sup>3</sup> hydrogen sulfide,<sup>4</sup> thiol under basic condition,<sup>5</sup> metal hydride,<sup>6</sup> and tetrathiomolybdate.<sup>7</sup> Most of these methods, however, often suffer from one or more limitations with respect to general applicability, selectivity, ready availability, operational convenience.<sup>6c</sup> Fe/NH<sub>4</sub>Cl in a mixed solution of ethyl acetate and water has been used for reducing nitroarenes,<sup>8</sup> but little attention has been focused on reduction of azides under this condition. During the course of our synthetic study on diazine-fused nitrogen-heterocycles,<sup>9</sup> we found that azidopyridazines substituted by various types of functional groups can be reduced chemoselectively and efficiently to the corresponding amines under this condition.

Table 1.



Entry.	X	R	yield (%) <sup>a</sup>	mp (°C) <sup>b</sup>
a	Cl	CH <sub>2</sub> COCH <sub>3</sub>	93	194-195
b	Br	CH <sub>2</sub> COCH <sub>3</sub>	95	192-193
c	Cl	CH <sub>2</sub> COOCH <sub>3</sub>	97	222-223
d	Br	CH <sub>2</sub> C(=NOH)CH <sub>3</sub>	92	204-205
e	Br	CH <sub>2</sub> CH <sub>2</sub> CN	98	219-220
f	Cl	CH <sub>2</sub> C[O(CH <sub>2</sub> ) <sub>2</sub> O]CH <sub>3</sub>	97	229-230
g	Cl	CH <sub>2</sub> (CH <sub>2</sub> ) <sub>3</sub> O-COC <sub>6</sub> H <sub>5</sub>	96	137-138

<sup>a</sup>Purified yields. <sup>b</sup>Uncorrected.

The results are summarized in Table 1. Azidopyridazines **1** bearing various functional groups<sup>10</sup> were simply stirred with the excess amount of Fe powder (3 equiv.) and NH<sub>4</sub>Cl (5 equiv.) in the two phase solution of ethyl acetate and water at room temperature to give amines **2** in excellent yields. Amines **2**<sup>11</sup> were purified as free amines by filtering the Fe powder, evaporating the organic phase, and recrystallizing or chromatographing the crude products. The reactions described here proceed in chemoselective fashion, keeping a variety of functional groups such as halogen, carbonyl, nitrile, oxime, ester, acetal and acyloxy, intact and complete within 2 hr at room temperature. The reactions can also be performed using other common organic phase such as benzene, chloroform or methanol in place of ethyl acetate to give the products in comparative yields. The present  $\alpha$ -amino- $\beta$ -halopyridazines **2** should find their way as useful synthons for the construction of a variety of fused nitrogen-heterocycles by a<sup>2</sup>, d<sup>0</sup> / a<sup>n</sup>, d<sup>0</sup> combination.<sup>12</sup>

This idea and the expanded results of this novel method to other heterocyclic azides will be published elsewhere.

### References and Notes;

- (a) Patai, S. Ed., "The Chemistry of the Azido group" Interscience Publishers, New York, 1971. (b) Scriven, E. F. V. Ed., "Azides and Nitrenes Reactivity and Utility" Academic Press, Inc., New York, 1984.
- (a) Malik, A. A.; Preston S. B.; Archibald, T. G.; Cohen, M. P.; Baum, K. *Synthesis*, **1989**, 450. (b) Sharma, G. V. M.; Chandrasekhar, S. *Synth. Commun.* **1989**, *19*, 3289.
- Koziara, A.; Zwierzak, A. *Synthesis*, **1992**, 1063.
- Rao, H. Surya Prakash; Doss, S. D. *Sulfur Lett.* **1992**, *14*, 61.
- Becher, J.; Pluta, K.; Krake, N.; Brondum, K.; Christensen, N. J.; Vinader, M. V. *Synthesis*, **1989**, 530. (b) Bartra, M.; Romea, P.; Urpi, F.; Vilarrasa, J. *Tetrahedron* **1990**, *46*, 587.
- (a) Boyer, J. H.; Ellzey Jr., S. E. *J. Org. Chem.* **1958**, *23*, 127 (b) Kabalka, G. W.; Wadgaonkar, P. P.; Chatla, N. *Synth. Commun.* **1990**, *20*, 293 (c) Ranu, B. C.; Sarkar, A.; Chakraborty, R. J. *Org. Chem.* **1994**, *59*, 4114.
- Ramesha, A. R.; Chandrasekaran, S. *J. Org. Chem.* **1995**, *60*, 7682.
- Ramadas, Krishnamurthy., Srinivasan, Natarajan. *Synth. Commun.* **1992**, *22*, 3189.
- (a) Lee, S.-G.; Choi, S. Y.; Yoon, Y.-J. *J. Heterocyclic Chem.* **1992**, *29*, 1409. (b) Shin, S. C.; Jeon, D. J.; Jang, K. A.; Lee, Y. Y. *Bull. Kor. Chem. Soc.* **1994**, *15*, 517.
- The starting azides used were prepared by nucleophilic displacement using sodium azide and the corresponding halides. The details with respect to the preparation of starting azides will be published elsewhere.
- For **2a**; white needles; IR (KBr) 3460, 3344, 3226(NH<sub>2</sub>), 1744(C=O), 1660(C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 2.10(s, 3H), 4.80(s, 2H), 6.77(br. s, NH<sub>2</sub>), 7.56(s, 1H) ppm; <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) 26.9, 60.1, 79.0, 130.0, 145.6, 156.9, 201.7 ppm; MS m/z (%) : 201 (10, M<sup>+</sup>), 159 (23, M<sup>+</sup>-42), 131 (7, M<sup>+</sup>-70), 32 (100, M<sup>+</sup>-169) : **2b**; white needles; IR (KBr) 3456, 3292, 3178(NH<sub>2</sub>), 1734(C=O), 1632(C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 2.12(s, 3H), 4.83(s, 2H), 6.80(br. s, NH<sub>2</sub>), 7.54(s, 1H) ppm; <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) 26.6, 60.2, 79.7, 129.8, 149.6, 157.3, 202.2 ppm; MS m/z (%) : 246 (14, M<sup>+</sup>), 203 (37, M<sup>+</sup>-43), 175 (10, M<sup>+</sup>-71), 68 (48, M<sup>+</sup>-178), 43 (100, M<sup>+</sup>-203) : **2c**; white needles; IR (KBr) 3436, 3332, 3218(NH<sub>2</sub>), 1754(C=O), 1658(C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 3.66(s, 3H), 4.75(s, 2H), 6.88(br. s, NH<sub>2</sub>), 7.62(s, 1H) ppm; <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) 51.9, 52.5, 79.0, 130.2, 145.7, 156.9, 168.2 ppm; MS m/z (%) : 217 (43, M<sup>+</sup>), 185 (25, M<sup>+</sup>-32), 158 (74, M<sup>+</sup>-59), 32 (100, M<sup>+</sup>-185) : **2d**; white needles; IR (KBr) 3444, 3322, 3200, (NH<sub>2</sub>), 1648(C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 1.67 & 2.14(s, 3H, syn & anti), 4.65 & 4.85(s, 2H, syn & anti), 6.75 & 6.80(s, NH<sub>2</sub>, syn & anti), 7.52(s, 1H), 10.71(s, OH) ppm; <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) 11.6, 26.9, 53.7, 60.3, 79.0, 129.4, 129.5, 147.2, 147.5, 150.9, 157.1 ppm; MS m/z (%) : 261 (48, M<sup>+</sup>), 246 (64, M<sup>+</sup>-15), 203 (64, M<sup>+</sup>-58), 122 (57, M<sup>+</sup>-139), 95(58, M<sup>+</sup>-166), 68(65, M<sup>+</sup>-193), 40(100, M<sup>+</sup>-221) : **2e**; white needles; IR (KBr) 3462, 3364, 3242(NH<sub>2</sub>), 2250(CN), 1646(C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 2.93(t, 2H), 4.20(t, 2H), 6.84(br. s, NH<sub>2</sub>), 7.56(s, 1H) ppm; <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) 16.2, 46.4, 79.0, 118.2, 129.7, 147.4, 156.9 ppm; MS m/z (%) : 242 (40, M<sup>+</sup>), 204(27, M<sup>+</sup>-38), 189 (72, M<sup>+</sup>-53), 132 (29, M<sup>+</sup>-110), 32 (100, M<sup>+</sup>-210), 40 (100, M<sup>+</sup>-203) : **2f**; white needles; IR (KBr) 3422, 3326, 3208(NH<sub>2</sub>), 1658(C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 1.25(s, 3H), 3.88(m, 4H), 4.06(s, 2H), 6.77(br. s, NH<sub>2</sub>), 7.58(s, 1H) ppm; <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) 23.0, 53.7, 64.1, 107.5, 129.4, 145.1, 157.2 ppm; Anal. Calcd. for C<sub>9</sub>H<sub>12</sub>N<sub>3</sub>O<sub>3</sub>Cl: C, 44.0; H, 4.92; N, 17.10. Found: C, 43.90 H, 4.79 N, 17.18 : **2g**; pale yellow needles; IR (KBr) 3502, 3320, 3204(NH<sub>2</sub>), 1740(C=O), 1658(C=O) cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) 1.75(m, 4H), 4.03(t, 2H), 4.27(t, 2H), 6.73(br. s, NH<sub>2</sub>), 7.49-7.96(m, 6H) ppm; <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) 24.6, 25.2, 49.9, 64.1, 128.5, 128.9, 129.5, 129.7, 133.0, 145.2, 156.9, 165.6 ppm; Anal. Calcd. for C<sub>15</sub>H<sub>16</sub>N<sub>3</sub>O<sub>3</sub>Cl: C, 55.99; H, 5.01; N, 13.06. Found: C, 56.15 H, 4.91 N, 13.24
- Seebach, D. *Angew. Chem. Int. Ed. Engl.* **1979**, *18*, 239

(Received in Japan 26 June 1996; revised 31 July 1996; accepted 6 August 1996)