

Reaction of Indole-2,3-dicarboxylic Anhydride with (3-Bromo-4-pyridyl)triisopropoxytitanium: Synthesis of Ellipticine

Yasuyoshi Miki,* Yoshitaka Tada, Norihide Yanase, Hiroko Hachiken,
and Ko-ichi Matsushita

Faculty of Pharmaceutical Sciences, Kinki University, 3-4-1 Kowakae, Higashi-Osaka 577, Japan

Abstract: *N*-Benzylindole-2,3-dicarboxylic anhydride (**1**) was reacted with (3-bromo-4-pyridyl)triisopropoxytitanium to give 2-(3-bromoisonicotinoyl)indole-3-carboxylic acid (**2**) as the sole product in high yield, which could be converted to ellipticine in six steps. Copyright © 1996 Elsevier Science Ltd


Ellipticine, 5,11-dimethyl-6*H*-pyrido[4,3-*b*]carbazole, has potent antitumor activity¹ and many useful methods for its synthesis have been reported.^{2,3} Although the Diels-Alder reaction of 4*H*-furo[3,4-*b*]indole,⁴ pyrrolo[3,4-*b*]indole,⁵ and pyrano[3,4-*b*]indol-3-one⁶ with 3,4-pyridyne is the attractive synthetic methods of ellipticine, it produced a mixture of ellipticine and isoellipticine. Reaction of 4*H*-furo[3,4-*b*]indole with 5,6-dihydro-2-pyridinone afforded a sole adduct, but the transformation of the adduct to ellipticine resulted in low yield.⁷ However, Gribble reported a useful synthesis of ellipticine by using the reaction of 2-lithio-1-(phenylsulfonyl)indole with 3,4-pyridinedicarboxylic anhydride.⁸

Recently we have reported a synthesis of murrayaquinone-A by using the selective reduction of *N*-benzylindole-2,3-dicarboxylic anhydride (**1**), in which the 2-carbonyl group is more reactive toward the nucleophile than the 3-carbonyl group,⁹ following cycloaddition reaction of the corresponding furo[3,4-*b*]indole with methyl acrylate.¹⁰ The nucleophilic reaction of *ortho*-halolithiopyridines is a convenient method for the preparation of numerous *ortho*-substituted halopyridines¹¹. However, the reaction of enolizable ketones or unstable compounds toward a base with *ortho*-halolithiopyridines gave sometimes unacceptable results.¹² Seebach¹³ reported that phenyltitanium compounds reacted with aldehydes to give the corresponding products in high yield, but the reactivity of *ortho*-haloltitaniumpyridines has not been extensively studied.

In this paper we report a simple and useful synthesis of ellipticine by regioselective reaction of **1** with a 4-pyridyltitanium derivative.

N-Benzylindole-2,3-dicarboxylic anhydride (**1**) reacted with a 3-bromo-4-pyridyltitanium compound (prepared from 3-bromo-4-lithiopyridine¹⁴ and ClTi(OPr^{*i*})₃¹⁵ in THF -96°C) to afford 2-acylindole-3-carboxylic acid (**2**) in 86% yield.¹⁶ Decarboxylation and debenylation (AlCl₃ in anisole,¹⁷ 100°C) of **2** furnished ketone (**3**), which was changed by Wittig reaction, followed by catalytic reduction to 1-(3-bromo-4-

pyridyl)-1-(2-indolyl)ethane (4). Treatment of 4 with (1-ethoxyvinyl)tributyltin in the presence of tetrakis(triphenylphosphine)palladium(0) in refluxing toluene gave the corresponding ethoxyvinyl derivative (5), which was converted to ellipticine¹⁸ in 87% yield by treatment with 10% hydrochloric acid in tetrahydrofuran.



References and Notes

- Dalton, L. K.; Demerac, S.; Elmes, B. C.; Loder, J. W.; Swan, J. M.; Teitei, T. *Aust. J. Chem.*, **1967**, *20*, 2715.
- For reviews, see Gribble, G. W. *Synlett*, **1991**, 289; Gribble, G. W. *Advances in Heterocyclic Natural Product Synthesis*; Vol. 1, p 43, Pearson, W., Ed., Jai Press Inc., London, **1990**; Kansal, V. K.; Potier, P. *Tetrahedron*, **1986**, *42*, 2389; Gribble, G. W.; Saulnier, M. G. *Heterocycles*, **1985**, *23*, 1277; Hewlins, M. J. E.; O-Campos, A.-M.; Shannon, P. V. R. *Synthesis*, **1984**, 289; Sainsbury, M. *Synthesis*, **1977**, 437.
- Hibino, S.; Sugino, E. *J. Heterocyclic Chem.*, **1990**, *27*, 1751; Modi, S. P.; Michael, M. A.; Archer, S. *Tetrahedron*, **1991**, *47*, 6539.
- Gribble, G. W.; Saulnier, M. G.; Sibi, M. P.; O-Nutaitis, J. A. *J. Org. Chem.*, **1984**, *49*, 4518.
- Sha, C.-K.; Yang, J.-F. *Tetrahedron*, **1992**, *48*, 10645.
- May, C.; Moody, C. J. *J. Chem. Soc., Perkin Trans. 1*, **1988**, 247.
- Davis, D. A.; Gribble, G. W. *Tetrahedron Lett.*, **1990**, *31*, 1081; Gribble, G. W.; Keavy, D. J.; Davis, D. A.; Saulnier, M. G.; Pelcman, B.; Barden, T. C.; Sibi, M. P.; Olson, E. R.; BelBruno, J. J. *J. Org. Chem.*, **1992**, *57*, 5878.
- Gribble, G. W.; Saulnier, M. G.; O-Nutaitis, J. A.; Ketcha, D. M. *ibid.*, **1992**, *57*, 5891.
- Jones, R. A. *Comprehensive Heterocyclic Chemistry*; Vol. 4, p 288, Bird, C. W.; Cheeseman, G. W. H., Ed., Pergamon Press, Oxford, **1984**.
- Miki, Y.; Hachiken, H. *Synlett*, **1993**, 333.
- Snieckus, V. *Chem. Rev.*, **1990**, *90*, 879; Rewcastle, G. W.; Katritzky, A. R. *Advances in Heterocyclic Chemistry*, Vol. 56, p 155, Katritzky, A. R., Ed., Academic Press Inc., **1993**.
- Gu, Y. G.; Bayburt, *Tetrahedron Lett.*, **1996**, *37*, 2565.
- Weidmann, B.; Wilder, L.; Olivero, A. G.; Maycock, C. D.; Seebach, D. *Helv. Chim. Acta*, **1981**, *64*, 357.
- Effenberger, F.; Daub, W. *Chem. Ber.*, **1991**, *124*, 2119.
- Reetz, M. T. *Organometallics in Synthesis*; John Wiley & Sons, p 195, Schlosser, M., Ed., **1994**.
- Reaction of the anhydride (1) with 3-bromo-4-lithiopyridine gave 2-acylindole-3-carboxylic acid (2) in 42% yield.
- Murakami, Y.; Watanabe, T.; Kobayashi, A.; Yokoyama, Y. *Synthesis*, **1984**, 738; Watanabe, T.; Kobayashi, A.; Nishiura, M.; Takahashi, H.; Usui, T.; Kamiyama, I.; Mochizuki, N.; Noritake, K.; Yokoyama, Y.; Murakami, Y. *Chem. Pharm. Bull.*, **1991**, *39*, 1152.
- Ellipticine: mp $>300^\circ\text{C}$ (MeOH) (lit.,⁶ 312-314°C); $^1\text{H-NMR}$ (DMSO-d_6) δ 2.80 (3H, s, 5- CH_3), 3.27 (3H, s, 11- CH_3), 7.22-7.33 (1H, m, Ar), 7.49-7.61 (2H, m, Ar), 7.92 (1H, d, $J = 6$ Hz, H-4), 8.39 (1H, d, $J = 8$ Hz, H-10), 8.43 (1H, d, $J = 6$ Hz, H-3), 9.69 (1H, s, H-1), 11.25 (1H, s, NH); HRMS m/z calcd for $\text{C}_{17}\text{H}_{14}\text{N}_2$ (M^+) 246.1157, found 246.1185.

(Received in Japan 27 May 1996; revised 7 August 1996; accepted 10 September 1996)