



An Approach Towards the Synthesis of Oligomers Containing a *N*-2-Hydroxyethyl-aminomethylphosphonate Backbone: A Novel PNA Analogue

Alexander C. van der Laan^a, Roger Strömberg^b, Jacques H. van Boom^a and Esther Kuyl-Yeheskiely^{a*}

^aLeiden Institute of Chemistry, Gorlaeus Laboratories, P.O. Box 9502, 2300 RA Leiden, The Netherlands


^bKarolinska Institute, Department of Medical Biochemistry and Biophysics, S-17177, Stockholm, Sweden

Vladimir A. Efimov and Oksana G. Chakhmakheva

Shemyakin & Ovchinnikov Institute of Bioorganic Chemistry, Miklukho-Maklaya str. 16/10, Moscow V-437, 117871, Russia

Abstract: A convenient route to the preparation of 4-methoxy-1-oxido-pyridine-2-methyl *N*-2-(4,4'-dimethoxytrityloxy)ethyl-*N*-thymine-1-yl-aminomethylphosphonate (**1a**, **T**⁺) and the corresponding *N*⁴-benzoylcytosine-1-yl derivative (**1b**, **C**⁺) is reported. These PPNA monomers proved to be suitable building blocks in a solid-support synthesis of the tetradecameric fragment (C⁺T⁺T⁺C⁺T⁺T⁺T⁺C⁺T⁺C⁺T⁺)dT. Copyright © 1996 Elsevier Science Ltd

In the last decade much effort has been directed towards the design and synthesis of natural and modified nucleic acids¹ that bind specifically to genes at the mRNA (antisense) or double stranded DNA (antigene) level. Recently it was revealed² that stacking interactions and Watson-Crick base pairing, as in B-DNA, could be effectively mimicked by replacing the deoxyribose phosphate backbone in DNA by an achiral polyamide backbone comprising *N*-(2-aminoethyl)glycine repeating units. This so-called PNA forms




highly stable PNA-DNA(RNA) complexes in a sequence specific manner and may therefore present a promising lead to therapeutics targeting at specific genes. However, the lack of negative charge in PNA, which is one of the factors responsible for the stability of PNA-DNA(RNA) duplexes, leads to poor solubility in a physiological environment³. It occurred to us that replacement of the amido group in PNA by a charged phosphonate linkage would give a water soluble PNA analogue (*i.e.* PPNA).

We here report the preparation of the *N*-(thymine-1-yl)- and *N*-(*N*⁴-benzoylcytosine-1-yl)-*N*-2-hydroxyethyl-aminomethylphosphonate building units **1a-b**, the 4-methoxy-1-oxido-pyridine-2-methyl

(MOPM) group of which will facilitate the introduction of the phosphonate linkages^{4,5} in PPNA. The use of the PPNA building units **1a-b** is further illustrated in a solid-support synthesis of the tetradecameric fragment (C^T*T^T*T^C*T^T*T^T*T^C*T^C*T^T)dT **15**.

Scheme 1




Reagents and Conditions: (i) CH₂O (1.2 eq) in ethyl acetate, 15 min; (ii) diphenyl phosphite (1.0 eq), toluene, 75°C, 2 h; (iii) Bromoacetic anhydride (1.0 eq), *N*-Me-morpholine (1.0 eq) in toluene (80% based on **2**); (iv) Thymine (1.2 eq), DBU (1.2 eq) [or *N*⁴-benzoylcytosine (1.2 eq), NaH (1.2 eq)] in dimethylformamide, 75°C, 1 h (**6a**: 75%, **6b**: 65%); (v) 0.4 M DBU in CH₃CN/H₂O (95/5), 15-30 min (90-95%); (vi) MOPM-OH (2.0 eq), TPS-Cl (2.0 eq), 4-methoxy-1-oxido-pyridine (6.0 eq) in CH₃CN, 30 min (70-75%); (vii) TPS-Cl (1.5 eq) in CH₃CN/C₆H₅N (4/1, v/v), 1 min (75%).

The preparation of the required PPNA building units **1a-b** could be realized by the sequence of reactions depicted in Scheme 1. Condensation of 2-(4,4'-dimethoxytrityloxy)ethylamine (**2**)⁶ in ethyl acetate with a slight excess of formaldehyde gave, after workup, the crude imino derivative **3**. Treatment of **3** at elevated temperature with an equimolar amount of diphenyl phosphite⁷ led, as gauged by ³¹P-NMR spectroscopy, to a near quantitative formation of the diphenyl phosphonate derivative **4** ($\delta_p = 20.2$ ppm). Acylation of crude **4** with bromoacetic anhydride⁸ in the presence of *N*-methyl-morpholine yielded, after purification by flash chromatography, homogeneous **5**⁹. Reaction of the intermediate bromoacetyl derivative **5** with thymine in the presence of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) resulted, after purification, in the isolation of homogeneous **6a** (B^{*} = T)⁹ in 75% yield. Similarly, treatment of **5** with *N*⁴-benzoylcytosine, using sodium hydride as a base, gave the corresponding cytosinyl derivative **6b** (B^{*} = C^{Bz})⁹ in 65% yield. Transformation of **6a,b** into the PPNA building block units **1a,b**, carrying the catalytic 4-methoxy-1-oxido-pyridine-2-methyl phosphonate protecting group, entailed the following three-step procedure. Conversion¹⁰ of the individual diphenyl phosphonates **6a,b** under the influence of DBU-H₂O led to corresponding monophenyl phosphonates **7a,b**⁹. Condensation¹¹ of the latter compounds with 2-hydroxymethyl-4-methoxy-1-oxido-pyridine (MOPM-OH) under the agency of 2,4,6-triisopropylbenzenesulfonyl chloride (TPS-Cl) and

4-methoxy-1-oxido-pyridine led to compounds **8a,b**⁹. Finally, removal¹¹ of the phenyl phosphonate protecting group from both **8a,b** with DBU-H₂O proceeded smoothly to give, after purification, **1a**⁹ and **1b**⁹ in an overall yield of 65% and 60%, respectively.

Scheme 2




Prior to the intended solid-support synthesis of the tetradecameric fragment **15**, the rate of the phosphorylation of the thymidine derivative **9** with **1a** was monitored by ³¹P-NMR spectroscopy. It was established that the TPS-Cl mediated condensation went to completion within 1 min. Workup and purification gave the homogeneous dimer **10** ($\delta_p = 22.7$ and 23.1 ppm) in 75% yield. The favourable outcome of the latter condensation was a stimulus to assemble tetradecamer **15** using a fully automated DNA synthesizer. The assembly of the target oligomer **15** comprises (see Scheme 2) extension of the thymidine derivative **11**, immobilized to controlled pore glass *via* a succinyl linker, with the PPNA units **1a-b**¹². Thus, sequential elongation of immobilized **11** with the appropriate units **1a-b** following the stepwise protocol summarized in Table 1 afforded, after thirteen elongation cycles, the fully protected and immobilized fragment **12**. The coupling efficiency of each elongation cycle was higher than 96%, as gauged spectrophotometrically by the released DMT-cation. Immobilized **12** was deblocked and released from the solid-support by the following three-step procedure. Acidolysis of the DMT group (R^1) from **12**, and subsequent removal of the MOPM group (R^2) in **13** with neat piperidine⁵, led to partially protected and immobilized **14**. Finally, *N*-debenzoylation and release from the solid-support was effected by ammonolysis of **14**. Purification of the resulting crude product by ion-exchange chromatography (Q-Sepharose) and subsequent desalting (Sephadex G-25) gave tetradecameric fragment **15**, the homogeneity and identity of which was established by fast protein liquid chromatography (FPLC, Figure 1)¹³ as well as mass spectro-

Table 1: Chemical steps involved in each elongation cycle of PPNA

Step	Manipulation	Solvents and reagents ^a	Time (min)
1	Detritylation	3% CHCl ₂ COOH in CH ₂ Cl ₂	1.5
2	Wash	CH ₃ CN	3.0
3	Coupling	1a-b ^b , TPS-Cl ^c in CH ₃ CN/ C ₆ H ₅ N (4/1, v/v)	5.0
4	Wash	CH ₃ CN	30.0
5	Capping	Ac ₂ O/ <i>N</i> -Me-imidazole/Collidine /THF, (2/3/2/32, v/v/v/v)	0.5
6	Wash	CH ₃ CN	2.0

^a Reactions were performed on 28 mg (1 μ mole) of resin. ^b 0.08 M **1a** (or **1b**) in CH₃CN/C₆H₅N (4/1, v/v), 5 eq. ^c 0.25 M TPS-Cl in CH₃CN/C₆H₅N, (4/1, v/v), 15 eq.

Figure 1: FPLC pattern of purified **15**¹³



scopy (MALDI-TOF).

In conclusion, the successful assembly of tetradecameric fragment **15** presented in this paper may open the way for a general solid-support synthesis of homogeneous PPNA.

A full report on the solid-support synthesis and biochemical properties of homogeneous PPNA will be published in due course.

Acknowledgements: We wish to thank Solvay Duphar N.V., Weesp, The Netherlands for financial support.

References and notes

1. a) Milligan, J.F.; Matteucci, M.D.; Martin, J.C. *J. Med. Chem.* **1993**, *36*, 1923-1937. b) Hunziker, J.; Leumann, C. in "Modern Synthetic Methods 1995", Ernst, B.; Leumann, E. Eds., Verlag Helv. Chim. Acta, Basel, p. 331-417 and references cited therein.
2. a) Nielsen, P.E.; Egholm, M.; Berg, R.H.; Buchardt, O. *Science* **1991**, *254*, 1497-1500. b) Egholm, M.; Nielsen, P.E.; Buchardt, O.; Berg, R.H. *J. Am. Chem. Soc.* **1992**, *114*, 9677-9678.
3. a) Bergmann, F.; Bannwarth, W.; Tam, S. *Tetrahedron Lett.* **1995**, *36*, 6823-6826. b) Egholm, M.; Buchardt, O.; Nielsen, P.E.; Berg, R.H. *J. Am. Chem. Soc.* **1992**, *114*, 1895-1897.
4. Szabo, T.; Kers, A.; Stawinski, J. *Nucl. Acids Res.* **1995**, *23*, 893-900.
5. Efimov, V.A.; Buryakova, A.A.; Dubey, I.Y.; Polushin, N.N.; Chakhmakhcheva, O.G.; Ovchinnikov, Y.A. *Nucl. Acids Res.* **1986**, *14*, 6526-6540.
6. Compound **2** was readily prepared by tritylation of commercially available *N*-(hydroxyethyl)phtalimide with DMT-Cl (1.1 eq) in pyridine and subsequent treatment of the tritylated product with hydrazine (2.0 eq) in 80% overall yield.
7. De Lombaert, S.; Erion, M.D.; Tan, J.; Blachard, L.; El-Chehabi, L.; Ghai, R.D.; Sakane, Y.; Berry, C.; Trapani, A.I. *J. Med. Chem.* **1994**, *37*, 498-511.
8. Meltzer, P.C.; Liang, A.Y.; Matasudaira, P. *J. Org. Chem.* **1995**, *60*, 4305-4308.
9. Relevant analytical data for **1a**, **1b**, **5**, **6a**, **6b**, **7a**, **7b**, **8a** and **8b**. **1a**: ³¹P NMR data (CDCl₃): δ = 15.0 and 15.7 ppm; ¹³C NMR data (CDCl₃): δ = 12.0 (CH₃, thymine), 46.7 (NCH₂), 48.4 (PCH₂), 55.0 (OCH₃, DMT), 56.3 (OCH₃, MOPM), 60.7 (CH₂, MOPM); EI (*m/z*): 760 [M+H]⁺. **1b**: ³¹P NMR data (CDCl₃): δ = 15.0 and 15.4 ppm; ¹³C NMR data (CDCl₃): δ = 46.6 (NCH₂), 48.4 (PCH₂), 55.1 (OCH₃, DMT), 56.2 (OCH₃, MOPM), 60.9 (CH₂, MOPM), 96.6 (C-5); EI (*m/z*): 848 [M+H]⁺. **5**: ³¹P NMR data (CDCl₃): δ = 14.8 ppm. **6a**: ³¹P NMR data (CDCl₃): δ = 14.6 ppm; ¹³C NMR data (CDCl₃): δ = 11.5 (CH₃, thymine), 46.7 (NCH₂), 47.6 (PCH₂), 54.7 (OCH₃, DMT), 119.8 (CH, Ph). **6b**: ³¹P NMR data (CDCl₃): δ = 14.6 ppm; ¹³C NMR data (CDCl₃): δ = 46.6 (NCH₂), 47.4 (PCH₂), 54.5 (OCH₃, DMT), 96.6 (C-5), 120.0 (CH, Ph). **7a**: ³¹P NMR data (CDCl₃): δ = 11.0 and 11.1 ppm; ¹³C NMR data (CDCl₃): δ = 11.9 (CH₃, thymine), 46.5 (NCH₂), 47.7 (PCH₂), 54.5 (OCH₃, DMT), 119.6 (CH, Ph). **7b**: ³¹P NMR data (CDCl₃): δ = 11.4 and 11.6 ppm; ¹³C NMR data (CDCl₃): δ = 46.7 (NCH₂), 47.6 (PCH₂), 55.0 (OCH₃, DMT), 96.8 (C-5), 120.1 (CH, Ph). **8a**: ³¹P NMR data (CDCl₃): δ = 19.2 ppm; ¹³C NMR data (CDCl₃): δ = 12.0 (CH₃, thymine), 46.3 (NCH₂), 47.5 (PCH₂), 54.6 (OCH₃, DMT), 56.0 (OCH₃, MOPM), 119.6 (CH, Ph). **8b**: ³¹P NMR data (CDCl₃): δ = 19.3 ppm; ¹³C NMR data (CDCl₃): δ = 46.5 (NCH₂), 47.5 (PCH₂), 54.6 (OCH₃, DMT), 56.2 (OCH₃, MOPM), 96.4 (C-5), 119.7 (CH, Ph).
10. To a stirred solution of **6a** (or **6b**) (1.0 mmol) in a mixture of CH₃CN/H₂O (95/5, v/v, 10 mL) was added DBU (600 μL, 4.0 mmol). After 15 min at 20°C, the reaction mixture was evaporated to dryness. The residue was coevaporated with CH₃CN, redissolved in CH₂Cl₂ (15 mL) and precipitated from diethyl ether (200 mL). The precipitate was collected by centrifugation and dried *in vacuo* to give **7a** (or **7b**).
11. To a solution of **7a** (or **7b**) (1.0 mmol) and 4-methoxy-1-oxido-pyridine (0.75 g, 6.0 mmol) in dry CH₃CN (7 mL) was added TPS-Cl (0.61 g, 2.0 mmol). After stirring for 5 min, 2-hydroxymethyl-4-methoxy-1-oxido-pyridine (0.31 g, 2.0 mmol) in CH₃CN (3 mL) was added. The mixture was stirred for another 30 min, quenched with aqueous NaHCO₃ (M, 10 mL) and extracted with CH₂Cl₂ (2x 50 mL). The organic layer was dried over MgSO₄ and evaporated to dryness. The phosphonate diester **8a** (or **8b**) was purified by silica gel column chromatography (eluent: CH₂Cl₂/CH₃OH, 10/0 to 9/1, v/v). Subsequently the phenyl group was selectively removed by treatment of **8a** (or **8b**) with 0.4 M DBU in CH₃CN/H₂O (95/5, v/v, 10 mL) for 30 min to give **1a** (or **1b**).
12. The solid-support synthesis was carried out on a Pharmacia Gene Assembler using preloaded 5-*O*-DMT-dT-succinyl-CPG (loading 35 μmole/g, purchased from Millipore) as support.
13. FPLC analysis was carried out on a Pharmacia mono-Q HR 5/5 column (anion exchange). Gradient elution was performed at 20°C by building up a gradient starting with buffer A (0.01 M NaOH, pH = 12) and applying buffer B (0.01 M NaOH, 1.2 M NaCl, pH = 12.0) at a flow rate of 2.0 mL/min.

(Received in UK 12 August 1996; revised 5 September 1996; accepted 6 September 1996)