




## Studies in Marine Macrolide Synthesis: Stereocontrolled Synthesis of the AB-Spiroacetal Subunit of Spongistatin 1 (Altohyrtin A).

Ian Paterson,\* Renata M. Oballa and Roger D. Norcross

University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW, UK.

**Abstract:** The C<sub>1</sub>–C<sub>13</sub> subunit **2**, containing the AB-spiroacetal ring system of spongistatin 1 (**1**), was prepared in 11 steps with 90% ds from 3-benzyloxypropanal. Key steps include (i) the aldol reaction between **4** and **11** using (–)-Ipc<sub>2</sub>BCl and (ii) the spiroacetalisation, **3** → **14**.  
Copyright © 1996 Elsevier Science Ltd

The spongistatins are a novel group of cytotoxic macrolides, isolated from marine sponges of the genus *Spongia*,<sup>1</sup> *Spirastrella*,<sup>2</sup> *Hyrtilios*<sup>3</sup> and *Cinachyra*,<sup>4</sup> which were first reported in 1993 by Pettit *et al.*<sup>1a</sup> The complete relative and absolute stereochemistry of spongistatin 1 (≡ altohyrtin A), as defined by elaborate NOESY and Mosher ester <sup>1</sup>H NMR experiments, has recently been proposed as that shown in structure **1** by Kitagawa and coworkers.<sup>3a</sup> However, this configurational assignment is partially in conflict with the stereochemistry proposed by the Pettit<sup>2b,5</sup> and Fusetani<sup>4</sup> groups. The spongistatins are reported to be among the most potent compounds tested against a distinctive subset of highly chemoresistant tumour types in the US National Cancer Institute's panel of 60 human cancer cell lines (mean GI<sub>50</sub> = 10<sup>-11</sup> M)<sup>5</sup> and are believed to function by inhibiting tubulin polymerisation.<sup>6</sup>




Scheme 1

Due to their extremely meagre natural supply (10<sup>-6</sup>–10<sup>-7</sup>% isolated yield from the whole sponge) and the painstaking isolation procedure, synthetic efforts towards the spongistatins are required to firmly establish the structures and provide further material for biological testing. As part of our studies in marine macrolide synthesis,<sup>7</sup> we now report a stereocontrolled synthesis of the C<sub>1</sub>–C<sub>13</sub> spiroacetal segment **2** of spongistatin 1 (**1**) by using the asymmetric boron aldol reactions<sup>8,9</sup> of methyl ketones.


Central to our synthetic strategy for **2** is the thermodynamically-controlled spiroacetalisation of the open-chain precursor **3** in **Scheme 1**, followed by manipulation of the C<sub>5</sub> and C<sub>9</sub> hydroxyl groups. Note that the AB-spiroacetal subunit of the spongistatins possesses an energetically favourable, double anomeric effect and has the C<sub>3</sub> and C<sub>11</sub> sidechains arranged equatorially on each tetrahydropyran ring. The stereocontrolled construction of the pseudo-C<sub>2</sub>-symmetric chain in **3** was planned around the sequential, two-directional, aldol coupling of acetone with the chiral aldehyde **4**. Previously, we have reported the use of (+)- and (-)-diisopinocampheylboron chlorides (Ipc<sub>2</sub>BCl) for asymmetric aldol reactions of methyl ketones with various aldehydes.<sup>9</sup> As part of a novel approach to 1,3-polyol synthesis,<sup>10</sup> we now extend this chemistry to double<sup>9a</sup> and triple<sup>11</sup> stereodifferentiating aldol reactions with β-oxygenated aldehydes such as **4** (**Scheme 2**).

First, we required a practical asymmetric synthesis of (*S*)-**4** in high enantiomeric purity.<sup>12</sup> Following the Brown protocol,<sup>13</sup> addition (Et<sub>2</sub>O, -78 °C) of the allylborane **5** (obtained from (+)-2-carene *via* **6**) to 3-benzyloxypropanal gave, after oxidative workup, a 93% yield of **7**. The homoallylic alcohol **7** was produced with 97% ee (MTPA ester) on a 4 g scale, which proved to be superior to all other enantioselective allylation methods we examined. Silyl protection as the TES ether, followed by ozonolysis, then gave the required (*S*)-aldehyde **4** in 87% yield.



**Scheme 2:** (a) Allylmagnesium bromide, Et<sub>2</sub>O, -78 °C, 15 min; 20 °C, 1 h; (b) **5**, -78 °C, 4 h; H<sub>2</sub>O<sub>2</sub>, NaOH, Δ, 19 h; (c) TESOTf, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, 2 h; (d) O<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, NaHCO<sub>3</sub>, -78 °C; PPh<sub>3</sub>, 20 °C, 3 h; (e) (-)-Ipc<sub>2</sub>BCl or (*c*-C<sub>6</sub>H<sub>11</sub>)<sub>2</sub>BCl or (+)-Ipc<sub>2</sub>BCl, Et<sub>3</sub>N, Et<sub>2</sub>O, 0 °C, 30 min; **4**, -78 °C, 5 h; H<sub>2</sub>O<sub>2</sub>, MeOH-pH 7 buffer; (f) TBSOTf, 2,6-lutidine, THF, -78 °C, 2 h.

We next explored the use of reagent control in the boron-mediated aldol reaction of acetone to aldehyde **4**. Using our standard conditions,<sup>9a,b</sup> the boron enolate **8a** was prepared by enolisation of acetone with (-)-Ipc<sub>2</sub>BCl (Et<sub>3</sub>N, Et<sub>2</sub>O, 0 °C), followed by addition to **4** at -78 °C. This matched reaction proceeded in 89% yield to give predominantly the 1,3-*syn* isomer **9** with 93% ds by *re*-face attack. In comparison, the dicyclohexylboron<sup>14</sup> enolate **8b** produced **9** with 75% ds, indicating a moderate level of 1,3-*syn* induction from the aldehyde.<sup>9a,15</sup> The enolate **8c** prepared from (+)-Ipc<sub>2</sub>BCl overturned this result, giving a small preference for formation of the corresponding 1,3-*anti* isomer **10**. After silyl protection of **9** as the TBS ether, methyl ketone **11** was obtained in 78% yield from **4**. The synthesis of the open-chain spiroacetal precursor **3** from **4** and **11** exploited *triple asymmetric induction*,<sup>11</sup> where the influence of all three chiral components (aldehyde, ketone and boron reagent) were matched. Using the boron enolate **12a** prepared from **11** with (-)-Ipc<sub>2</sub>BCl, the aldol coupling with **4** proceeded in 81% yield to provide a 97 : 3 mixture of adducts, in favour of the 1,3-*syn* isomer **3**.<sup>16</sup> Here, the 1,3-*syn* preference of aldehyde **4** (*i.e.* *re*-face attack) and the remote 1,5-*anti* preference<sup>17</sup> of methyl ketone **11** act in a synergistic fashion, thereby providing excellent levels of stereocontrol. When using the dicyclohexylboron enolate **12b** alone, the selectivity of this aldol reaction was reduced to 91% ds, while the enolate **12c** formed from (+)-Ipc<sub>2</sub>BCl gave 89% ds. In contrast to these boron aldol results, the corresponding lithium enolate (LDA, THF, -78 °C) led to a reversal in selectivity to give **13** with 66% ds (83% yield).



**Scheme 3:** (a) PPTS, CH<sub>2</sub>Cl<sub>2</sub>/MeOH (1:1), 20 °C, 3 h; (b) TPAP, NMO, MeCN, 4Å powdered mol. sieves, 20 °C, 1 h; (c) MeMgBr, THF, -78 °C → -20 °C, 2 h; (d) CSA, MeOH, 20 °C, 2 h; (e) Ac<sub>2</sub>O/CH<sub>2</sub>Cl<sub>2</sub> (4:1), DMAP, 20 °C, 90 min.

As shown in **Scheme 3**, treatment of **3** with PPTS in MeOH/CH<sub>2</sub>Cl<sub>2</sub> led to clean removal of both TES protecting groups and *in situ* acetalisation to produce a single spiroacetal **14** in 77% yield. Note that although the C<sub>9</sub> stereocentre is later destroyed by oxidation, efficient thermodynamic spiroacetalisation only proceeds with the (9*S*)-configuration.<sup>18</sup> A possible rationale for this observation is the extra stabilisation obtained by the intramolecular hydrogen bond which develops in **14** between the C<sub>9</sub> *axial* hydroxyl and the spiroacetal oxygen. TPAP oxidation<sup>19</sup> of the alcohol **14** and equatorial addition of MeMgBr to the resultant ketone then gave the tertiary alcohol **15** in 85% yield. Finally, conversion of the TBS-protected alcohol at C<sub>5</sub> to the corresponding acetate was accomplished in 2 steps (95%) to provide the desired spiroacetal **2**, [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -67.2° (c 1.17, CHCl<sub>3</sub>). The <sup>1</sup>H and <sup>13</sup>C NMR spectral data obtained<sup>12</sup> for **2** closely resembles that reported for the corresponding segment of the spongistatins. In particular, the NOESY results and coupling constants for **2** are in full accord with that described for spongistatin 1 (althoyrtin A), which supports the structure proposed for the C<sub>1</sub>-C<sub>13</sub> region containing the AB-spiroacetal ring system (*cf.* boxed region of **1**).

In summary, this synthesis of the C<sub>1</sub>–C<sub>13</sub> spiroacetal subunit **2** of spongistatin 1 (altohrytin A) proceeds in 11 steps (32% yield) with 90% overall diastereoselectivity. This work, together with that reported in the accompanying paper,<sup>17</sup> demonstrates that high diastereoselectivity can be obtained for methyl ketone aldol reactions using chiral boron enolates, providing a practical entry into the synthesis of 1,3-polyols. Studies directed towards the total synthesis of spongistatin 1 (**1**) are currently underway.

**Acknowledgement:** We thank the NSERC of Canada (Postdoctoral Fellowship to RMO), the EPSRC and Merck, Sharp & Dohme for support.

## References and Notes

- (a) Pettit, G. R.; Cichacz, Z. A.; Gao, F.; Herald, C. L.; Boyd, M. R.; Schmidt, J. M.; Hooper, J. N. A. *J. Org. Chem.* **1993**, *58*, 1302. (b) Pettit, G. R.; Cichacz, Z. A.; Gao, F.; Herald, C. L.; Boyd, M. R. *J. Chem. Soc., Chem. Commun.* **1993**, 1166.
- (a) Pettit, G. R.; Herald, C. L.; Cichacz, Z. A.; Gao, F.; Schmidt, J. M.; Boyd, M. R.; Christie, N. D.; Boettner, F. E. *J. Chem. Soc., Chem. Commun.* **1993**, 1805. (b) Pettit, G. R.; Cichacz, Z. A.; Herald, C. L.; Gao, F.; Boyd, M. R.; Schmidt, J. M.; Hamel, E.; Bai, R. *J. Chem. Soc., Chem. Commun.* **1994**, 1605.
- Some of the altohrytin macrolides reported in the following papers are assumed to be structurally identical to the spongistatins (*i.e.* altohrytin A corresponds to spongistatin 1): (a) Kobayashi, M.; Aoki, S.; Kitagawa, I. *Tetrahedron Lett.* **1994**, *35*, 1243. (b) Kobayashi, M.; Aoki, S.; Sakai, H.; Kawazoe, K.; Kihara, N.; Sasaki, T.; Kitagawa, I. *Tetrahedron Lett.* **1993**, *34*, 2795. (c) Kobayashi, M.; Aoki, S.; Sakai, H.; Kihara, N.; Sasaki, T.; Kitagawa, I. *Chem. Pharm. Bull.* **1993**, *41*, 989.
- Cinachyrolide A is assumed to be structurally identical to spongistatin 4 (see ref 2a): Fusetani, N.; Shinoda, K.; Matsunaga, S. *J. Am. Chem. Soc.* **1993**, *115*, 3977.
- Pettit, G. R. *Pure Appl. Chem.* **1994**, *66*, 2271 and references therein.
- (a) Bai, R.; Cichacz, Z. A.; Herald, C. L.; Pettit, G. R.; Hamel, E. *Mol. Pharmacol.* **1993**, *44*, 757. (b) Bai, R.; Taylor, G. F.; Cichacz, Z. A.; Herald, C. L.; Kepler, J. A.; Pettit, G. R.; Hamel, E. *Biochemistry* **1995**, *34*, 9714.
- Norcross, R. D.; Paterson, I. *Chem. Rev.* **1995**, *95*, 2041.
- Cowden, C. J.; Paterson, I. *Org. React.* **1997**, in press.
- (a) Paterson, I.; Cumming, J. G.; Smith, J. D.; Ward, R. A. *Tetrahedron Lett.* **1994**, *35*, 441. (b) Paterson, I.; Smith, J. D. *Tetrahedron Lett.* **1993**, *34*, 5351. (c) Paterson, I. *Pure Appl. Chem.* **1992**, *64*, 1821. (d) Paterson, I.; Goodman, J. M.; Lister, M. A.; Schumann, R. C.; McClure, C. K.; Norcross, R. D. *Tetrahedron* **1990**, *46*, 4663. (e) Paterson, I.; Goodman, J. M. *Tetrahedron Lett.* **1989**, *30*, 997.
- Oishi, T.; Nakata, T. *Synthesis* **1990**, 635.
- Duplantier, A. J.; Nantz, M. H.; Roberts, J. C.; Short, R. P.; Somfai, P.; Masamune, S. *Tetrahedron Lett.* **1989**, *30*, 7357.
- All new compounds gave spectroscopic data in agreement with the assigned structures. NMR data for spiroacetal **2**: <sup>1</sup>H NMR δ (500 MHz, DMSO-d<sub>6</sub>) 1.03 (3H, s, C<sub>9</sub>-Me), 1.21 (1H, t, *J* = 12.3 Hz, H<sub>10A</sub>), 1.38 (1H, d, *J* = 13.9 Hz, H<sub>8A</sub>), 1.51–1.46 (2H, m, H<sub>4A</sub> and H<sub>10B</sub>), 1.71–1.55 (7H, m, H<sub>2A</sub>, H<sub>2B</sub>, H<sub>4B</sub>, H<sub>6A</sub>, H<sub>8B</sub>, H<sub>12A</sub> and H<sub>12B</sub>), 1.77 (1H, d, *J* = 15 Hz, H<sub>6B</sub>), 1.89 (3H, s, -OC(O)CH<sub>3</sub>), 3.44–3.35 (2H, m, -CH<sub>2</sub>OBn), 3.56 (2H, dt, *J* = 5.7, 1.8 Hz, -CH<sub>2</sub>OBn), 3.89 (1H, br t, *J* = 12.3 Hz, H<sub>11</sub>), 3.99–3.95 (1H, m, H<sub>3</sub>), 4.21 (1H, s, 9-OH), 4.36–4.27 (4H, m, 2 x -OCH<sub>2</sub>Ph), 4.89 (1H, br s, H<sub>5</sub>), 7.32–7.22 (10H, m, aromatic); <sup>13</sup>C NMR δ (100 MHz, DMSO-d<sub>6</sub>) 21.2, 30.4, 34.2, 35.0, 35.4, 36.7, 43.6, 45.4, 62.2, 62.6, 66.5, 66.7, 67.0, 67.4, 72.1, 72.2, 97.3, 127.4, 127.5, 127.7, 128.3, 138.3, 138.6, 169.9.
- Brown, H. C.; Randad, R. S.; Bhat, K. S.; Zaidlewicz, M.; Racherla, U. S. *J. Am. Chem. Soc.* **1990**, *112*, 2389.
- Brown, H. C.; Dhar, R. K.; Ganesan, K.; Singaram, B. *J. Org. Chem.* **1992**, *57*, 499.
- The 1,3-induction obtained with such boron-mediated aldol reactions of methyl ketones is influenced by several factors, one of which is the nature of the β-hydroxyl protecting group in the aldehyde. See also: (a) Evans, D. A.; Duffy, J. L.; Dart, M. J. *Tetrahedron Lett.* **1994**, *35*, 8537. (b) Gustin, D. J.; VanNieuwenhze, M. S.; Roush, W. R. *Tetrahedron Lett.* **1995**, *36*, 3443. (c) Paterson, I.; Bower, S.; Tillyer, R. D. *Tetrahedron Lett.* **1993**, *34*, 4393.
- The configuration of the newly introduced stereocentre at C<sub>9</sub> was verified by TBS protection of the hydroxyl group of **3**, thus generating a C<sub>2</sub>-symmetric compound.
- Paterson, I.; Gibson, K. R.; Oballa, R. M. *Tetrahedron Lett.* **1996**, *37*, 0000 (following paper).
- Alternatively, when **3** was oxidised to the acyclic β-diketone, all attempts to effect efficient spiroacetalisation failed.
- Ley, S. V.; Norman, J.; Griffith, W. P.; Marsden, S. P. *Synthesis* **1994**, 639.

(Received in UK 30 August 1996; accepted 4 October 1996)