



MICROWAVE-ASSISTED OXIDATION OF ALCOHOLS UNDER SOLVENT-FREE CONDITIONS USING CLAYFEN

Rajender S. Varma* and Rajender Dahiya

Department of Chemistry and Texas Regional Institute for Environmental Studies (TRIES),
Sam Houston State University, Huntsville, TX 77431-2117, USA

Key Words: Alcohols; Carbonyl compounds; Clayfen; Iron(III) nitrate nonahydrate; Montmorillonite K 10 clay; Microwave irradiation.

Abstract: Adsorbed on clayfen, alcohols are readily oxidized to carbonyl compounds upon exposure to microwaves under solvent-free conditions. This rapid, selective and environmentally benign method conserves the use of excess solvents and toxic oxidants usually employed. © 1997 Elsevier Science Ltd.

The oxidation of alcohols to carbonyl compounds is an important transformation in organic synthesis and several methods have been explored to accomplish this conversion.¹ In this process, the use of heavy metals and peracids is very common, although selective methods are sought that are not detrimental to the environment.^{2,3} In this context, the use of inexpensive supported reagents^{4,5} has attracted attention because of the selectivity and the ease of manipulation. Specifically, clay-supported reagents have found widespread acceptance in a variety of heterogeneous reactions.^{6,7}

The microwave (MW) enhanced chemical reactions,^{8,9} in general and on inorganic solid supports⁹⁻¹¹ in particular, have gained popularity over the usual homogeneous and heterogeneous reactions^{6,7} as they can be conducted rapidly and provide pure products in quantitative yields without the use of solvents. Since only the polar reactants adsorbed on the surface of the solid support absorb microwaves, a variety of reagents supported on such surfaces^{6,7} can be utilized for the enhancement of organic reactions using an unmodified MW oven.⁸ We report here a facile oxidation of alcohols to carbonyl compounds using montmorillonite K10 clay-supported iron(III) nitrate, clayfen,^{6,7} under solvent-free conditions in a process that is accelerated by microwave irradiation. The reaction presumably proceeds *via* the intermediacy of nitrosonium ions,^{6b} and no formation of carboxylic acids occurs. The experimental procedure involves a simple mixing of neat alcohols with clayfen and irradiating the reaction mixtures in a microwave oven for 15-60 seconds in the absence of any solvent. This extremely rapid, manipulatively simple, inexpensive and selective protocol avoids the use of excess solvents and toxic oxidants. The amount of clayfen used in our reactions is half of that used by earlier workers⁷ under heterogeneous conditions that employs large excess of hydrocarbon solvents. Our results are summarized in the **Table**.

The oxidation of benzoin is representative of the general procedure employed. Clayfen (0.125 g) was thoroughly mixed with neat benzoin (0.106 g, 0.5 mmol) in the solid state using a vortex mixer and the

material was placed in an alumina bath inside the MW oven¹² and irradiated. Upon completion of the reaction, monitored on TLC (hexane:AcOEt, 10:1, v/v), the product was extracted into methylene chloride. That the effect is not purely *thermal*¹² is supported by the fact that this reaction could be completed in 18 hours in an oil bath at a comparable temperature of 65 °C. The notable exception to these conditions was the case of *p*-methoxybenzyl alcohol which was completely oxidized to an aldehyde at room temperature. The limitation with aliphatic alcohols, however, still persists and the protocol gives a mixture of compounds.




Table. Oxidation of alcohols by clayfen under microwave irradiation

Entry	R ₁	R ₂	Time (sec)	Yields (%)	I.R. ν _{C=O} (cm ⁻¹)
1.	C ₆ H ₅	H	15	92	1705
2.	C ₆ H ₅	CH ₂ CH ₃	30	87	1670
3.	C ₆ H ₅	C ₆ H ₅ CO	60	93	1660
4.	<i>p</i> -CH ₃ C ₆ H ₄	H	15	94	1690
5.	<i>p</i> -CH ₃ OC ₆ H ₄	H	–	96	1685
6.	<i>p</i> -CH ₃ OC ₆ H ₄	<i>p</i> -CH ₃ OC ₆ H ₄ CO	60	94	1655
7.	Tetrahydrofurfuryl	H	30	90	1730
8.	–(CH ₂) ₅ –		30	89	1715

In conclusion, clayfen-mediated solvent-free microwave thermolysis is a convenient, selective and environmentally benign oxidative protocol when compared to the conventional solution phase or heterogeneous reactions.

ACKNOWLEDGMENT

We are grateful to the Texas Advanced Research Program (ARP) in chemistry (Grant # 0036060-023) and TRIES under contract from Office of Naval Research/SERDP (Grant # N00014-96-1-1067) for financial support.

REFERENCES AND NOTES

1. *Comprehensive Organic Synthesis (Oxidation)*, Trost B.M., Ed., Pergamon; New York, 1991, Vol 7.
2. Einhorn, J.; Einhorn, C.; Ratajczak, F.; Pierre, J.-L. *J. Org. Chem.* **1996**, *61*, 7452.
3. Muzart, J.; N'Ait Ajjou, A.; Ait-Mohand, S. *Tetrahedron Lett.* **1994**, *35*, 1989.
4. McKillop, A.; Young, D.W. *Synthesis* **1979**, 401 and 481.
5. Posner, G.H. *Angew. Chem.* **1978**, *90*, 527; *Angew. Chem. Int. Ed. Engl.* **1978**, *17*, 487.
6. a) Delaude, L.; Laszlo, P. *J. Org. Chem.* **1996**, *61*, 6360; b) Cornelis, A.; Laszlo, P. *Synthesis* **1985**, 909; c) Balogh, M.; Laszlo, P. *Organic Chemistry Using Clays*, Springer-Verlag, Berlin, **1993**.
7. Cornelis, A.; Laszlo, P. *Synthesis* **1980**, 849.
8. For recent reviews on microwave-assisted chemical reactions, see a) Abramovich, R.A. *Org. Prep. Proced. Int.* **1991**, *23*, 683; b) Whittaker, A.G.; Mingos, D.M.P. *J. Microwave Power Electromagn. Energy* **1994**, *29*, 195; c) Majetich, G.; Hicks, R. *J. Microwave Power Electromagn. Energy* **1995**, *30*, 27; d) Caddick, S. *Tetrahedron* **1995**, *51*, 10403; e) Bose, A.K.; Jayaraman, M.; Okawa, A.; Bari, S.S.; Robb, E.W.; Manhas, M.S. *Tetrahedron Lett.* **1996**, *37*, 6989 and references from this group cited therein.
9. a) Varma, R.S.; Chatterjee, A.K.; Varma, M. *Tetrahedron Lett.* **1993**, *34*, 3207. b) Varma, R.S.; Chatterjee, A.K.; Varma, M. *Tetrahedron Lett.* **1993**, *34*, 4603. c) Varma, R.S.; Varma, M.; Chatterjee, A.K. *J. Chem. Soc., Perkin. Trans. I* **1993**, 999. d) Varma, R.S.; Lamture, J.B.; Varma, M. *Tetrahedron Lett.* **1993**, *34*, 3029.
10. Varma, R.S.; Varma, M. *Tetrahedron Lett.* **1992**, *33*, 5937.
11. Marrero-Terrero, A.L.; Loupy, A. *Synlett* **1996**, 245.
12. For a critical evaluation of activation process by microwaves see: Raner, K.D.; Strauss, C.R.; Vyskoc, F.; Mokbel, L. *J. Org. Chem.* **1993**, *58*, 950. The temperature of the alumina bath (heat sink) inside a Sears Kenmore microwave oven equipped with a turntable at full power (900 Watts) found to be ~65 °C after 30 seconds of irradiation.

(Received in USA 16 December 1996; revised 28 January 1997; accepted 3 February 1997)