



## Selective Dealkylations of Alkyl Aryl Ethers in Heterogeneous Basic Media under Microwave Irradiation.


Abdelouahad Oussaïd, Le Ngoc Thach <sup>1</sup> and André Loupy \*

Laboratoire des Réactions Sélectives sur Supports associé au CNRS  
ICMO Université Paris XI - Bâtiment 410 - 91405 ORSAY France

**Abstract:** Ethyl isoeugenol and 2-ethoxyanisole could be selectively deethylated using potassium t-butoxide in the presence of crown ether under solvent-free conditions. By addition of ethylene glycol, selectivity was reversed to demethylation. In both cases, strong enhancements were observed under focused microwaves. © 1997 Published by Elsevier Science Ltd.

Dealkylation of alkyl aryl ethers is a synthetically very useful reaction as it constitutes an important method for deprotection of hydroxyl groups from substituted phenols <sup>2</sup>. A wide variety of systems has been proposed in homogeneous phase using more generally Lewis acids such as AlCl<sub>3</sub> <sup>3</sup> and BX<sub>3</sub> <sup>4</sup> or strong bases such as alkaline hydroxides <sup>5</sup> or amides <sup>6</sup> and lithium diphenyl phosphide <sup>7</sup>. These conditions suffer from harsh reaction conditions and present a lot of inconvenience connected to their handling, removal and toxicity of catalyts.

Heterogeneous conditions were thus developed with significant advantages. Potassium fluoride coated on alumina, previously widely described as a strong base <sup>8</sup>, was shown by Singh et al. to be also an effective reagent for the selective O-demethylation of aromatic methyl ethers <sup>9</sup>.



In connection with our interest for mild heterogeneous solvent-free conditions <sup>10</sup>, we checked the behaviour of ethyl isoeugenol **1a** using different basic systems. We especially examined phase transfer catalysis (PTC) conditions in the absence of solvent under microwave irradiation as it was shown of great efficiency in a lot of cases including Krapcho decarboxylation <sup>11</sup>. A system potassium t-butoxide/transfer agent was selected as it behaves at the same time as a strong base (H-nucleophilicity) <sup>12</sup> or a strong nucleophile (C-nucleophilicity) <sup>13</sup>.

Ethyl isoeugenol **1a** is able to lead to either propenylguaethol **2a** by O-demethylation or isoeugenol **3a** by O-deethylation. The main results are given in Table I.




Table I: Dealkylation of ethyl isoeugenol **1a** (5 mmoles) using a base (2 eq, 10 mmoles) in the presence of a phase transfer agent (10%) under monomode microwave activation <sup>14</sup>.

Base	Phase transfer agent	Exp. cdt			Yields <sup>a</sup>		
		P W	t <sub>mn</sub>	T °C <sup>b</sup>	% <b>1a</b>	% <b>2a</b>	% <b>3a</b>
KF	Aliquat 336	270	120	177	95	2	-
LiF	"	90	60	116	99	-	-
"	12-crown-4	90	60	123	97	2	-
KOtBu	-	60	20	93	92	-	-
"	Aliquat 336	90	20	175	20	-	60
"	Kryptofix [2.2.2]	90	10	135	15	-	68
"	18-crown-6	60	20	139	9	-	82

a) g. c. yields using a standard; b) maximum in temperature during microwave irradiation measured by infrared detection <sup>15</sup> using a monomode Synthwave 402 reactor (Prolabo)

It was noteworthy that we obtained selective deethylation under these conditions, whereas KF-alumina in ethylene glycol leads to selective demethylation. The two methods seem to be complementary. The effect of this solvent was thus checked under PTC conditions. The more significant results are indicated in Table II.

Table II: Effect of ethylene glycol (EG) on the dealkylation of **1a** (5 mmoles) induced by base (10 mmoles) and 18-crown-6 (10%) under microwave (MW) irradiation or under conventional heating ( $\Delta$ ) using a thermoregulated oil or sand bath.

Base	E G (Volume / ml)	mode	Exp. cdt			Yields		
			P W	t <sub>mn</sub>	T °C	% <b>1a</b>	% <b>2a</b>	% <b>3a</b>
KOtBu	-	MW	60	20	139	9	-	<b>82</b>
"	-	$\Delta$		20	139	32	-	52
"	2	MW	60	60	228	6	<b>56</b>	3
"	2	$\Delta$		60	228	70	2	-
NaOH	5	MW	60	180	200	5	<b>70</b>	2
"	5	$\Delta$		180	200	70	-	-

The addition of ethylene glycol was determinant from selectivity point of view in the case of KOtBu and crown ether reaction. This is at the origin of a total reversal in selectivity, allowing O-demethylation.

To avoid the possible influence of the double bond on aromatic ring and check the generality of our observation, same conditions were extrapolated to 2-ethoxyanisole **1b** (Table III).

Table III: Dealkylation of 2-ethoxyanisole **1b** (5 mmoles) using base (10 mmoles, 2 eq.) in the presence of a phase transfer agent (10%) with or without ethylene glycol (E G) under microwave irradiation <sup>14</sup> (MW) or using a thermoregulated oil bath ( $\Delta$ ).

Base	Phase transfer agent	E G (ml)	mode	Exp. cdt's			Yields		
				P <sub>w</sub> <sup>a</sup>	t <sub>mn</sub>	T °C	% <b>1b</b>	% <b>2b</b>	% <b>3b</b>
KOtBu (2 eq.)	-	-	MW	60	20	120	39	-	60
"	Aliquat 336	-	MW	60	20	120	28	-	66
"	18-crown-6	-	MW	60	20	120	7	-	<b>90</b>
"	P E G 400	-	MW	60	20	120	13	-	80
"	T D A - 1	-	MW	60	20	120	7	-	<b>91(90)</b>
"	"	-	$\Delta$		20	120	48	-	50
"	"	-	$\Delta$		120	120	28	-	60
KOtBu (2 eq.)	Aliquat 336	2	MW	60	60	180	86	10	3
"	T D A - 1	2	MW	60	60	180	93	5	1
"	18-crown-6	2	MW	60	60	180	20	58	22
"	"	2	MW	60	75	180	-	<b>72</b>	23
"	"	2	$\Delta$		75	180	<b>98</b>	-	-
"	"	2	$\Delta$		1200	180	63	26	10
NaOH (2 eq.)	15-crown-5	-	MW	60	30	180	<b>93</b>	-	2
"	"	-	$\Delta$		75	180	<b>99</b>	-	-
NaOH (4 eq.)	15-crown-5	5	MW	20	120	205	<b>5</b>	<b>77</b>	10
"	"	5	$\Delta$		120	205	<b>94</b>	-	-

a) maximal in power during microwave irradiation.

The conclusions were rather the same for those two substrates:

- deethylation is the dominant reaction with KOtBu and crown ether (or TDA-1) in the absence of any organic solvent. A strong microwave effect on reactivity was observed as the yield fell to only 50% under conventional heating, instead of 90% under microwave irradiation, without any change on selectivity.
- demethylation became the dominant reaction when ethylene glycol was added. A very strong microwave influence (not purely thermal) was shown as no reaction occurred under conventional heating (same conditions of time and temperature). This observation was in accordance with some results from literature <sup>16</sup>.

In terms of mechanism, as KOtBu is known as a very strong base, we can envisage a  $\beta$ -elimination for deethylation whereas KOtBu and EG (leading probably to  $K^+$ ,  $-OCH_2CH_2OH$  or at least a decrease in basicity of KOtBu due to his solvation by EG) leads to a nucleophilic attack on the less hindered methyl group.



We are grateful to Prolabo, France for useful discussions and financial support. We are also indebted to Patrice Moreau for helpful contacts.

#### REFERENCES AND NOTES:

1. On leave from Department of Organic Chemistry, University of Ho-Chi-Minh-City (Viet-Nam).
2. a) Sala T.; Sargent M. V. *J. Chem. Soc. Perkin Trans I*, **1979**, 2593-2598.  
b) Bhatt M. V.; Kulkarni S. U. *Synthesis*, **1983**, 249-282.
3. Burwell R. L.; Jr. *Chem. Rev.* **1954**, *54*, 615-685.
4. Gerrard W.; Lappert M. F. *Chem. Rev.* **1958**, *58*, 1081-1111.
5. Sainsbury M.; Dyle S. F.; Moon B. J. *J. Chem. Soc. (C)*, **1970**, 1797-1800.
6. Brotherton T. K.; Bunnett J. F. *Chem. & Ind. (London)*, **1957**, 80.
7. Veriot G.; Collet A. *Acros Organics Acta*, **1995**, *1*, 40-41.
8. Clark J. H. *Chem. Rev.*, **1980**, *80*, 429-452.
9. Radhakrishna A. S.; Prasad Rao K. R. K.; Suri S. K.; Sivaprakash K.; Singh B. B. *Synth. Commun.* **1991**, *21*, 379-383.
10. Loupy A.; Bram G.; Sansoulet J. *New J. Chem.* **1992**, *16*, 233-242.
11. a) Loupy A.; Pigeon P.; Ramdani M.; Jacquault P. *J. Chem. Res. (S)*, **1993**, 36-37.  
b) Barnier J. P.; Loupy A.; Pigeon P.; Ramdani M.; Jacquault P. *J. Chem. Soc. Perkin Trans I*, **1993**, 397-398.
12. Le Ngoc T.; Duong-Lieu H.; Nguyen Ba H.; Radhakrishna A. S.; Singh B. B.; Loupy A. *Synth. Commun.* **1993**, *23*, 1379-1384.
13. Barry J.; Bram G.; Decodts G.; Loupy A.; Pigeon P.; Sansoulet J. *Tetrahedron*, **1984**, *40*, 2945-2950.
14. Loupy A. *Spectra Analyse*, **1993**, *175*, 33-38, and references cited therein.  
Reactions were performed under continuous stirring using a monomode Synthewave 402 reactor from Prolabo (France) fitted with an infrared temperature detector <sup>15</sup> calibrated and controlled with a luxtron optical fiber inside the medium.
15. Jacquault P. (Prolabo, France), European patent n° 92-420477.9 (21.12.92) **1992**.
16. a) Bougrin K.; Kella Bennani A.; Fkih Tetouani S.; Soufiaoui M. *Tetrahedron Lett.* **1994**, *35*, 8373-8376.  
b) Bougrin K.; Soufiaoui M.; Loupy A.; Jacquault P. *New J. Chem.* **1995**, *19*, 213-219.  
c) Perez E. R.; Marrero A. L.; Perez E.; Autie M.A. *Tetrahedron Lett.* **1995**, *36*, 1779-1783.  
d) Loupy A.; Pigeon P.; Ramdani M. *Tetrahedron*, **1996**, *52*, 6705-6712.

(Received in France 23 January 1997; accepted 21 February 1997)