

General Syntheses of Ethyl 2-Ulosonates and 3-Deoxy-D-manno-oct-2-ulosonic Acid from Carbohydrate Lactones with 1-Ethoxyvinyl lithium


Shende Jiang,^a Anthony D. Rycroft,^a Gurdial Singh,^{a*} Xiang-Zhu Wang^{a,b} and Yu-Lin Wu^b

^aDepartment of Chemistry, University of Sunderland, Sunderland, SR1 3SD, U.K. ^bState Key Laboratory of Bio-organic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, China

Received 2 February 1998; revised 11 March 1998; accepted 13 March 1998

Abstract: Reactions of 1-ethoxyvinyl lithium with isopropylidene or benzyl protected sugar lactones afforded the corresponding hemiacetals which, on ozonolysis, gave the ethyl 2-ulosonates. This allowed a rapid and efficient synthesis of 3-deoxy-D-manno-oct-2-ulosonic acid (KDO). © 1998 Elsevier Science Ltd. All rights reserved.


The 3-deoxy-D-manno-oct-2-ulosonic acid (KDO) **1** is an essential constituent of lipopolysaccharides (LPS) and acidic exopolysaccharides (K antigens) found in the cell surface of Gram-negative bacteria.¹ The incorporation of KDO is essential if cell growth is to occur in the bacteria. Thus the design and synthesis of potential inhibitors of lipopolysaccharide biosynthesis based upon analogues of KDO has attracted much attention, as has the synthesis of KDO itself.² Another ulosonic acid, 3-deoxy-D-arabino-hepta-2-ulosonic acid (DAH) **2**, plays an important role in the biosynthesis of amino acids in plants and micro-organisms as a key intermediate in the shikimate pathway.³ It is therefore not surprising that these structures have invoked immense interests from both biological and synthetic standpoints.




As part of an ongoing programme to study the shikimate pathway, in particular, the syntheses of key biosynthetic intermediates and their analogues from carbohydrates,⁴ we wished to develop a general strategy for the synthesis of a range of 2-ulosonic acids.

Since the first report on the alpha lithiation of ethyl vinyl ether,⁵ the 1-alkoxyvinyl lithium reagents have been employed in 1,2 and 1,4-addition reactions,⁶ as well as for natural product syntheses.⁷ As a result of this we were attracted to the use of 1-ethoxyvinyl lithium,⁸ because it is an ideal acyl anion equivalent that is easy to prepare and the precursor, ethyl vinyl ether, is available in commercial quantities. Combination of the reactivity of 1-ethoxyvinyl lithium with readily available sugar lactones thus provides a concise and highly attractive approach to the synthesis of 2-ulosonic acids. In this letter we report on our findings employing this strategy.


Our initial studies centred on the addition of 1-ethoxyvinyl lithium to 2,3-*O*-isopropylidene-D-erythro-1,4-lactone **3**⁹ and the subsequent ozonolysis. The reaction of lactone **3** with 1-ethoxyvinyl lithium proceeded smoothly in THF at $-78\text{ }^{\circ}\text{C}$ to furnish the hemiacetal **4** as an anomeric mixture in 83% yield.¹⁰ Ozonolysis of lactol **4** in mixed solvents of dichloromethane and ethanol at $-78\text{ }^{\circ}\text{C}$ afforded the ethyl D-erythro-pent-2-ulosonate **5** in 88% yield (scheme 1).



Scheme 1 Reagents and conditions: i, 1-ethoxyvinyl lithium, THF, $-78\text{ }^{\circ}\text{C}$ (83%); ii, O_3 , CH_2Cl_2 , EtOH, $-78\text{ }^{\circ}\text{C}$; then Me_2S (88%)



Following this success, we investigated the reactions of 1-ethoxyvinyl lithium with isopropylidene protected furanone lactones **6**,¹¹ **7**¹² and **8**.¹³ Thus 1-ethoxyvinyl lithium readily added to the lactones **6**, **7**, and **8** and the resulting hemiacetals were converted to the corresponding ethyl ulosonates. The hemiacetals derived from these lactones proved somewhat labile towards chromatography on silica gel. However, in all cases they were successfully isolated when the silica gel was deactivated with triethylamine and the chromatography was conducted rapidly. Subsequent cleavage of the double bond in the hemiacetals with ozone furnished the corresponding ethyl ulosonates as anomeric mixtures (Table 1).




Scheme 2 Reagents and conditions: i, 1-ethoxyvinyl lithium, THF, $-78\text{ }^{\circ}\text{C}$ (67%); ii, O_3 , CH_2Cl_2 , EtOH, $-78\text{ }^{\circ}\text{C}$; then Me_2S (74%)

Attention was then turned to the six-membered pyranone lactones. Treatment of 2,3,4-tri-*O*-benzyl-L-arabinono-1,5-lactone **9**¹⁴ with 1-ethoxyvinyl lithium in THF at $-78\text{ }^{\circ}\text{C}$ afforded the hemiacetal **10** in 67% yield as a mixture of anomers. Ozonolysis of **10** proceeded uneventfully and gave the ethyl ulosonate **11** in 74% yield with the β -anomer as the major isomer.

Table 1

Lactone	Yield of hemiacetal (%)	Yield of ethyl 2-ulosonate (%)
3	D-erythro hemiacetal 4 (83)	D-erythro 2-ulosonate 5 (88)
6	D-ribo hemiacetal (84)	D-ribo 2-ulosonate (80)
7	D-lyxo hemiacetal (85)	D-lyxo 2-ulosonate (85)
8	D-manno hemiacetal (86)	D-manno 2-ulosonate (84)
9	L-arabino hemiacetal (67)	L-arabino 2-ulosonate (74)

Application of this strategy to the pyranonolactone **12**^{2h} gave the corresponding hemiacetal **13** in 60% yield. Cleavage of the double bond in compound **13** with ozone afforded the ester **14**^{2d} (84% yield), $[\alpha]_D +49.4$ (c 0.57, CHCl₃). Deprotection of ethyl ulosonate **14** with aqueous trifluoroacetic acid, followed by treatment with aqueous ammonia gave the ammonium KDO **15**, the spectral and physical properties of which were in excellent agreement with those reported in the literature.^{2g,2j,15}



Scheme 3 Reagents and conditions: i, 1-ethoxyvinyl lithium, THF, $-78\text{ }^{\circ}\text{C}$ (60%); ii, O₃, CH₂Cl₂, EtOH, $-78\text{ }^{\circ}\text{C}$; then Me₂S (84%); iii, aq. TFA (10%), 80 $^{\circ}\text{C}$, 30 min; iv, aq. NH₃

In summary, we have demonstrated that 1-ethoxyvinyl lithium adds to a wide range of sugar derived lactones in an efficient manner and this strategy provides ready access to ulosonic acids in a short number of steps.

Acknowledgements: We thank Dr. Richard H. Wightman (Heriot-Watt University) for helpful discussions and Dr. M. Batley (Macquarie University) for the provision of some spectral data on KDO derivatives. We also thank the EPSRC for a ROPA grant and for access to mass spectrometry service at the University of Wales, Swansea (Director, Dr. J. A. Ballantine).

References and Notes

- Unger, F. M. *Adv. Carbohydr. Chem. Biochem.* **1981**, *38*, 323.
- (a) Danishefsky, S. J.; Pearson, W. H.; Segmuller, B. E. *J. Am. Chem. Soc.* **1985**, *107*, 1280; (b) Ramage, R.; Rose, G. W. *Tetrahedron Lett.* **1988**, *29*, 4877; (c) Branchaud, B. P.; Meier, M. S. *J. Org. Chem.* **1989**, *54*, 1320; (d) Boons, G. J. P. H.; van der Klein, P. A. M.; van der Marel, G. A.; van Boom, J. H. *Recl. Trav. Chim. Pays-Bas* **1990**, *109*, 273; (e) Dondoni, A.; Merino, P. *J. Org. Chem.* **1991**, *56*, 5294; (f) Shing, T. K. M. *Tetrahedron Lett.* **1992**, *33*, 1307; (g) Giese, B.; Linker, T. *Synthesis* **1992**, 47; (h) Haudrechy, A.; Sinaÿ, P. *J. Org. Chem.* **1992**, *57*, 4142; (i) Shing, T. K. M. *Tetrahedron: Asymmetry* **1994**, *5*, 2405; (j) Gao, J.; Härter, R.; Gordon, D. M.; Whitesides, G. M. *J. Org. Chem.* **1994**, *59*, 3714; (k) Craig, D.; Pennington, M. W.; Warner, P. *Tetrahedron Lett.* **1995**, *36*, 5815; (l) Barton, D. H. R.; Jaszberenyi, J. Cs.; Liu, W.; Shinada, T. *Tetrahedron* **1996**, *52*, 2717; (m) Coutrot, P.; Grison, C.;

- Lecouvey, M. *Tetrahedron Lett.* **1996**, *37*, 1595; (n) Du, S.; Plat, D.; Baasov, T. *Tetrahedron Lett.* **1996**, *37*, 3545; (o) López-Herrera, F. J.; Sarabia-García, F. *Tetrahedron* **1997**, *53*, 3325; (p) Pakulski, Z.; Zamojski, A. *Tetrahedron* **1997**, *53*, 3723; (q) Tsukamoto, H.; Takahashi, T. *Tetrahedron Lett.* **1997**, *38*, 6415.
3. Haslam, E. *Shikimic Acid, Metabolism and Metabolites*, Wiley, Chichester, 1993.
 4. (a) Jiang, S.; Mekki, B.; Singh, G.; Wightman, R. H. *Tetrahedron Lett.* **1994**, *35*, 5505; (b) Jiang, S.; McCullough, K. J.; Mekki, B.; Singh, G.; Wightman, R. H. *J. Chem. Soc., Perkin Trans. 1* **1997**, 1805.
 5. Schöllkopf, U.; Hänssle, P. *Liebigs Ann. Chem.* **1972**, *763*, 208.
 6. (a) Paquette, L. A.; Andrew, J. F. P.; Vanucci, C.; Lawhorn, D. E.; Negri, J. T.; Rogers, R. D. *J. Org. Chem.* **1992**, *57*, 3956; (b) Lesimple, P.; Beau, J.-M.; Jaurand, G.; Sinaÿ, P. *Tetrahedron Lett.* **1986**, *27*, 6201; (c) Baldwin, S. W.; Mazzuckelli, T. J. *Tetrahedron Lett.* **1986**, *27*, 5975; (d) Boeckman, R. K., Jr.; Bruza, K. J. *Tetrahedron* **1981**, *37*, 3997; (e) Soderquist, J. A.; Hassener, A. *J. Am. Chem. Soc.* **1980**, *102*, 1577; (f) Corey, E. J.; Boger, D. L. *Tetrahedron Lett.* **1978**, *13*; (g) Boeckman, R. K., Jr.; Bruza, K. J.; Baldwin, J. E.; Lever, O. W., Jr.; *J. Chem. Soc., Chem. Commun.* **1975**, 519.
 7. (a) Boger, D. L.; Zhu, Y. *J. Org. Chem.* **1994**, *59*, 3453; (b) Maleczka, R. E., Jr.; Paquette, L. A. *J. Org. Chem.* **1991**, *56*, 6538; (c) Maleczka, R. E., Jr.; Paquette, L. A. *J. Org. Chem.* **1991**, *56*, 912.
 8. Meyers, A. I.; Shimano, M. *J. Am. Chem. Soc.* **1994**, *116*, 10815.
 9. Cohen, N.; Banner, B. L.; Lopresti, R. J.; Wong, F.; Rosenberger, M.; Liu, Y. Y.; Thom, E.; Liebman, A. A. *J. Am. Chem. Soc.* **1983**, *105*, 3661.
 10. All new compounds gave satisfactory spectral and microanalytical data. Selected data: **5**: Anomeric mixture (4:1); $[\alpha]_D -80.5$ (c 3.18, CHCl₃); ν_{\max} (neat film) 3432, 1743 cm⁻¹; δ_H (270 MHz) 1.34 (3H, t, *J* 7.25 Hz, major), 1.35 (3H, appt, *J* 7.25 Hz, minor), 1.30(3H, s, major), 1.40 (3H, s, minor), 1.45 (3H, s, major), 1.58 (3H, s, minor), 4.11 (1H, appdd, minor), 4.14 (1H, dd, *J* 10.56, 3.3 Hz, major), 4.30 (2H, q, *J* 7.26 Hz, major), 4.32 (2H, q, *J* 7.25 Hz, minor), 4.39 (1H, d, 1.98 Hz, minor, OH), 4.43 (1H, bs, OH), 4.68 (1H, d, *J* 5.94 Hz, major), 4.79 (1H, d, *J* 5.93 Hz, minor), 4.90 (1H, ddd, *J* 5.92, 3.93, 3.3 Hz, major, minor); δ_C (67.8 MHz) *major anomer* 13.94, 24.82, 25.81, 62.04, 72.17, 79.81, 86.69, 103.95, 113.15, 168.31; *minor anomer* 13.86, 24.74, 25.92, 61.96, 70.03, 80.01, 80.18, 103.95, 113.96, 168.45; m/z HRMS (CI, NH₃), Found MNH₄⁺, 250.1291, C₁₀H₂₀NO₆ requires MNH₄, 250.1291. **13**: mixture of anomers; ν_{\max} (neat film) 3498, 1612 cm⁻¹; δ_H (270 MHz) 1.14-1.57 (15H, m), 2.03-2.26 (2H, m), 3.09-3.39 (1.7H, m), 3.40-3.53 (0.3H, m), 3.60-3.80 (2H, m), 3.81-4.07 (3H, m), 4.08-4.22 (0.4H, m), 4.23-4.54 (2H, m), 4.56-4.77 (1H, m), 5.03-5.20 (0.6H, m); δ_C (67.8 MHz) *major anomer* 157.14, 109.07, 109.01, 95.60, 76.33, 75.55, 72.87, 69.95, 66.42, 63.60, 31.91, 26.56, 26.27, 25.14, 24.21, 14.04; *minor anomer* 162.00, 109.01, 107.43, 95.60, 74.05, 73.50, 72.54, 71.33, 67.05, 62.95, 33.59, 26.88, 25.92, 25.05, 24.07, 14.18; m/z HRMS (CI, NH₃), Found MH⁺, 345.1913 C₁₇H₂₉O₇ requires MH, 345.1913. **14**: $[\alpha]_D +49.4$ (c 0.57, CHCl₃); ν_{\max} (neat film) 3397, 1743 cm⁻¹; δ_C (67.8 MHz) *major anomer* 13.92, 25.29, 26.82, 26.96, 32.31, 66.19, 66.74, 69.86, 70.50, 71.25, 73.90, 94.32, 109.11, 109.28, 169.62; *minor anomer* 14.03, 24.18, 25.00, 25.29, 25.98, 30.89, 61.88, 67.07, 70.61, 72.28, 73.27, 73.81, 95.48, 109.28, 109.48, 169.62; m/z HRMS (CI, NH₃), Found MNH₄⁺, 364.1971, C₁₆H₃₀NO₈ requires MNH₄, 364.1971.
 11. Jiang, S.; Singh, G.; Wightman, R. H. *Chem. Lett.* **1996**, 67.
 12. Lactone **7** was prepared from D-lyxose in three steps (overall yield 61%) *via* isopropylideneation (acetone, conc. H₂SO₄, rt), silylation (TBDPSCl, Et₃N, DMAP, CH₂Cl₂, rt) and PCC oxidation.
 13. (a) Buchanan, J. G.; Smith, D.; Wightman, R. H. *Tetrahedron* **1984**, *40*, 119; (b) Csuk, R.; Dorr, P. *J. Carbohydr. Chem.* **1995**, *14*, 35.
 14. Lactone **9** was prepared from the PCC oxidation of 2,3,4-tri-*O*-benzyl-L-arabinopyranose. For the precursor, see: Finch, P.; Iskander, G. M.; Siriwardena, A. H. *Carbohydr. Res.*, **1991**, *210*, 319.
 15. McNicholas, P. A.; Batley, M.; Redmond, J.W. *Carbohydr. Res.* **1986**, *146*, 219.