

Total Synthesis of (-)-Decarestrictine D Through a Stereoselective Intramolecular Nozaki-Hiyama-Kishi Reaction

Ronaldo A. Pilli* and Mauricio M. Victor

Instituto de Química, Unicamp. Cx. Postal 6154, 13083-970 Campinas, SP Brasil


Received 11 March 1998; revised 13 April 1998; accepted 14 April 1998

Abstract: A concise total synthesis of (-)-decarestrictine D (**1**) from 1,3-propanediol and polyhydroxybutyrate (PHB) is described. The approach involves the stereoselective intramolecular Nozaki-Hiyama-Kishi coupling to construct the decanolide ring and to set the proper configuration at C-7. © 1998 Published by Elsevier Science Ltd. All rights reserved.

Decarestrictine D (**1**) is a 10-membered lactone isolated from *Penicillium corylophilum* and *P. simplicissimum*¹ and from *Polyporus tuberaster*² which has been demonstrated to inhibit cholesterol biosynthesis in HEP-G2 liver cells and *in vivo* studies with normolipidemic rats. Recently, Andrus and Shih reported the first total synthesis of decarestrictine D (**1**) which also established its absolute configuration.³

Here we would like to report our approach to decarestrictine D (**1**) which features a stereoselective construction of the 10-membered lactone and of its stereogenic center at C-7 through an intramolecular Nozaki-Hiyama-Kishi coupling reaction⁴ with vinylic iodide **2** (Scheme 1).


Scheme 1



We reasoned that the proper configuration at C-7, which requires the approach of the vinylic Cr(II) species to the *re* face of the aldehyde, would be favored due to the minimization of torsional and transannular interactions in the forming decanolide.⁵ Additionally, we hoped that such topology would be reinforced by the conformational bias imposed by the bulky OTBS groups at C-3 and C-4.

Our synthesis started with the reductive depolymerization of polyhydroxybutyrate (PHB), according to the procedure described by Seebach⁶, which afforded (-)-1,3-butanediol in 85% yield. Selective protection of the primary hydroxyl group with TBSCl gave (+)-**3** in 83% yield and >99% ee, as determined by chiral GC analysis.⁷

Scheme 2




Key: a) i. NaH, THF, 0°C; ii. TBSCl(91%); b) i. (COCl)₂, DMSO, CH₂Cl₂, -78°C; ii. Et₃N, -78°C->rt; c) (EtO)₂P(O)CHNaCH₃, THF, 0°C (70%, 2 steps); d) AD-mix α , CH₃SO₂NH₂, H₂O/ *tert*-BuOH (94%); e) TBSCl, DMF, imidazole (100%); f) DIBAL-H (2.0 equiv.), toluene, -95°C; g) CrCl₂ (12 equiv.), CHI₃ (4 equiv.), THF, 55°C; h) Jones reagent, acetone, 0°C (53%, 3 steps).

The preparation of (-)-4 required monosilylation of 1,3-propanediol (**5**)⁸ (91% yield), followed by Swern oxidation of **6** and Wittig olefination to give the unsaturated ester **7** as a 22:1 mixture of *E* and *Z* isomers (Scheme 2). The *E* isomer **7** was isolated in 70% yield after column chromatography on silica gel and submitted to Sharpless dihydroxylation⁹ (AD-mix α reagent in the presence of methanesulfonamide). Diol (-)-**8** was formed in 94% yield (91% ee by chiral GC analysis⁷) and it was fully protected as the corresponding TBS ether (-)-**9** (quantitative yield).

The final transformation of (-)-**9** to the C1-C6 fragment in (-)-**4** required extensive experimentation: DIBAL-H reduction of (-)-**9** to the corresponding aldehyde required 2.0 equivalents of the reducing agent at -95°C and the crude aldehyde was immediately submitted to modified Takai's conditions¹⁰ (CrCl₂, CHI₃, THF, 55°C) to afford *E*-**10** in 53% yield (2 steps). Selective desilylation (HF.pyr, pyridine, THF), followed by Jones oxidation afforded carboxylic acid (-)-**4** in 64% yield (2 steps). As the final steps in the

sequence above (desilylation and oxidation to the carboxylic acid) could conceivably be carried out sequentially with Jones reagent¹¹ and we had observed partial desilylation of (-)-10 during its purification by silica gel chromatography, we employed crude iodide (-)-10 in a one-pot desilylation-oxidation step with Jones reagent, under which conditions the carboxylic acid (-)-4 was obtained in 53% overall yield from (-)-9 after column chromatography on silica gel.

Scheme 3



Key: a) i. 2,4,6-Cl₃C₆H₂COCl, Et₃N, THF; ii. (-)-3, DMAP (2 equiv.), C₆H₆ (83%); b) HF.pyr, C₅H₅N, THF; c) Dess-Martin periodinane, CH₂Cl₂, H₂O; d) CrCl₂-0.5% NiCl₂ (15 equiv.), 5x10⁻³ M in DMF (30% yield from (-)-11; e) TBAF, HF, CH₃CN (80%).

At this stage the two requisite fragments were coupled using Yamaguchi's protocol¹² to afford (-)-11, in 83% yield. Attempts to purify by column chromatography on silicagel the alcohol resulting from the deprotection of (-)-11 with HF.pyr led to extensive migration of the acyl moiety from the secondary hydroxyl group to the primary one. Therefore, we employed the crude mixture in the Dess-Martin oxidation step to afford 2 which was immediately submitted to the Nozaki-Hiyama-Kishi coupling reaction in DMF to give (-)-12 (30% yield, 3 steps) as a single diastereoisomer, as determined by capillary GC analyses and from the ¹H-NMR spectrum (300 MHz) of the crude reaction mixture (Scheme 3).¹³

The stereochemistry at C-7 in (-)-12 was revealed by NOE difference spectroscopy (4.3% enhancement of the H-7 signal upon irradiation of H-9) and by its conversion to (-)-decastrictine D (1), in 80% yield. The optical rotation ([α]_D = -70.9 (c 1.0, CHCl₃)) and the ¹H- and ¹³C-NMR spectra of the synthetic material¹⁴ fully agreed with those reported by Andrus and Shih.³

In summary, an efficient (13 steps, 6.3% overall yield from 1,3-propanediol) and stereoselective route to (-)-decastrictine D (1) was developed with a reduced number of purification steps. Studies are

currently underway aiming to probe the structural features controlling the stereochemical outcome of the Nozaki-Hiyama-Kishi coupling as applied to the formation of 10-membered lactones.

Acknowledgements. The authors gratefully acknowledge financial support from FINEP (Brasil), Fapesp (Brasil), CNPq (Brasil), Volkswagen Stiftung (Germany) and IFS(Sweden).

References and Notes.

1. (a) Grabley, S.; Granzer, E.; Hütter, K.; Ludwig, D.; Mayer, M.; Thiericke, R.; Till, G.; Wink, J.; Phillips, S.; Zeeck, A. *J. Antibiotics* **1992**, *45*, 56. (b) Göhr, A.; Zeeck, A.; Hütter, K.; Kirsch, R.; Kluge, H.; Thiericke, R. *J. Antibiotics* **1992**, *45*, 66. (c) Grabley, S.; Hamann, P.; Hütter, K.; Kirsch, R.; Kluge, H.; Thiericke, R.; Mayer, M.; Zeeck, A. *J. Antibiotics* **1992**, *45*, 1176.
2. Ayer, W. A.; Sun, M.; Browne, L. M.; Brinen, L. S.; Clardy, J. *J. Nat. Prod.* **1992**, *55*, 649.
3. Andrus, M. B.; Shih, T.-L. *J. Org. Chem.* **1996**, *61*, 8780.
4. (a) Cintas, P. *Synthesis* **1992**, 248. (b) Kishi, Y. *Pure & Appl. Chem.* **1992**, *64*, 343.
5. (a) Still, W. C.; Galynger, I. *Tetrahedron* **1981**, *37*, 3981. (b) Rossseau, G. *Tetrahedron* **1995**, *51*, 2777.
6. Seebach, D.; Züger, M. *Helv. Chem. Acta* **1982**, *65*, 495.
7. Column: heptakis(2,6-methyl-3-pentyl)- β -cyclodextrine 20% in OV 1701, 25m, 0.25mm id.
8. McDougal, P. G.; Rico, J. G.; Oh, Y. I.; Condon, B. C. *J. Org. Chem.* **1986**, *51*, 3388.
9. Sharpless, H. B.; Amberg, W.; Bennani, Y. L.; Crispino, G. A.; Hartung, J.; Jeong, K.-S.; Kwong, H.-L.; Morikawa, K.; Wang, Z.-M.; Xu, D.; Zhang, X.-L. *J. Org. Chem.* **1992**, *57*, 2768.
10. Takai, K.; Nitta, K.; Utimoto, K. *J. Am. Chem. Soc.* **1986**, *108*, 7408.
11. Evans, P. A.; Roseman, J. D.; Garber, L. T. *Synth. Commun.* **1996**, *26*, 4685.
12. Inanaga, J.; Hirata, K.; Saeiki, H.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 1989.
13. When the Nozaki-Hiyama step in the above sequence was carried out in DMSO the formation of a 2:1 mixture of (-)-**12** and its C-7 epimer was observed.
14. ¹H NMR (500MHz, CD₃OD) of (-)-**1**: δ 5.83 (1H, ddd, J=15.9Hz, J=9.3Hz, J=1.5Hz), 5.74 (1H, dd, J=15.9Hz, J=3.1Hz), 5.17 (1H, dqd, J=11.3Hz, J=6.5Hz, J=1.6Hz), 4.19 (1H, ddd, J=4.5Hz, J=3.2Hz, J=1.5Hz), 4.07 (1H, ddd, J=10.7Hz, J=9.3Hz, J=3.4Hz), 3.94 (1H, ddd, J=6.8Hz, J=4.6Hz, J=2.4Hz), 2.59 (1H, dd, J=14.0Hz, J=2.3Hz), 2.31 (1H, dd, J=14.1Hz, J=6.9Hz), 1.85 (1H, ddd, J=13.9Hz, J=3.6Hz, J=1.5Hz), 1.72 (1H, dt, J=13.9Hz, J=11.2Hz), 1.21 (3H, d, J=6.7Hz); ¹³C NMR (75MHz, CDCl₃) of (-)-**1**: δ 175.3, 133.9, 130.1, 73.9, 72.5, 72.2, 68.2, 42.9, 33.0, 21.0.