



Pergamon

Tetrahedron Letters 39 (1998) 5903–5906

TETRAHEDRON
LETTERS

NEIGHBORING GROUP EFFECTS IN THE REGIOSELECTIVE CYCLIZATION OF VICINAL *trans*-1,2-BROMOHYDRINS TO EPOXIDES

Fengrui Lang, Darren J. Kassab, and Bruce Ganem*


Department of Chemistry and Chemical Biology
Baker Laboratory, Cornell University
Ithaca, NY 14853-1301 U. S. A.

Received 5 May 1998; revised 13 May 1998; accepted 8 June 1998


Abstract: Bromocyclopentitols and amino (or amido) bromocyclopentitols having a C-Br bond *trans* to two different vicinal hydroxyl groups show selectivity in base-promoted epoxide formation. The role of adjacent polar substituents in directing bromohydrin cyclization is discussed. © 1998 Elsevier Science Ltd. All rights reserved.

Epoxides are versatile functional groups in synthesis, and can generally be prepared from alkenes either directly, using peracids, or indirectly, by HOBr addition and cyclization of the corresponding bromohydrin.¹ The two methods are stereochemically complementary, with the bromohydrin route providing access in cyclic systems to the sterically more hindered epoxide.² Adjacent polar groups, especially allylic alcohol and amide NH groups can direct peracid-mediated alkene epoxidations.³ Proximal hydroxy and acetoxy groups can also affect the course of epoxide ring opening.⁴ However, the effect of nearby polar substituents on the cyclization of *trans*-1,2-bromohydrins to epoxides has not been investigated.

In connection with synthetic studies on trehazolin,^{5,6} we had occasion to prepare a number of bromocyclopentitols having a C-Br bond *trans* to two different vicinal hydroxyl groups. With their overlapping 1,2-bromohydrins, such systems are capable of forming regioisomeric epoxides. However, in each instance we observed high levels of selectivity favoring nucleophilic attack by the hydroxyl group vicinal to the X-group shown.



The regiochemistry of bromohydrin cyclization was examined in a series of bromocyclopentitols embodying the requisite (1,3/2)-dihydroxybromocycloalkane unit. Bromocyclopentitols **2**, **4**, and **6** having an additional hydroxy, amido, or amino substituent *cis* to one of the bromohydrin OH groups were prepared as shown in Scheme 1. Reaction of the known⁷ epoxide **1** with 48% HBr in Et₂O afforded **2** in 95% yield. Amidoalcohol **3**, prepared by the reaction of (benzyloxymethyl)cyclopentadiene⁸ with R-mandelohydroxamic acid and Bu₄NIO₄ following a published protocol,⁹ afforded **4**⁶ upon treatment with N-bromosuccinimide in wet 1,4-dioxane. Hydrolysis of **4** (0.5 N HCl) generated **6** in 98% yield.




(a) 48% HBr, Et₂O, (b) Na₂CO₃-CH₃OH, (c) NBS, 20:1 dioxane-H₂O; (d) 0.5 N HCl, CH₃OH, reflux; (e) 0.5 N NH₃-H₂O, (f) (CCl₃CO)₂O, pyr-CH₂Cl₂; (g) OxoneTM, H₂O-acetone.

Scheme 1

Cyclization of **2** and **4** to the corresponding epoxide was achieved by stirring a methanol solution of each bromohydrin with powdered Na_2CO_3 for 16-24 h. Cyclization of **6** was performed using 0.5 *N* $\text{NH}_3\text{-H}_2\text{O}$ for 2 h. Under these kinetically-controlled conditions, each bromohydrin efficiently formed a single epoxide (**1**, **5**, **7**) whose identity was conclusively established by comparison with a reference sample. Authentic **5**¹⁰ was obtained by *syn*-epoxidation of **3** with *m*-chloroperoxybenzoic acid (MCPBA, CH_2Cl_2 , rt). The structure of epoxide **7** was confirmed by comparing its trichloroacetylated derivative **9**¹¹ with an authentic sample prepared by *syn*-epoxidation of **8** (Scheme 1).

Earlier studies indicate the complex nature of polar substituent effects on epoxide opening and rearrangement reactions.¹² The related bromohydrin cyclizations reported here might involve ring conformational change, alkoxide formation, or nucleophilic attack as the rate-determining step (see Scheme 2), with each of those processes potentially affected by a vicinal alcohol, amine, or amide group. While intramolecular hydrogen bonding¹³ has been shown to increase the acidity of hydrogen-bond-accepting OH groups,¹⁴ cyclization of **2** in either 9:1 or 1:1 methanol:water also gave **1** as the exclusive product, indicating that internal H-bonding is not a factor in selectivity.



Scheme 2

Additional studies are needed to elucidate the role of structural, conformational, and inductive effects on bromohydrin closure. The methodology described here may find utility in the synthesis of bioactive cyclic polyols and aminocyclitols, which are components of a variety of potent antibiotics (e.g. kanamycins, streptomycins, neomycins) and glycosidase inhibitors (trehalosin, allosamidin, mannosatin).

ACKNOWLEDGMENT: We thank the National Institutes of Health (GM 35712) for generous financial assistance, and Professors A. Vasella and L. Liebeskind for helpful discussions. Support of the Cornell NMR Facility has been provided by the NSF (CHE 7904825; PGM 8018643) and NIH (RR02002).

REFERENCES AND NOTES

1. Berti, G in *Topics Stereochem.* **1973**, *7*, 93-251.
2. (a) Girard, J. P.; Vidal, J. P.; Granger, R.; Rossi, J. C.; Chapat, J. P. *Tetrahedron Lett.* **1974**, 943-946; (b) for a recent example see: Garofalo, A.; Hursthouse, M. B.; Abdul Malik, K. M.; Olivo, H. F.; Roberts, S. M.; Sik, V. *J. Chem. Soc. Perkin Trans. I* **1994**, 1311-1321.
3. (a) Narula, A. S. *Tetrahedron Lett.* **1983**, *24*, 5421-5424; (b) Ohfune, Y.; Kurokawa, N. *Tetrahedron Lett.* **1984**, *25*, 1587-1590; (c) Shaw, K. J.; Luly, J. R.; Rapoport, H. *J. Org. Chem.* **1985**, *50*, 4515-4523; (d) Askin, D.; Wallace, M. A.; Vacca, J. P.; Reamer, R. A.; Volante, R. P.; Shinkai, I. H. *J. Org. Chem.* **1992**, *57*, 2771-2773.
4. (a) Campion, T. A.; Morrison, G. A.; Wilkinson, J. B. *J. Chem. Soc. Perkin Trans. I* **1976**, 2508-2516; (b) Baldwin, D. A.; Hanson, J. R.; Raines, D. *J. Chem. Soc. Perkin Trans. I* **1979**, 344-348.
5. Goering, B. K.; Li, J.; Ganem, B. *Tetrahedron Lett.* **1995**, *36*, 8905-8908.
6. Li, J.; Lang, F.; Ganem, B. *J. Org. Chem.*, **1998**, *63*, 3403-3410.
7. Madhavan, G. V. B.; Martin, J. C. *J. Org. Chem.* **1986**, *51*, 1287-1293.
8. Corey, E. J.; Koelliker, U.; Neuffer, J. *J. Am. Chem. Soc.* **1971**, 1489-1490.
9. King, S. B.; Ganem, B. *J. Am. Chem. Soc.* **1994**, *116*, 562-570.
10. For **5**: ^1H NMR (CDCl_3 , 300 MHz) δ 7.39-7.24 (m, 10 H), 5.07 (d, 1 H, $J = 8$ Hz), 4.45 (s, 2 H), 4.31 (br t, 1 H), 4.11 (m, 1 H), 3.65-3.45 (m, 4 H), 1.65, (m, 1 H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 172.4, 139.2, 137.7, 128.8, 128.6, 128.4, 127.8, 127.6, 126.7, 74.3, 73.8, 73.4, 68.6, 57.3, 56.2, 50.6, 45.3.
11. For **9**: ^1H NMR (CDCl_3 , 300 MHz) δ 7.49-7.32 (m, 5 H), 5.32 (br d, 1 H), 4.64-4.48 (m, 3 H), 3.89, 3.78 (br m, each 1 H), 3.55 (m, 2 H), 2.09 (m, 1 H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 161.6, 137.8, 137.0, 131.4, 128.5, 127.8, 127.7, 78.3, 73.5, 70.3, 58.3, 56.3.
12. Buchanan, J. G.; Sable, H. Z. in *Selective Organic Transformations*, Thyagarajan, B. S.; Ed.; Wiley-Interscience, New York, **1972**; Vol. 2, 1-95.
13. Tichy, M. in *Adv. Org. Chem. Methods and Results* **1965**, *5*, 115-298.
14. (a) Muddasani, P. R.; Bernet, B.; Vasella, A. *Helv. Chim. Acta* **1994**, *77*, 334-350; (b) Weber, M.; Vasella, A. *Helv. Chim. Acta* **1997**, *80*, 2352-2367; (c) Griller, D.; Liu, M. T. H.; Scaiano, J. C. *J. Am. Chem. Soc.* **1982**, *104*, 5549-5551.