

DMAP-Catalyzed Hydroxymethylation of 2-Cyclohexenones in Aqueous Medium Through Baylis-Hillman Reaction

Farhat Rezgui and Mohamed Moncef El Gaied*

Laboratoire de Chimie Organique, Faculté des Sciences de Tunis
Campus Universitaire, 1060 Tunis, Tunisie. Fax: (1) 88 50 08

Received 1 April 1998; accepted 10 June 1998


Abstract: Reaction of 2-cyclohexenones **1a-d** with aqueous formaldehyde, catalyzed by DMAP in THF, affords the corresponding 2-(hydroxymethyl)-2-cyclohexenones **2a-d** in good yields.

© 1998 Elsevier Science Ltd. All rights reserved.

2-(Hydroxymethyl)-2-cyclohexenone **2a** is a potential intermediate for the synthesis of products having biological properties^{1,2}. For instance, acetate **3a** is effective against HELA cells growth³ while amines **3b** derivatives have an analgesic activity⁴. Furthermore, we have recently developed a simple and convenient method for the synthesis of allylic amines **3b**⁵ and bicyclic dienones **4a-b**⁶ via the reaction of synthon **3a** and a variety of nucleophiles. However, up to now, 2-(hydroxymethyl)-2-cyclohexenone **2a** was prepared in two steps and in small-scale by reaction of 1,3-cyclohexanedione with paraformaldehyde in the presence of BF₃ etherate, followed by Dibal-H reduction of intermediate 1,3-dioxine vinylogous ester⁷.

We assume that, under the influence of a suitable catalyst, Baylis-Hillman^{8,9} coupling reaction of cyclic enone **1a** with formaldehyde, which has been reported rarely¹⁰, would be a simple method for preparing alcohol **2a**. We report herein, a new and efficient procedure for direct α -hydroxymethylation of enones **1a-d** in one step, in aqueous medium and relatively in much higher scale.

Reaction of **1a** and HCHO in THF, catalyzed by 1,4-diazabicyclo[2.2.2]octane (DABCO)¹¹⁻¹³, commonly used as a catalyst in the Baylis-Hillman reaction, led to the recovery of the starting materials. However, we have found that, at room temperature and in the presence of a catalytic amount (10 mol %) of 4-(dimethylamino)pyridine (DMAP) or 4-(pyrrolidino)pyridine (PPY), α,β -unsaturated ketone **1a** reacted with aqueous formaldehyde (2 equiv) in THF, to give 2-(hydroxymethyl)-2-cyclohexenone **2a** in 75-82 % yield.




	a	b	c	d	3	a	b	4	a	b
R ¹	H	H	Me	H	Y	OAc	NR ₂	R	H	Me
R ²	H	Me	Me	Ph						
Yield (%)	82	76	68	71						

To our knowledge, this process is the first Baylis-Hillman reaction of cyclic enones with aldehydes catalyzed by a pyridine derivative. In order to explore the scope and limitations of the reaction described above, we have prepared a variety of 5-substituted-2-cyclohexenones **1b-d**¹⁴⁻¹⁵ and we have studied their catalytic

hydroxymethylation with aqueous formaldehyde. Our experimental results and the required compounds **2b-d** obtained in 68-76 % yields, are summarized in the table below:

α,β -enones (mmol)	HCHO (x equiv)	DMAP (molar %)	T (°C)	Time (day)	Products	Yield (%)
1b (20)	4	15	25	3	2b	76
1c (20)	6	20	60	5	2c	68
1d (12)	4	20	25	6	2d	71

It is noteworthy that in the presence of 0.2 equiv of DMAP, the treatment of β -substituted 2-cyclohexenones **5a-c** with a large excess (6 equiv) of aqueous formaldehyde at 60 °C for 10 days, led to the recovery of the starting materials; this may be explained by the C (β) steric effects which prevent, in the first step of the Baylis-Hillman reaction, the Michael addition of DMAP to α,β -enones **5a-c**¹⁶⁻¹⁷.



A representative experimental procedure for α -hydroxymethylation of enone **1a** is presented below:

A mixture of compound **1a** (24 g, 0.25 mol), 30 % aqueous formaldehyde (50 mL, 0.5 mol), 50 mL of THF and DMAP (3,05 g, 25 mmol), was stirred for 15 h at room temperature, then acidified dropwise with 1.5 N aqueous HCl. The reaction mixture was extracted with methylene chloride, washed successively with NaHCO₃ and brine. After usual work up, chromatography of the crude product on a silica gel column, using ether-methylene chloride (3:1) as eluent, gave a 82 % yield of pure **2a** as yellow oil.

IR (CHCl₃): 1670, 3580 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 6.90 (t, J = 4, 1H); 4.07 (s, 2H); 2.67-2.23 (m, 4H); 2.13-1.83 (m, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 200.4; 146.8; 138.4; 60.8; 38.2; 25.7; 22.8; MS (m/e): 39 (100); 41 (88); 55 (82); 69 (82); 97 (44); 111 (29); 126 (M⁺, 60).

In conclusion, 2-(hydroxymethyl)-2-cyclohexenones **2a-d** were prepared in one step and in mild conditions from the corresponding α,β -enones **1a-d** in good yields. This convenient methodology is inexpensive and can be successfully used in large-scale syntheses of cyclic 2-(hydroxymethyl) enones. Extension of this process to coupling several substituted cyclic α,β -enones with a variety of electrophiles, is in progress in our laboratory.

References

- Smith, III, A. B.; Branca, S. J.; Pilla, N. N.; Guaciaro, M. A. *J. Org. Chem.* **1982**, *47*, 1855-1869 and references cited in.
- Smith, III, A. B.; Branca, S. J.; Guaciaro, M. A.; Wovkulich, P. M.; Korn, A. *Org. Synth. Coll. Vol. VII*, **1990**, 271-275.
- Takao, I.; Kunimoto, K.; Masanobu, S.; Kiyohiro, N.; Takemitsu, N.; *Chem. Abstr.* **1987**, *106*, 49660m.
- Kato, T.; Ishii, H.; Kawai, K.; Sawa, Y. *Chem. Pharm. Bull.* **1984**, *32*, 2279-2289.
- Rezgui F.; EL Gaïed M. M. *J. Soc. Chim. Tunisie* **1993**, *3*, 293-298; *Chem. Abstr.* **1994**, *121*, 8725c.
- Rezgui F.; EL Gaïed M. M. *Tetrahedron* **1997**, *53*, 15711-15716.
- Smith, III, A. B.; Dorsey, B. D.; Ohba, M.; Lupo, A. T., Jr.; Malamas, M. S. *J. Org. Chem.* **1988**, *53*, 4314-4325.
- Baylis, A. B.; Hillman, M. E. D. German Patent 2155113, 1972; *Chem. Abstr.* **1972**, *77*, 34174q.
- For reviews of the Baylis-Hillman reaction, see: a) Drewes, S. E.; Roos, G. H. P. *Tetrahedron* **1988**, *44*, 4653-4670. b) Basavaiah, D.; Dharma Rao, P.; Suguna Hyma, R. *Tetrahedron* **1996**, *52*, 8001-8062.
- Kabat, M. M.; Kiegiel, J.; Cohen, N.; Toth, K.; Wovkulich, P. M.; Uskokovic, M. R. *J. Org. Chem.* **1996**, *61*, 118-124.
- Hwu, J. R.; Hakimelahi, G. H.; Chou, C.-T. *Tetrahedron Lett.* **1992**, *33*, 6469-6472 and references cited in.
- Roth, F.; Gygax, P.; Frater, G. *Tetrahedron Lett.* **1992**, *33*, 1045-1048.
- Rafel, S.; Leahy, J. W. *J. Org. Chem.* **1997**, *62*, 1521-1522.
- House, H. O.; Fischer, W. F., Jr. *J. Org. Chem.* **1968**, *33*, 949-956.
- Ames, G. R.; Davey, W. *J. Chem. Soc.* **1957**, 3480-3487.
- Piancatelli, G.; Scettri, A.; D'auria, M. *Synthesis* **1982**, *4*, 245-258.
- Clark, R. D.; Heathcock, C. H. *J. Org. Chem.* **1976**, *41*, 636-643.