



Oxoanion Recognition by a Thiuronium Receptor

Woon-Seok Yeo and Jong-In Hong*

Department of Chemistry, Seoul National University, Seoul 151-742, Korea

Received 8 July 1998; accepted 28 August 1998

Abstract: The relative binding affinities of various oxoanions for a simple bis(thiuronium) receptor **2** derived from 1,3-bis(aminomethyl)benzene were determined to be $\text{ArPO}_3^{2-} > \text{ArOPO}_3^{2-} > \text{H}_2\text{PO}_4^- > \text{ArCOO}^- > \text{ArP}(\text{OH})\text{O}_2^- > \text{ArOP}(\text{OH})\text{O}_2^- > \text{ArSO}_3^-$ in strongly solvating solvent like DMSO on the basis of ^1H NMR titrations. The relative binding affinity of **2** for various oxoanions can be explained on the basis of the Brønsted basicity of substrate oxoanions and complex structures.

© 1998 Elsevier Science Ltd. All rights reserved.

In biological systems, the selective recognition of oxoanion substrates such as phosphate and sulfate often takes place by a combination of hydrogen bonds and electrostatic interactions.¹ Synthetic receptors for the biologically relevant oxoanions have recently been developed using these binding forces and others in strongly solvating solvents such as DMSO, methanol, or water by several research groups.²⁻⁵ In particular, bis(urea) and bis(thiourea) groups have been shown to bind dihydrogenphosphate (H_2PO_4^-) selectively over various other anions.^{4a} Recently, Smith and co-workers showed that polarization of a urea group by intramolecular coordination with a Lewis acidic boronate as shown in **1a** and **1b**, improved acetate binding affinities by up to 3.0 kcal/mol because of larger host dipole moment and the increased positive surface potential at the urea NH residues induced by internal Lewis acid coordination.⁶ Therefore we reasoned that the thiuronium group⁷ generated by S-alkylation of the thiourea group would possess relatively larger dipole moment and enhanced acidity of thiourea NH residues and therefore can function as a better oxoanion binder compared to the thiourea group.⁸

In this paper, we report on the complexation behavior of thiuronium receptors **2** and **3** for various oxoanions in DMSO, which is a strong hydrogen bonding acceptor and strongly precludes the formation of hydrogen bond based complex.




Fig. 1


Table 1. Association constants of **2** with oxoanions in DMSO- d_6 ^a

Oxoanion substrate	pK_a of substrate•H(approx.) ^b	$K_a(M^{-1})$ ^c
PhSO ₃ ⁻	-6	~5 ^d
PhOP(OH)O ₂ ⁻	1	50(3)
PhP(OH)O ₂ ⁻	2	150(108)
H ₂ PO ₄ ⁻	2	1080(180)
PhCOO ⁻	4	590(130)
PhOPO ₃ ²⁻	6	3700
PhPO ₃ ²⁻	7	4350

a. Oxoanions were used as their *n*-Bu₄N⁺ salts.

b. See refs. 4a and 4b for sources of pK_a data.

c. The number in parenthesis indicates 1:2 complexation constant.



d. Not accurately determined.

Receptors **2** and **3** were prepared from reaction of *p*-nitrobenzyl bromide with the corresponding bis(thiourea) and mono(thiourea) precursors,^{4a} respectively.⁹ The association constants of sulfonate, carboxylates, phosphate, and phosphonates for thiuronium receptors were determined by proton NMR titrations in DMSO- d_6 (Table 1). For solubility reasons, counteranions of anionic guests were replaced by tetrabutylammonium cations.

The new thiuronium receptors bound oxoanions through hydrogen bonding and electrostatic interaction. For the receptor **2**, upfield shifts (~0.5 ppm) of the two different benzylic proton (ArCH₂NH- and Ar'CH₂S-) resonances were observed upon complexation with anionic substrates, while the two different kinds of NH proton peaks rapidly disappeared with increasing anionic substrate concentrations. The relative binding strengths correlate reasonably well with the net charge on the anionic guest, indicating electrostatic interactions as major binding force. Thus, in DMSO- d_6 the dianionic substrates were bound more tightly than their monoanionic counterparts. But that correlation does not explain the difference among substrates possessing the same net charge. The best correlation for the binding strengths of the substrates is with their Brønsted basicity,^{4a,4b} as shown in Table 1. Thus, benzenesulfonate with the weakest basicity is the weakest binder, and the phosphate and phosphonate dianions with the strongest basicity are the strongest binders.

The one exceptional result was the degree of complexation selectivity for H₂PO₄⁻. The observed selectivity for H₂PO₄⁻ can be explained in terms of the complexation geometry. Because PhCOO⁻ has only two oxygens interacting with the receptor, it must have a different binding mode. The fact that H₂PO₄⁻ displayed stronger binding affinity than PhCOO⁻ in spite of its substantially smaller basicity indicated that H₂PO₄⁻ was not bound in a similar manner to PhCOO⁻.^{4a} Based on the structural similarity to bis(guanidinium)/H₂PO₄⁻,^{2h} bis(thiourea)/H₂PO₄⁻ and bis(thiourea)/PhCOO⁻ complexes,^{4a} we propose structures **4** and **5** for the complexes of bis(thiuronium) **2** with H₂PO₄⁻ and PhCOO⁻, respectively. This hypothesis was also supported by ¹H NMR titration spectra. In fact, the aromatic region signals of the receptor **2** were broad, which indicated that a bis(thiuronium) salt **2** was asymmetrical but slightly distorted in substrate-free condition since two bromides in the anion receptor **2** could play a role of substrate. However, the fact that these signals were sharpened with increasing substrate concentration illustrated that receptor-substrate complex structure was symmetrical as

depicted in **4**. The weaker binding of H_2PO_4^- to mono(thiuronium) receptor **3** (340 M^{-1}) suggested that both thiuronium groups of **2** were involved in complexation as shown in **4**.




Fig. 2

In spite of the same basicity, large difference in K_a between $\text{PhP}(\text{OH})\text{O}_2^-$ and H_2PO_4^- induced an abstruse problem. It was not clear why phenyl was deleterious, but we might argue that the diminished binding strength of $\text{PhP}(\text{OH})\text{O}_2^-$ could be rationalized by the different number of oxygens which could interact with the receptor **2**.

Comparing to thiourea groups as an oxoanion binder,^{4a,4c} thiuronium groups turned out to be a relatively stronger binder for oxoanion substrates. For example, a mono(thiuronium) receptor **3** bound acetate in $\text{DMSO-}d_6$ more strongly ($K_a = 800 \text{ M}^{-1}$) than a mono(thiourea) ($K_a = 340 \text{ M}^{-1}$) and a bis(thiourea) ($K_a = 470 \text{ M}^{-1}$)^{4a}, respectively. However, it showed much poorer binding affinity than guanidinium based ones, although it was very similar to those in structure. This fact implied the cationic power of thiuronium was weaker than that of guanidinium, which was well reasoned by relatively large size of sulfur. Since anion basicity is a measure of the propensity to form hydrogen bonds, the observed stability trend suggested that the major importance in the formation of thiuronium-oxoanion complex was hydrogen bond formation rather than electrostatic interaction.

UV absorption experiments in 1,2-dichloroethane showed that the absorbance of **2** above 280 nm increased upon complexation with H_2PO_4^- ($K_a = 34000 \text{ M}^{-1}$). In the low H_2PO_4^- concentration range (< 2.5 equiv.), a clear isosbestic point was observed. However, the deviation from the clear isosbestic point at high substrate concentration indicated the formation of some amount of 1:2 complexes from one molecule of **2** and two H_2PO_4^- beside the dominant 1:1 complex.

In summary, we have shown that the thiuronium salt can function as an oxoanion receptor and a bis(thiuronium) group can strongly bind H_2PO_4^- in strongly solvating solvent like DMSO. The relative binding affinity of a bis(thiuronium) receptor **2** for various oxoanions can be explained on the basis of the Brønsted basicity of substrate oxoanions and complex structures.

Acknowledgements. We thank Ministry of Education (BSRI-97-3416) and KOSEF (Grant No. 961-0302-008-2) for financial support.

References and Notes

- (a) Luecke, H.; Quioco, F. A. *Nature* **1990**, *347*, 402-406. (b) Mangani, S.; Ferraroni, M.; Orioli, P. *Inorg. Chem.* **1994**, *33*, 3421-3423. (c) Malashkevich, V.; Kammerer, R. A.; Efimov, V. P.; Schulthess, T.;

Engel, J. *Science* **1996**, *274*, 761-765. (d) Kanyo, Z. F.; Christianson, D. W. *J. Biol. Chem.* **1991**, *266*, 4246-4268.

2. (a) Hosseini, M. W.; Lehn, J.-M. *Helv. Chim. Acta.* **1988**, *71*, 749-756. (b) Sessler, J. L.; Cyr, M.; Furuta, H.; Král, V.; Mody, T.; Morishima, T.; Shionoya, M.; Weghorn, S. *Pure Appl. Chem.* **1993**, *65*, 393-398. (c) Berger, M.; Schmidtchen, F. P. *J. Am. Chem. Soc.* **1996**, *118*, 8947-8948. (d) Bencini, A.; Bianchi, A.; Paoletti, P.; Paoli, P. *Pure Appl. Chem.* **1993**, *65*, 381-386. (e) Worm, K.; Schmidtchen, F. P.; Schier, A.; Schäfer, A.; Hesse, M. *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 327-329. (f) Jung, Y.-G.; Yeo, W.-S.; Lee, S. B.; Hong, J.-I. *Chem. Commun.* **1997**, 1061-1062. (g) Kimura, E.; Kuramoto, Y.; Koike, T.; Fujioka, H.; Kodama, M. *J. Org. Chem.* **1990**, *55*, 42-46. (h) Dixon, R. P.; Geib, S. J.; Hamilton, A. D. *J. Am. Chem. Soc.* **1992**, *114*, 365-366. (i) Yeo, W.-S.; Hong, J.-I. *Tetrahedron Lett.* **1998**, *39*, 3769-3772.

3. (a) Reetz, M. T.; Niemeyer, C. M.; Harms, K. *Angew. Chem. Int. Ed. Engl.* **1991**, *30*, 1472-1474. (b) Zheng, Z.; Yang, X.; Knobler, C. B.; Hawthorne, M. F. *J. Am. Chem. Soc.* **1993**, *115*, 5320-5321. (c) Newcomb, M.; Horner, J. H.; Blanda, M. T.; Squatrito, P. J. *ibid.* **1989**, *111*, 6294-6301. (d) Konishi, K.; Yahara, K.; Toshishige, H.; Aida, T.; Inoue, S. *ibid.* **1994**, *116*, 1337-1344. (e) Rudkevich, D. M.; Verboom, W.; Brzozka, Z.; Palys, M. J.; Stauthamer, W. P. R. V.; van Hummel, G. J.; Franken, S. M.; Harkema, S.; Engbersen, J. F. J.; Reinhoudt, D. N. *ibid.* **1994**, *116*, 4341-4351. (f) Jung, M. E.; Xia, H. *Tetrahedron Lett.* **1988**, *29*, 297-300.

4. (a) Nishizawa, S.; Buehlmann, P.; Iwao, M.; Umezawa, Y. *ibid.* **1995**, *36*, 6483-6486. (b) Kelly, T. R.; Kim, M. H. *J. Am. Chem. Soc.* **1994**, *116*, 7072-7080. (c) Fan, E.; Arman, S. A. V.; Kincaid, S.; Hamilton, A. D. *ibid.* **1993**, *115*, 369-370. (d) Garcia-Tellado, F.; Goswami, S.; Chang, S. K.; Geib, S. J.; Hamilton, A. D. *ibid.* **1990**, *112*, 7393-7394. (e) Smith, P. J.; Reddington, M. V.; Wilcox, C. S. *Tetrahedron Lett.* **1992**, *33*, 6085-6088. (f) Beer, P. D.; Szemes, F.; Balzani, V.; Salà, C. M.; Drew, M. G. B.; Dent, S. W.; Maestri, M. *J. Am. Chem. Soc.* **1997**, *119*, 11864-11875. (g) Jeong, K.-S.; Park, J. W.; Cho, Y. L. *Tetrahedron Lett.* **1996**, *37*, 2795-2798.

5. For positively charged receptors using electrostatic interactions, see refs 2a-2i. For neutral receptors having Lewis acid centers, see refs 3a-3f. For neutral receptors having only hydrogen bonding binding sites, see refs 4a-4g.

6. (a) Hughes, M. P.; Smith, B. D. *J. Org. Chem.* **1997**, *62*, 4492-4499. (b) Hughes, M. P.; Shang, M.; Smith, B. D. *ibid.* **1996**, *61*, 4510-4511.

7. Thiuronium salts have been used as a reagent for the identification of organic acids. See ref 8.

8. (a) Donleavy, J. J. *J. Am. Chem. Soc.* **1936**, *58*, 1004-1005. (b) Dewey, B. T.; Sperry, R. B. *ibid.* **1939**, *61*, 3251-3252.

9. Selected spectral data for **2**: ^1H NMR (300 MHz, DMSO- d_6) δ 9.87 (br, 2H), 9.60 (br, 2H), 8.16 (d, $J = 8.3$ Hz, 4H), 7.58 (br d, 4H), 7.40-7.00 (m, 4H), 4.66 (s, 4H), 4.56 (s, 4H), 3.34 (m, 4H), 1.55-0.75 (m, 14H); FABMS 637 (M^{2+} -1). **3**: ^1H NMR (300 MHz, DMSO- d_6) δ 8.19 (d, $J = 8.5$ Hz, 2H), 7.55 (br d, 2H), 7.40-7.00 (m, 5H), 4.70-4.50 (m, 4H), 3.33 (m, 2H), 1.55-0.75 (m, 7H); FABMS 358(M^+).