

Extension of the Tandem Pauson-Khand Reaction to [5.6.6.5]Tetracycles: A Unique Entry into *cis*-Fused Decalins.

Michelle M. Bruendl,^a Scott G. Van Ornum,^a Tze-Ming Chan,^b and James M. Cook^{a*}


^aDepartment of Chemistry, University of Wisconsin-Milwaukee, Milwaukee, WI 53201

^bSchering Plough Research Institute, Kenilworth, NJ 07033

Received 10 November 1998; accepted 7 December 1998

Abstract: The tandem Pauson-Khand reaction has been extended to the synthesis of [5.6.6.5]dicyclopenta[*b,g*]decalins. This approach provides a unique entry into a *cis*-fused decalin-based ring system *via* an acyclic precursor. © 1999 Elsevier Science Ltd. All rights reserved.

The intramolecular Pauson-Khand reaction^{1,2} is one of the most versatile tools for the construction of bicyclo[3.3.0]octen-ones. Previous interest in the tandem version of this reaction has resulted in the synthesis of the parent [5.5.5.5]enone ring systems such as **1**; potential intermediates en route to 14 π polyquinenes represented by dicyclopenta[*a,f*]pentalene (**2**) and dicyclopenta[*a,e*]pentalene.³ We wish to report the first extension of this dicobalt octacarbonyl-mediated *tandem* process which has permitted facile entry into [5.6.6.5]dicyclopenta[*b,g*]decalins (see **3**) with a *cis*-fused decalin core.




There are several methods utilized for the formation of *cis*-fused decalin rings. Most commonly, inter- and intramolecular⁴ Diels-Alder reactions are employed, as well as anionic cycloadditions;⁵ samarium diiodide-promoted constructions are also known.⁶ In addition, *cis*-decalins have also been prepared *via* an intramolecular aldol reaction⁷ and an asymmetric Heck-type reaction.⁸ In contrast, this tandem Pauson-Khand strategy begins from acyclic molecules in which 6 carbon-carbon bonds are formed in a one-pot process, while providing an internal *cis*-decalin within the tetracyclic system.


The synthesis of the diene-diyne required for the tandem Pauson-Khand reaction began with bisalkyne **4**, available by addition of diethyl oxalate to a stirred suspension of trimethylsilyl propargyl bromide and zinc in THF with a catalytic amount of mercury(II)chloride.⁹ When the dialkyne ester **4** was stirred with allylmagnesium chloride the diene-diyne **5** was furnished in 88% yield (Scheme 1). The desired diol **6** was obtained by stirring **5**

with $K_2CO_3/MeOH$ (95%) and this was followed by protection of the hydroxyl groups *via* DMAP and TMSCl to furnish the *seco*-Pauson-Khand intermediate **7** (99%).

Scheme 1



Application of the Livinghouse photochemically-mediated Pauson-Khand protocol¹⁰ to the synthesis of the parent ring system of the dicyclopentalenenes had previously provided a convenient, semi-catalytic tandem reaction sequence. Similar reaction conditions were employed here [1 equiv. $Co_2(CO)_8$ added to a 0.10 M solution of **7** in DME under a CO atmosphere, irradiation with Q-Beam for 12 h]^{3c} to convert **7** into *cis*-decalin **8** with somewhat limited success (38% yield). The stereochemistry of the trimethylsilyl protected diol **8** at the 6-6 ring fusion was determined to be exclusively *cis* *via* 2D NMR techniques (Figure 1).¹¹ Removal of the silyl groups in **8** with TBAF hydrate in THF provided diol **3** in 89% yield.



Since only *cis*-fused decalins were formed in this process, it was felt the trimethylsilyl groups in **7** may have retarded the reaction progress. For this reason, the Pauson-Khand reaction of the unprotected diol **6** was then carried out. Unfortunately, the method of catalytic photolysis furnished low yields of the [5.6.6.5]dicyclopenta[*b,g*]decalin tetracycle **3a/b** [(26%), Table 1, entry *b*].


**8a** H_a, H_b , trans


Figure 1

As shown in Table 1, a variety of reaction conditions were investigated in attempts to increase the reaction yield from the initial photochemical process. Recently, Livinghouse¹² has reported that the thermal (60°C) version of the catalytic Pauson-Khand procedure (entry *c*, Table 1) occurred in yields comparable to the photolytic catalytic method.¹⁰ In the study outlined below, yields for the tetracyclic products were also similar to the light-promoted reactions. The modified conditions of Schreiber,¹³ [2.5 equivalents of $\text{Co}_2(\text{CO})_8$ followed by excess *N*-methylmorpholine-*N*-oxide (NMO) in CH_2Cl_2] employed earlier in our laboratory,^{3a,b} were successfully employed here and the yield of **8** was increased to 53% (3:1 ratio of diastereomers at H_α). Importantly, diol **6** was also submitted to the same reaction conditions which provided tetracycle **3** in 52% yield. Success with this approach prompted the study of other solvents (see Table 1, entries *f-i*) but CH_2Cl_2 remained the solvent of choice. It was felt the polarity of CH_2Cl_2 contributed to the near-homogeneous nature of the NMO reaction in contrast to the heterogeneous mixture present when heptane was the solvent.

Table 1

Entry	Substrate	Reaction Conditions	% Yield	Product
<i>a</i>	OTMS 7	DME, hv	38	8
<i>b</i>	diol 6		26	3
<i>c</i>	diol 6	DME, 65°C	31	3
<i>d</i>	OTMS 7	CH_2Cl_2 , NMO ¹⁴	53	8
<i>e</i>	diol 6		52	3
<i>f</i>	diol 6	CH_3CN , NMO	44	3
<i>g</i>	diol 6	$\text{CF}_3\text{CH}_2\text{OH}$, NMO	33	3
<i>h</i>	diol 6	1:1 heptane/ CH_2Cl_2 , NMO	<5	3
<i>i</i>	diol 6	heptane, NMO	0	3
<i>j</i>	acetal 10	CH_2Cl_2 , NMO	30	11
<i>k</i>	acetal 10	DME, hv	45	11

Finally, the success of the tandem Pauson-Khand reaction (53%) in the sterically hindered case, coupled with no yield increase when the diol itself was employed, prompted the construction of a fixed diene-diyne conformation *via* a benzylidene acetal, which substantially reduced rotation about the diol carbon-carbon bond. Treatment of diol **5** with *p*-anisaldehyde dimethyl acetal in refluxing toluene, (catalytic amount of PTSA), readily provided acetal **9** in 90% yield. The trimethylsilyl groups were removed on stirring with $\text{K}_2\text{CO}_3/\text{MeOH}$ (90%) to afford diene-diyne **10**. This constrained, eclipsed diene-diyne was subjected to the NMO conditions for the tandem Pauson-Khand reaction to provide the *cis*-fused decalin **11** in 30% yield, isolated as a 1.2:1 mixture of acetal isomers (entry *j*). Similarly, reaction of acetal **10** *via* the photochemically-mediated Pauson-Khand procedure afforded 45% of the desired decalin **11** in a 1:1 ratio (entry *k*).



In summary, the scope of the tandem Pauson-Khand reaction has been extended to the [5.6.6.5] dicyclopenta[*b,g*]decalin ring system. A *cis*-fused decalin core was obtained exclusively from the formation of 6 carbon-carbon bonds in this one-pot process. This tandem Pauson-Khand reaction is unique in that it provides a *cis*-decalin ring system, moreover, the yield of each of the six carbon-carbon bonds formed in this procedure is at least 89%.

References:

- Schore, N. E. *Org. React.* John Wiley and Sons, N.Y., **1991**, 40, 1.
- (a) Khand, I. U.; Knox, G. R.; Pauson, P.L.; Watts, W. E. *J. Chem. Soc., Chem. Commun.* **1971**, 36.
(b) Khand, I. U.; Knox, G. R.; Pauson, P.L.; Watts, W. E. *J. Chem. Soc., Perkin Trans. 1.* **1973**, 975.
- (a) Van Ornum, S. G.; Cook, J. M. *Tetrahedron Lett.* **1996**, 37, 7185. (b) Van Ornum, S. G.; Cook, J. M. *Tetrahedron Lett.* **1997**, 38, 3657. (c) Van Ornum, S. G.; Bruendl, M. M.; Cook, J. M. *Tetrahedron Lett.* **1998**, 39, 6649.
- Rémy, S.; Jaeckel, F.; Olesker, A.; Lukacs, G. *Carbohydrate Res.* **1992**, 229, 167.
- Reul, R.; Hogan, K. T.; Deslongchamps, P. *Synlett* **1990**, 516.
- (a) Kito, M.; Sakai, T.; Haruta, N.; Shirahama, H.; Matsuda, F. *Synlett* **1996**, 1057. (b) Kito, M.; Sakai, T.; Yamada, K.; Matsuda, F.; Shirahama, H. *Synlett* **1993**, 158.
- Renard, P.-Y.; Lallemand, J.-Y. *Synlett* **1993**, 163.
- Sato, Y.; Sodeoka, M.; Shibasaki, M. *Chem. Lett.* **1990**, 1953.
- Scott G. Van Ornum, Ph.D. Thesis, University of Wisconsin-Milwaukee, 1998.
- Pagenkopf, B. L.; Livinghouse, T. *J. Am. Chem. Soc.* **1996**, 118, 2285.
- Major NOEs are shown here. Complete NMR results will be published elsewhere, in preparation.
- Belanger, D. B.; O'Mahony, D. J. R.; Livinghouse, T. *Tetrahedron Lett.* **1998**, 39, 7637.
- Shambayati, S.; Crowe, W. E.; Schreiber, S. L. *Tetrahedron Lett.* **1990**, 31, 5289.
- A typical procedure is as follows: A solution of the diene-diyne **7** (0.255 g, 0.704 mmol) and dicobalt octacarbonyl (0.602 g, 1.76 mmol, 2.5 equiv) in CH_2Cl_2 (10 mL) was stirred at rt under an atmosphere of carbon monoxide. After stirring for 2 h, examination of TLC indicated an absence of starting material and the reaction mixture was diluted with CH_2Cl_2 (10 mL) and *N*-methylmorpholine-*N*-oxide (0.825 g, 7.04 mmol, 10 equiv) was added portionwise. After stirring 6 h, additional NMO was added (0.400 g, 3.41 mmol) until no metal complexes were observed by TLC. The purple solution was concentrated to one-quarter of its original volume under reduced pressure and loaded onto a silica gel column and eluted (flash chromatography) with 55% EtOAc/hexane to provide tetracycles **8a** and **8b** (3:1) as off-white solids, 0.156 g (53%). **8a**: ^{13}C NMR (125 MHz, CDCl_3) δ 208.3, 207.7, 180.4, 179.6, 130.4, 129.2, 80.4, 77.2, 42.4, 41.6, 41.51, 41.48, 40.7, 40.3, 38.7, 37.0, 2.7, 2.5; LRMS (CI) *m/z* relative intensity 419 (*M*+1, 100), 329 (72). **8b**: ^{13}C NMR (75 MHz, CDCl_3) δ 207.9, 180.2, 129.7, 81.1, 77.2, 41.3, 39.9, 38.9, 36.7, 2.6, 2.2; LRMS (CI) *m/z* relative intensity 419 (*M*+1, 100), 329 (25).