

Copper- and Manganese-Catalyzed Homocoupling of Organostannanes in the Presence of Iodine

Suk-Ku Kang,* Tae-Gon Baik, Xiang Hua Jiao, and Yong-Tack Lee

Department of Chemistry, Sungkyunkwan University, Natural Science Campus, Suwon 440-746, Korea

Received 21 November 1998; accepted 22 January 1999


Abstract

The copper- and manganese-catalyzed homocoupling of aryl-, alkenyl-, and alkynylstannanes to afford biaryls, 1,3-dienes, and 1,3-diynes were achieved in the presence of iodine at 100 °C in good yields.

© 1999 Elsevier Science Ltd. All rights reserved.

Keywords: biaryls; coupling reactions; tin and compounds; copper and compounds; manganese and compounds

The oxidative homocoupling of aryl, alkenyl-, and alkynyl-substituted stannanes is one of the useful synthetic methods to form symmetrical biaryls, 1,3-dienes, and 1,3-diynes. Thus, the palladium-catalyzed homocoupling of organostannanes was reported employing *t*-BOOH [1], benzoquinone [2], HMPA [3], dichloroethane [4], diiodoethene [5], chloroacetone [6], acrylate dibromide [7], and air [8,9] as oxidant. Efficient homocoupling of stoichiometric amounts of copper salts have been already reported [10]. However, copper-catalyzed reactions have not been realized, in spite of their great potentials. Here we report that copper(II) chloride and manganese bromide efficiently catalyze the homocoupling of aryl-, alkenyl-, and alkynylstannanes by the use of iodine as an oxidant.



Scheme 1

To find suitable reaction conditions, we have examined the homocoupling of 1-phenylalkynylstannanes **1g**. Using CuI or CuCl (10 mol %) as catalyst in DMF at 60 °C or 100 °C the yield became lower (20–30 %) even in the presence of CsF or KF as additive. We found that by slow addition of I₂ (0.5 equiv) for 4 h the yield improved to more than >95% after isolation. As suitable catalysts, CuI, CuCl, CuCl₂, and CuF₂·2H₂O tested, CuCl₂ was the best of

choice. At 65 °C for 4 h we could get the product a rather lower (45%) yield.

However at 100 °C almost quantitative yield of the dimerized product was obtained. In the case of MnBr₂ (10 mol %) as catalyst, of the solvent THF, dioxane, DMF, and NMP, NMP was the most suitable. In considering the role of I₂ in these catalytic reactions, it is presumed that iodine might act as an oxidant or a scavenger for organostannanes. The results are summarized in Table 1 [11].

In conclusion, the catalytic oxidative homocoupling of organostannanes was accomplished by CuCl₂ or MnBr₂ adding slowly iodine as an oxidant.

Table 1. Copper- and Manganese-Catalyzed Homocoupling of Organostannanes in the Presence of Iodine

Entry	Substrate	Reaction Conditions ^a	Time	Product	Yield (%) ^b
1	1a	A	4	2a	93
		B	5		70
2	1b	A	4	2b	85
3	1c	A	4	2c	80
		B	5		80
4	1d	A	4	2d	72
		B	5		83
5	1e	A	4	2e	74
		B	5		85
6	1f	A	4	2f	68
		B	5		90
7	1g	A	4	2g	95
		B	5		81

^a A: CuCl₂ (10 mol %), I₂ (0.5 equiv), DMF, 100 °C. B: MnBr₂ (10 mol %), I₂ (0.5 equiv), NMP, 100 °C.

^b The isolated yields.

Acknowledgment. Generous financial support by KOSEF(97-0501-02-01-3) and Korea Research Foundation (BSRI-98-3420) is gratefully acknowledged.

Reference and Notes

- [1] Kanemoto S, Matsubara S, Oshima K, Utimoto K, Nozaki H. *Chem. Lett.* 1987;5-6.
- [2] Liebeskind LS, Riesinger SW. *Tetrahedron Lett.* 1991;32:5681-5682.
- [3] Tolstikov GA, Miftakhov MS, Danilova NA, LV Ya, Spirikhin LV. *Synthesis* 1989;633-634.
- [4] Kang S-K, Namkoong E-Y, Yamaguchi T. *Synth. Commun.* 1996;27:641-646.
- [5] Wright ME, Porch MJ, Buckley C, Cochran BB. *J. Am. Chem. Soc.* 1997;119:8393-8394.
- [6] Tamao K, Ohno S, Yamaguchi S. *J. Chem. Soc., Chem. Commun.* 1996;1873-1874.
- [7] Yamaguchi S, Ohno S, Tamao K. *Synlett* 1997;1199-1201.
- [8] Shirakawa E, Murata Y, Nakao Y, Hiyama T. *Synlett* 1997;1143-1144.
- [9] Alcaraz L, Taylor RJK. *Synlett* 1997;791-792.
- [10] (a) Ghosal S, Luke GP, Kyler KS. *J. Org. Chem.* 1987;52:4296-4298. (b) Beddoes RL, Cheeseright T, Wang J, Quayle P. *Tetrahedron Lett.* 1995;36:283-286. (c) Piers E, McEachern EJ, Romero MA. *Tetrahedron Lett.* 1996;37:1173-1176. (d) Piers E, Romero MA. *J. Am. Chem. Soc.* 1996;118:1215-1216.
- [11] The typical procedure is as follows. **Method A:** To a stirred solution of CuCl₂ (17.2 mg, 10 mol %) and DMF (5 mL) was added **1g** (500 mg, 1.28 mmol). The reaction mixture was heated to reflux at 100 °C and then I₂ (162 mg, 0.50 equiv) in DMF (5 mL) was added *via* syringe pump for 4 h. When the addition was completed checked by tlc or gc, the reaction mixture was cooled to room temperature and quenched with saturated KF solution. The reaction mixture was extracted with ether and the organic layer was dried over anhydrous MgSO₄ and evaporated *in vacuo*. The crude product was separated by SiO₂ column chromatography (hexanes, R_f = 0.38) to afford 1,4-diphenylbutadiyne (**2g**) (123 mg, 95%). **Method B:** To a stirred solution of MnBr₂ (29 mg, 10 mol %) and NMP (5 mL) was added **1a** (500 mg, 1.36 mmol). The reaction mixture was heated to reflux at 100 °C and then I₂ (173 mg, 0.50 equiv) in NMP (5 mL) was added *via* syringe pump for 5 h. When the addition was completed checked by tlc or gc, the reaction mixture was cooled to room temperature and quenched with saturated KF solution. The reaction mixture was extracted with ether and the organic layer was dried over anhydrous MgSO₄ and evaporated *in vacuo*. The crude product was separated by SiO₂ column chromatography (hexanes, R_f = 0.50) to afford biphenyl (**2a**) (74 mg, 70%).