

Influence of the Chiral Auxiliary on the Stereoselectivity of the $S_{RN}1$ C-Alkylation of 2-Nitropropionate Anions


Robert Nouguier*, Valérie Béraud, Patrice Vanelle, Michel P. Crozet

Laboratoire de Chimie Moléculaire Organique - UMR 6517 Universités d'Aix-Marseille 1 et 3, Avenue Normandie-Niemen
13397 Marseille Cedex 20 - France

Received 6 April 1999; accepted 4 May 1999

Abstract : A series of carbohydrate-derived 2-nitropropionate anions was reacted with *p*-nitrobenzyl chloride under $S_{RN}1$ reactions conditions, the diastereomeric ratios were moderate, but increased when the carbohydrate moiety was replaced by the more bulky 8-phenylmenthol. © 1999 Elsevier Science Ltd. All rights reserved.

We have recently described¹ an efficient method for the synthesis of C^{α} -unsymmetrically disubstituted nitroesters by electron transfer C-alkylation of ethyl 2-nitropropionate anion. To demonstrate the utility of the method, the nitroester **II** obtained from $S_{RN}1$ reaction of **I** with *p*-nitrobenzyl chloride, was transformed into 2-(±)-amino-3-(4-aminophenyl)-2-methylpropionic acid **IV**. The synthetic potential of this reaction is now extended even further in the synthesis of optically pure aminoacids.



To this goal, we first have screened a series of easily accessible chiral auxiliaries **1a-e** derived from carbohydrates (**1a** from D-arabinose², **1b-c** from D-fructose^{3,4}, **1d-e** from D-glucose⁵). The bromoesters **2** were obtained as a mixture of diastereomers by reaction of **1** with racemic 2-bromopropanoic bromide. By the Kornblum procedure⁶ the nitro esters **3** were obtained and then submitted to the $S_{RN}1$ reaction with *p*-nitrobenzyl chloride. The C-alkylation products **4a-f** were isolated in good to moderate yields as a mixture of diastereomers. The diastereomeric ratios (*dr*) were determined by ¹H NMR. As no asymmetric induction occurred with **3a**, we decided to increase the bulkiness at the anomeric position of the carbohydrate. With the spiro sugars **1b-c** easily accessible from cheap D-fructose the *dr* was increased to 60:40 and the diastereomers of the alkylated products were easily separable. To try to understand the role of the nature of the substituents adjacent to the carbon bearing the nitropropionate, we synthesized the two isomeric 2-*O* and 3-*O*-benzylethers of the methyl 4,6-*O*-benzylidene- α -D-glucopyranoside by the stannylene method⁵ but the *dr* was always 60:40.

As 8-phenylmenthol has been successfully used as chiral auxiliary for radical⁷ and ionic⁸ alkylations of malonic-type derivatives, we anticipate that the anion of the nitroester **3f** will react with the *p*-nitrobenzyl radical with high diastereoselectivity. At room temperature the *dr* was 80:20, at lower temperatures either the reaction was inhibited (-40 °C) or $S_{N}2$ *O*-alkylation became preponderant (0 °C).

Therefore, based on these findings, the design of new chiral auxiliaries was envisaged to reach high levels of induction. The camphor-derived alcohol **1g** was synthesized, nevertheless it has been impossible to prepare the corresponding bromoester **2g**.

$$\text{R-OH} \xrightarrow{i} \text{R-O}_2\text{C}-\underset{\text{H}}{\overset{\text{CH}_3}{\text{C}}}-\text{Br} \xrightarrow{j} \text{R-O}_2\text{C}-\underset{\text{H}}{\overset{\text{CH}_3}{\text{C}}}-\text{NO}_2 \xrightarrow{k} \text{R-O}_2\text{C}-\underset{\text{CH}_2\text{C}_6\text{H}_4\text{pNO}_2}{\overset{\text{CH}_3}{\text{C}}}-\text{NO}_2$$

1a-g **2a-f** **3a-f** **4a-f**

entry	R=	(i) yield	(ii) yield	(iii) yield - diastereomeric ratio
a		82%	55%	87% - 50:50
b		97%	50%	50% - 60:40*
c		96%	70%	51% - 60:40*
d		58%	32%	84% - 60:40
e		85%	40%	86% - 62:38
f		97%	80%	22% - 80:20**
g		0%	-	-

i: $\text{CH}_3\text{CHBrCOBr}$, CH_2Cl_2 , Py, $-5^\circ\text{C} \rightarrow \text{RT}$, 2 h; *j*: NaNO_2 , phloroglucinol, DMF, RT, 2 h;

k: $\text{ClCH}_2\text{-C}_6\text{H}_4\text{-}p\text{-NO}_2$, NaH, DMF, 150 Watt sun lamp, RT, 1 h; see ref 1 for exp. details.

* Separated by semi-prep. HPLC on SiO_2 , 4b: mp 161°C and 113°C , 4c: mp 96°C and 88°C .

** The newly created asymmetric carbon is *S* in the major isomer, see ref 8 for a discussion.

In conclusion, we have tested the ability of six chiral alcohols for the asymmetric induction in the $\text{S}_{\text{RN}}1$ reaction of *p*-nitrobenzyl chloride and 2-nitropropionate anions. The *dr* was not very high, but in the case of **4b** and **4c** the diastereomers were easily separated. As there is only one example of asymmetric induction in the literature⁹ ($\text{S}_{\text{RN}}1$ reaction of a chiral amide enolate and the radical from 1-iodonaphthalene), the design of efficient other chiral alcohols with bulky substituents on the adjacent carbon (camphor-derived auxiliaries) is now in progress for $\text{S}_{\text{RN}}1$ synthesis of enantiomerically pure disubstituted aminoacids.

References

- Béraud, V.; Perfetti, P.; Pfister, C.; Kaafarani, M.; Vanelle, P.; Crozet, M. P. *Tetrahedron* **1998**, *54*, 4923-4934.
- Paquette, L. A.; Bailey, S. *J. Org. Chem.* **1995**, *60*, 7849-7856.
- Nouguier, R.; Mignon, V.; Gras, J.-L. *J. Org. Chem.* **1999**, *64*, 1412-14.
- Mio, S.; Kumagawa, Y.; Sugai, S. *Tetrahedron* **1991**, *47*, 2133-2144.
- Jenkins, D.; Dubreuil, D.; Potter, B. *J. Chem. Soc. Perkin Trans. 1* **1996**, 1365-1372.
- Kornblum, N.; Blackwood, R.; Powers, J. *J. Am. Chem. Soc.* **1957**, *79*, 2507-2515.
- Hamon, D.; Massy-Westropp, R.; Razzino, P. *Tetrahedron* **1995**, *51*, 4183-4194.
- Ihara, M.; Takahashi, M.; Tanigushi, N.; Yasui, K.; Niitsuma, H.; Fukumoto, K. *J. Chem. Soc. Perkin Trans. 1* **1991**, 525-535.
- Lotz, G.; Palacios, S.; Rossi, R. *Tetrahedron Lett.* **1994**, *35*, 7711-7714.