

[1,5]SIGMATROPIC SHIFT OF HYDROGEN IN AMINATION
OF 3-PYRROLIDINO-1,2,4-TRIAZINE 4-OXIDE[†]

Oleg N. Chupakhin, Valery N. Kozhevnikov, Dmitry N. Kozhevnikov, Vladimir L. Rusinov*

Urals State Technical University, 620002, Ekaterinburg, Russian Federation

Fax: +7 3432 740458, E-mail: mike@htf.ustu.ru

Received 16 April 1999; accepted 22 June 1999

Abstract: The reaction of 3-pyrrolidino-1,2,4-triazine 4-oxide with ammonia leads to the product of *tele*-substitution of pyrrolidine - 5-amino-1,2,4-triazine 4-oxide. Sigmatropic shift of hydrogen postulated for such reactions has been proved by isolation of key intermediates.

© 1999 Published by Elsevier Science Ltd. All rights reserved.


Nucleophilic attack on an azine ring is often accompanied by cleavage of the C-C bond resulting in open-chain or ring-transformation products.¹ This can be illustrated by some classical examples, such as the formation of glutamic aldehyde derivatives in the reactions of pyridinium or quinolinium cations with a variety of nucleophiles,² nucleophilic substitution reactions in pyrimidines according to the ANRORC (Addition of Nucleophile, Ring Opening, Ring Closing) mechanism,³ or the formation of 3-aminopyridazine as a result of the ring transformation of 3-chloro-6-phenyl-1,2,4-triazine by action of the phenylacetonitrile anion.⁴ Often, ring-opening is observed in aromatic N-oxides, as exemplified by hydrolysis of 1,2,4-triazine-4-oxides,⁵ by reactions of pyridine N-oxides⁶ and 1,2,4-triazine-4-oxides⁷ with amines, as well as by the reactions of pyrimidine N-oxides⁸ and 1,2,4-triazine-4-oxides⁵ with active methylene compounds, etc.

Continuing our studies on nucleophilic substitution reactions in 1,2,4-triazine 4-oxides, we have found that heating 3-pyrrolidino-6-phenyl-1,2,4-triazine-4-oxide **1** with ethanolic ammonia leads to 5-amino-6-phenyl-1,2,4-triazine-4-oxide **2** in 70% yield. The melting point and spectral data for the oxide **2** proved to be identical to those for the compound obtained by amination of 6-phenyl-1,2,4-triazine-4-oxide in liquid ammonia in the presence of potassium permanganate.⁹ Obviously in this case, we deal with *tele*-substitution of the pyrrolidino group, as an example of the so-called odd *tele*-substitution (an odd number of atoms lie between the position of nucleophilic attack and that from which a nucleofuge departs). In accordance with the mechanism postulated for such reactions,¹⁰ the [1,5]-sigmatropic shift of hydrogen is a crucial step in conversion of the adduct **A** resulting from the addition of ammonia at C⁵ of the 1,2,4-triazine-4-oxide **1** into the 3,4-dihydro intermediate **B**. This is aromatized into the *tele*-substitution product **2** by elimination of pyrrolidine.

In this paper we present new experimental evidence for the generally accepted mechanism mentioned above by isolation of key intermediates. Indeed, the reaction of 1,2,4-triazine-4-oxide **1** with ammonia, dimethylamine or cycloalkylamines has unexpectedly been found to lead under the mild conditions (room temperature) to 6-pyrrolidino-2-amino-1-hydroxy-3-phenyl-1,4,5-triazahexatrienes **3a-d**.

[†] Dedicated to Professor H.C. van der Plas on the occasion of his 70-th birthday

The structures of products **3a-d** are in good agreement with NMR spectral data, and for 1-hydroxy-2,6-dipyrrolidino-3-phenyl-1,4,5-triazahexatriene **3b** was confirmed by X-ray diffraction. In their ^1H NMR spectra chemical shifts of the phenyl protons and H^6 (8.2-8.3 ppm) are identical to those for 1-hydroxy-6-dialkylamino-3-phenyl-1,4,5-triazahexatrienes.⁷



Refluxing triazahexatriene **3a** in ethanol leads to its full conversion into 5-amino-1,2,4-triazine 4-oxide **2**. Taking into account that open-chain 1-hydroxy-1,4,5-triazahexatrienes **3** exist in equilibrium with the cyclic 3,4-dihydro-4-hydroxy-1,2,4-triazines¹¹ one can see that compounds **3a-d** are just open-chain forms of σ^{H} -adducts **B**. The location of the substituents and protons in the compounds **3** confirms the conversion of **A** into **B** through the [1,5]-sigmatropic shift of hydrogen.

Thus, a reversible opening of the 1,2,4-triazine ring in the reaction of 3-pyrrolidino-1,2,4-triazine 4-oxide with amines removes the intermediates **B** from the reaction. It allows a study in detail of their structure. The conclusion resulting from this structure is undoubtedly that the odd *tele*-substitution proceeds through the [1,5]-sigmatropic shift of hydrogen.

References

- H.C. van der Plas, *Ring Transformations of Heterocycles*. N.Y.: Acad. Press, 1973
- I.S.Poddubny, *Khim. Geterotsycl. Soed. (Chem. Heterocycl. Comp.-Rus)*, **1996**, 774-815
- H.C. van der Plas, *Acc. Chem. Res.*, **1978**, 11, 462-468
- A. Rykowski, and E. Wolinska, *Tetrahedron Lett.*, **1996**, 37, 5795-5796
- D.N. Kozhevnikov, V.L. Rusinov, O.N. Chupakhin, *Russian Chem. Reviews*, **1998**, 67, 633-647
- J. Schnekenburger, H. Riedel, *Arch. Pharmazie*, **1983**, 316, 27-33
- Chupakhin O.N., Kozhevnikov V.N., Kozhevnikov D.N., Rusinov V.L. *Zh. Org. Khim. (Russ. Journal of Organic Chemistry)*. **1998**, 34, 418-422
- H. Yamanaka, T. Sakamayo, and S. Nitauma, *Heterocycles*, **1990**, 31, 923-928
- A.Rykowski, H.C. van der Plas, *Synthesis*, **1985**, 884-885
- H.C. van der Plas, M. Wozniak, A. van Veldhuizen, *Recl. Trav. Chim. (Pays-Bas)*, **1978**, 97, 130-136
- D.N. Kozhevnikov, V.N.Kozhevnikov, V.L.Rusinov, O.N.Chupakhin, E.O.Sidorov, N.A.Kl'uev, *Zh. Org. Khim. (Russ. Journal of Organic Chemistry)*. **1998**, 34, 423-428