



Pergamon

Tetrahedron Letters 40 (1999) 6603–6607

TETRAHEDRON
LETTERS

Rhodium(II)-catalyzed reactions of 3-diazo-2,4-chromenediones. First one-step synthesis of pterophyllin 2

Yong Rok Lee,* Jung Yup Suk and Byung So Kim

School of Chemical Engineering and Technology, College of Engineering, Yeungnam University, Kyongsan 712-749, Korea


Received 6 May 1999; revised 28 June 1999; accepted 2 July 1999

Abstract

Rhodium(II)-catalyzed reactions of 3-diazo-2,4-chromenediones 1–3 with a variety of substrates are described. Reactions of 1 with nitriles, isocyanates, and ketones give the corresponding 5-membered heterocycles in moderate yields, whereas those of 1 with acid chlorides afford α -chloroenones in moderate yields. Reactions of 1 with electron-rich and -deficient olefins such as vinyl ethers and allyl iodide give dihydrofurocoumarins as a single compound, whereas those of 3 with vinyl esters afford either angular dihydrofurocoumarin and a linear product or a rearranged product as an isomer. Furthermore, this novel method has been applied to the synthesis of the natural pterophyllin 2 isolated from *Ekebergia pterophylla*. © 1999 Elsevier Science Ltd. All rights reserved.


The rhodium-catalyzed decomposition of diazocarbonyl compounds has become an important method in organic and natural product synthesis.¹ The rhodium(II)-catalyzed reactions of acyclic and cyclic diazodicarbonyl compounds with several substrates such as olefins, nitriles, isocyanates, carbon disulfides, furans, benzofurans, thiophenes, and pyrroles have been extensively studied by many groups.² The reactions include C-H and X-H insertion,³ cyclopropanation,⁴ ylide formation,⁵ and cycloaddition.⁶ However, little study has been made of the reactions of 3-diazo-2,4-chromenedione 1–3 to substrates. We have been interested in reactions of 2-diazo-1,3-dicarbonyl compounds with several substrates.⁷ As an extension of our work, we have investigated the rhodium-catalyzed carbenoid reactions of 3-diazo-2,4-chromenediones with substrates to synthesize the skeletons of naturally-occurring coumarins and dihydrofurocoumarins.⁸ They are reported to show various biological activities such as anticoagulant, insecticide, anthelmintic, hypnotic, antifungal, phytoalexin, and provide HIV protease inhibition.⁹ This wide range of biological properties has stimulated the development of the new synthetic methods for the construction of coumarins and dihydrofurocoumarins.

* Corresponding author.



- 1 $R_1=R_2=H$
 2 $R_1=CH_3, R_2=H$
 3 $R_1=H, R_2=CH_3$


3-Diazo-2,4-chromenediones 1–3 were prepared by the diazotransfer reaction of the corresponding 4-hydroxycoumarins with mesyl azide according to Taber's method.¹⁰ 3-Diazo-2,4-chromenediones 1–3 are fairly stable and can be stored in a refrigerator for a long time without any decomposition. In order to check the reactivity of 1 as carbenoid reactions, several nitriles were first used. When acetonitrile was heated at 60°C for 5 h as a solvent and a reactant in the presence of 1 mol% of $Rh_2(OAc)_4$, oxazole 4a was isolated in a 50% yield (Scheme 1). Under the same conditions, chloroacetonitrile and phenylacetonitrile also afforded oxazoles 4b (95%) and 4c (73%) in increased yields. When ethyl isocyanate and propyl isocyanate were used, oxazolones 5a and 5b were produced in 30 and 54% yields, respectively, as a single compound. Reactions of 1 with 2-butanone and 3-methyl-2-butanone, 1,3-dioxoles 6a and 6b were obtained in 63 and 35% yields, respectively. In particular, there is no direct precedent for Rh(II)-catalyzed reaction of diazodicarbonyl compounds with ketones, so it is noteworthy that reactions of 1 with ketones give the 1,3-dioxoles in moderate yields. In related work, Alonso has reported that the copper(II)-catalyzed reaction of acyclic methyl 2-diazo-3-oxobutyrates with ketones gave 1,3-dioxole adducts in moderate yields.¹¹ However, reaction of 1 with other carbonyl compounds such as acetyl chloride and propionyl chloride, α -chloroenones 7a and 7b were produced in 35 and 63% yields, respectively, without a trace of expected 1,3-dioxole adducts. Support for the structural assignment comes from their spectroscopic analysis. The α -chloroenone 7a is identified by the 1H NMR peak of the methyl group of the vinyl acetate as a singlet at δ 2.49, and by the two IR carbonyl absorptions of the lactone at 1730 cm^{-1} and the vinyl ester at 1784 cm^{-1} .



Scheme 1.

To synthesize the skeletons of naturally-occurring coumarins and dihydrofurocoumarins, additional reactions with several olefins were investigated. Reaction of 1 with an excess of cyclohexene used as


a solvent and a reagent, in the presence of 1 mol% of $\text{Rh}_2(\text{OAc})_4$ at 60°C for 5 h gave the unexpected rearranged product **8** in a 51% yield (Scheme 2). The two olefinic protons of **8** were observed at δ 6.29 and δ 6.04. When allyl iodide with electron-deficient olefin was used, the dihydrofurocoumarin **9** was obtained in 78% yield as a single compound. This result is also in clear contrast to that of Doyle, who reported that the rhodium(II)-catalyzed reactions of acyclic ethyl diazoacetate with allyl iodide afforded the [2,3]-rearranged product without forming the cycloaddition adduct as a single compound.¹² Another approach for dihydrofurocoumarin formation was obtained by using vinyl ethers. On heating neat ethyl vinyl ether and isobutyl vinyl ether in the presence of 1 mol% of $\text{Rh}_2(\text{OAc})_4$ for 3 h, dihydrofurocoumarins **10a** and **10b** were produced in 69 and 64% yields, respectively. These results indicate that reactions with more electron-rich and electron-deficient olefins than simple alkenes such as cyclohexene provide an efficient route for synthesizing the dihydrofurocoumarin skeletons.



Scheme 2.


Next, reactions with vinyl acetates were also examined. Reaction of **3** with vinyl acetate at 60°C for 5 h afforded the two regioisomeric cycloadducts **11** and **12** in 16 and 25% yields, respectively (Scheme 3). The two isomers were easily assigned from spectroscopic data. The ^1H NMR spectrum of angular dihydrofurocoumarin **11** showed methine peaks at δ 7.00 as a doublet associated with the dihydrofuran ring, whereas linear adduct **12** showed peaks at δ 6.87 as a doublet. The clear assignment was based on comparison of IR carbonyl absorption. In angular isomer **11**, there are two $\text{C}=\text{O}$ peaks stretching at 1753 and 1711 cm^{-1} corresponding to the two ester groups, while in linear isomer **12** the two $\text{C}=\text{O}$ stretches are shown at 1660 cm^{-1} for an enone and at 1765 cm^{-1} for an ester group. On the other hand, reaction of **3** with isopropenyl acetate afforded angular dihydrofurocoumarin **13** (35%) and rearranged product **14** (32%) as a mixture. Surprisingly, in this case, the expected linear adduct was not detected. These mixtures were easily separated by column chromatography, and the assignment of their structures of **13** and **14** was made on the basis of spectral data. The methylene protons of **14** are observed as a singlet at δ 3.55, and the two methyl protons of the ester and ketone are shown at δ 2.43 and δ 2.41, also as singlets. Further support for the structural assignment of **14** is accomplished from its IR spectrum, which clearly shows the expected three $\text{C}=\text{O}$ absorptions at 1776 , 1730 , and 1709 cm^{-1} due to two esters and a ketone.

Finally, in order to extend the utility of these reactions, one-step synthesis of a furocoumarin natural product was attempted. Pterophyllin 2 (**15**) has been isolated from a small evergreen tree of *Ekebergia pterophylla*, found in South Africa.¹³ The structure of pterophyllin 2 was established by spectroscopic analysis, but no synthetic methods are known. Reaction of **2** with 2-methyl-1-buten-3-yne (15-fold



Scheme 3.

excess) under 1 mol% of rhodium catalysis at reflux for 3 h afforded pterophyllin **2** (33%) together with linear isomer **16** (35%) as a mixture (Scheme 4). These products were easily separated by column chromatography, and the spectroscopic properties of our synthetic pterophyllin **2** agree well with those reported in the literature.¹³



Scheme 4.

In conclusion, the rhodium(II)-catalyzed reactions of 3-diazo-2,4-chromenedione **1–3** with several substrates is described. This method provides a novel entry into the naturally-occurring biologically active coumarin and dihydrofurocoumarin skeletons, and demonstrates an interesting application of one-step synthesis of the natural product, pterophyllin **2**. Further mechanistic studies and applications of these carbenoid reactions are in progress in our laboratory.

Acknowledgements

This work was supported by the Korea Science and Engineering Foundation (981-0303-016-2).

References

- (a) Doyle, M. P.; Mckerverey, M. A.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds: from Cyclopropanes to Ylides*; Wiley, 1997. (b) Ye, T.; Mckerverey, M. A. *Chem. Rev.* **1994**, *94*, 1091. (c) Padwa, A.; Weingarten, M. D. *Chem. Rev.* **1996**, *96*, 223. (d) Miller, D. J.; Moody, C. J. *Tetrahedron* **1995**, *51*, 10811. (e) Padwa, A.; Krumpe, K. E. *Tetrahedron* **1992**, *48*, 5385. (f) Doyle, M. P. *Acc. Chem. Res.* **1986**, *19*, 348.
- (a) Connell, R. D.; Tebbe, M.; Helquist, P.; Akermark, B. *Tetrahedron Lett.* **1986**, *27*, 5559. (b) Yoo, S.-K. *Tetrahedron Lett.* **1992**, *33*, 2159. (c) Shi, G.; Xu, Y. *J. Chem. Soc., Chem. Commun.* **1989**, 607. (d) Pirrung, M. C.; Zhang, J.; McPhail, A. T. *J. Org. Chem.* **1991**, *56*, 6269.

3. (a) Taber, D. F.; Ruckle Jr., R. E. *J. Am. Chem. Soc.* **1986**, *108*, 7686. (b) Taber, D. F.; Raman, K. *J. Am. Chem. Soc.* **1983**, *105*, 5935. (c) Heslin, J. C.; Moody, C. J.; Slawin, A. M. Z.; Williams, D. J. *Tetrahedron Lett.* **1986**, *27*, 1403.
4. (a) Doyle, M. P.; Bagheri, V.; Wandless, T. J.; Harn, N. K.; Brinker, D. A.; Eagle, C. T.; Loh, K.-L. *J. Am. Chem. Soc.* **1990**, *112*, 1906. (b) Petiniot, N.; Anciaux, A. J.; Noels, A. F.; Hubert, A. J.; Teyssie, P. *Tetrahedron Lett.* **1978**, 1239.
5. (a) Doyle, M. P.; Bagheri, V.; Harn, N. K. *Tetrahedron Lett.* **1988**, *29*, 5119. (b) Kirmse, W.; Kund, K. *J. Am. Chem. Soc.* **1989**, *111*, 1465. (c) Padwa, A. *Acc. Chem. Res.* **1991**, *24*, 22.
6. (a) Davies, H. M. L.; Romines, K. R. *Tetrahedron* **1988**, *44*, 3343. (b) Wenkert, E.; Ananthanarayan, T. P.; Ferreira, V. F.; Hoffman, M. G.; Kim, H. *J. Org. Chem.* **1990**, *55*, 4975.
7. (a) Lee, Y. R.; Suk, J. Y. *Heterocycles* **1998**, *48*, 875. (b) Lee, Y. R.; Suk, J. Y. *Chem. Commun.* **1998**, 2621. (c) Lee, Y. R. *Tetrahedron* **1995**, *51*, 3087. (d) Lee, Y. R.; Morehead Jr., A. *Tetrahedron* **1995**, *51*, 4909.
8. (a) Miski, M.; Jakupovic, J. *Phytochemistry* **1990**, *29*, 1995. (b) Schuster, N.; Christiansen, C.; Jakupovic, J.; Mungai, M. *Phytochemistry* **1993**, *34*, 1179. (c) Bittner, M.; Jakupovic, J.; Grenz, M.; Silva, M. *Phytochemistry* **1988**, *27*, 3263. (d) Rustalyan, A.; Nazarians, L.; Bohlmann, F. *Phytochemistry* **1980**, *19*, 1254. (e) Zdero, C.; Bohlmann, F.; King, R. M.; Robinson, H. *Phytochemistry* **1986**, *25*, 509.
9. (a) Feuer, G. In *Progress in Medicinal Chemistry*; Ellis, G. P.; West, G. B., Eds; North-Holland Publishing: New York, 1974. (b) Deana, A. A. *J. Med. Chem.* **1983**, *26*, 580. (c) Wenkert, E.; Buckwalter, B. L. *J. Am. Chem. Soc.* **1972**, *94*, 4367.
10. Taber, D. F.; Ruckle Jr., R. E.; Hennessy, M. *J. Org. Chem.* **1986**, *51*, 4077.
11. (a) Alonso, M. E.; Chitty, A. W. *Tetrahedron Lett.* **1981**, *22*, 4181. (b) Alonso, M. E.; Garcia, M. C.; Chitty, A. W. *J. Org. Chem.* **1985**, *50*, 3445.
12. Doyle, M. P.; Tamblyn, W. H.; Bagheri, V. *J. Org. Chem.* **1981**, *46*, 5094.
13. Mulholland, D. A.; Iourine, S. E.; Taylor, D. A. H.; Dean, F. M. *Phytochemistry* **1998**, *47*, 1641.