



## Corrigendum









Corrigendum to "Direct displacement of -OH by nucleophiles  
in hydroxymethylimidazoles"[Tetrahedron Letters 41 (2000) 2777]<sup>†</sup>

John E. Macor,\* Lyndon A. M. Cornelius and Jacques Y. Roberge

*Bristol-Myers Squibb, Pharmaceutical Research Institute, Discovery Chemistry, PO Box 5400,  
Princeton, NJ 08543-5400, USA*

Table 1 contained ambiguous 'Nucleophiles' for entries **D**, **E**, **F** and **G**. All other entries were correct. The corrected portion of the Table is shown below.

Table 1  
Synthesis of functionalized imidazoles

Entry	Imidazole	Nucleophile	Product	Yield (%)
<b>D</b>		NH(CH <sub>3</sub> ) <sub>2</sub>		<b>99</b>
<b>E</b>		NH(CH <sub>3</sub> ) <sub>2</sub>		<b>95</b>
<b>F</b>		NH(CH <sub>3</sub> ) <sub>2</sub>		<b>81</b>
<b>G</b>		CH <sub>3</sub> ONHCH <sub>3</sub>		<b>100</b>

\* Corresponding author. E-mail: john.macor@bms.com

† PII of original article: S0040-4039(00)00262-8