



Corrigendum

Corrigendum to "An efficient one-pot synthesis of (+)-deoxypyridinoline" [Tetrahedron Letters 40 (1999) 8993] and "Bone collagen cross-links: an efficient one-pot synthesis of (+)-pyridinoline and (+)-deoxypyridinoline" [Tetrahedron *Asymmetry* 11 (2000) 2289][†]

Maciej Adamczyk,* Donald D. Johnson and Rajarathnam E. Reddy

Department of Chemistry (9NM, Bldg AP20-2), Diagnostics Division, Abbott Laboratories, 100 Abbott Park Road, Abbott Park, IL 60064-6016, USA

We wish to acknowledge that a one-pot protocol first described by Allevi, P.; Longo, A. and Anastasia, M. ("Total synthesis of deoxypyridinoline, a biochemical marker of collagen turnover", *Chem. Commun.* **1999**, 559) for the preparation of 3-hydroxypyridinium compound **1**, was adapted for one-pot conversion of (S)-(-)-tert-butyl-6-bromo-[bis-(2-tert-butoxycar-bonyl)amino]-5-oxohexanoate to (+)-deoxypyridinoline (2) (*Tetrahedron Lett.* **1999**, 40, 8993), and (+)-pyridinoline (3) (*Tetrahedron: Asymmetry* **2000**, 11, 2289).

(+)-Pyridinoline (X=OH, 3)

(+)-Deoxypyridinoline (X=H, 2)

0040-4039/00/\$ - see front matter © 2000 Elsevier Science Ltd. All rights reserved. PII: S0040-4039(00)01533-1

^{*} Corresponding author. E-mail: maciej.adamczyk@abbott.com

[†] PII of original article: S0040-4039(99)01936-X