



Available online at www.sciencedirect.com



Tetrahedron
Letters

Tetrahedron Letters 47 (2006) 1487

Corrigendum

Corrigendum to “Synthesis of chiral vicinal C_2 symmetric and unsymmetric bis(sulfonamide) ligands based on *trans*-1,2-cyclohexanediamine by aminolysis of *N*-tosylaziridines”

[Tetrahedron Lett. 46 (2005) 7935]

Alakesh Bisai, B. A. Bhanu Prasad and Vinod K. Singh*

Department of Chemistry, Indian Institute of Technology, Kanpur 208 016, India

Available online 27 December 2005

We apologize for the oversight of omitting the following references on the cleavage of *N*-alkylaziridines towards the synthesis of *trans*-1,2-cyclohexanediamine derivatives in the above paper.

References

1. For the synthesis of *trans*-1,2-cyclohexanediamine derivatives through the ring cleavage of cyclohexyl-*N*-alkylaziridine, see: (a) de Parrodi, C. A.; Moreno, G. E.; Quintero, L.; Juaristi, E. *Tetrahedron: Asymmetry* **1998**, *9*, 2093–2099; (b) de Parrodi, C. A.; Vazquez, V.; Quintero, L.; Juaristi, E. *Synth. Commun.* **2001**, *31*, 3295–3302.
2. For the use of C_2 -symmetric *trans*-1,2-cyclohexanediamine based ligands in enantioselective reactions, see: (a) Mastranzo, V. M.; Quintero, L.; de Parrodi, C. A.; Juaristi, E.; Walsh, P. J. *Tetrahedron* **2004**, *60*, 1781–1789; (b) de Parrodi, C. A.; Walsh, P. J. *Synlett* **2004**, 2417–2420.

DOI of original article: 10.1016/j.tetlet.2005.09.089

* Corresponding author. Tel.: +91 512 2597291; fax: +91 512 2597436; e-mail: vinodks@iitk.ac.in