

Efficient synthesis of chiral phenethylamines: preparation, asymmetric hydrogenation, and mild deprotection of ene-trifluoroacetamides

Shawn P. Allwein,^{a,*} J. Christopher McWilliams,^{b,*} Elizabeth A. Secord,^a
Dale R. Mowrey,^a Todd D. Nelson^a and Michael H. Kress^a

^aMerck Research Laboratories, Department of Process Chemistry, 466 Devon Park Drive, Wayne, PA 19087, USA

^bMerck Research Laboratories, Department of Process Chemistry, 126 East Lincoln Avenue, Rahway, NJ 07065, USA

Received 7 June 2006; revised 24 June 2006; accepted 26 June 2006

Abstract—A mild and efficient route to enantioenriched aryl alkyl amines from ketones has been developed. The first successful synthesis and asymmetric hydrogenation of ene-trifluoroamides from oximes gave highly enantioenriched trifluoroacetamides (94–98% ee). The corresponding phenethyl amides are liberated under mild conditions (K_2CO_3 , MeOH/H₂O). In addition, a new application of Josiphos ligands toward the asymmetric hydrogenation of both ene-acetamides and ene-trifluoroacetamides was discovered.
© 2006 Elsevier Ltd. All rights reserved.

Optically active amines are found in abundance within biologically active natural products and pharmaceutical agents.¹ They have also been established as a useful class of resolving agents for organic synthesis.² While numerous synthetic routes to enantioenriched amines have been developed, catalytic asymmetric hydrogenation of the corresponding enamides is arguably one of the most valuable methods due to the inherent efficiency of these atom-economical processes. Originally communicated by Kagan,³ asymmetric hydrogenation of enamides allows for convenient arrival to enantioenriched acetamides that can be converted to the amines by removal of the acetate functionality. Since that initial discovery, the overwhelming focus of subsequent research in this area has been on the design of new chiral ligands for asymmetric hydrogenations. Consequently, many catalytic systems have been discovered that can reduce a variety of enamides in high yield and with very high enantioselectivity.⁴ While the field of enamide asymmetric hydrogenation has matured, there has not been a corresponding application of these methods to complex synthetic targets. The reason for this discrepancy

may lie in the fact that little attention has been placed on the preparation and design of practical functionality for the enamide itself. With few exceptions,⁵ enamides used in these processes are in the form of acetamides derived from oximes and acetic anhydride. These enamides suffer from a significant limitation: the products of the asymmetric hydrogenation require harsh conditions to convert them to amines. It has been stated that acetamides are ‘worthless as protecting groups because the conditions required to remove them are harsh’.⁶ From our own experience, we have seen removal of such acetates required prolonged exposure to 6 N HCl in refluxing MeOH. Consequently, we sought to develop a more practical enamide that could take advantage of the highly efficient asymmetric hydrogenation methods currently available. Aware of the clear advantages of trifluoroacetate over acetate as a removable protecting group, we were hopeful that an extension of methodology could be made. A previous attempt to reduce an ene-trifluoroacetate with ruthenium catalysis was unsuccessful.⁵ In that communication, the formyl group was employed to achieve milder deprotection. However, this protocol still required strongly basic conditions at elevated temperatures. Herein, we report our successful approach to developing a mild and practical route to enantioenriched amines via ene-trifluoroamides.

The benchmark synthesis of enamides reported by Burk utilized an iron mediated reduction of α -arylalkyl-oximes

Keywords: Asymmetric; Hydrogenation; Amines; Amides; Homogeneous; Catalysis.

* Corresponding authors. Tel.: +1 610 883 5969 (S.P.A.); +1 908 656 0285 (J.C.M.); e-mail addresses: sallwein@cephalon.com; chrismcwilliams@msn.com

that are derived from the corresponding ketones.^{7,8} This approach is an extension of Barton's earlier work.⁹ To date, this reaction has only been applied to the synthesis of ene-acetamides.^{4,10} Our modification to Burk's original enamide synthesis has led to a series of TFA-protected enamides in high overall yield. By simply substituting Ac₂O and HOAc with TFAA and TFA, ene-trifluoroacetamides **3** were generated in high overall yield (Table 1, entries a–f). Acetophenones bearing electron donating groups (entry g) were found to be problematic. Under the acidic reaction conditions, the oximes underwent a Beckmann rearrangement to form the corresponding amide.¹¹

An alternative entry to ene-trifluoroacetamides is the Blaise reaction,¹² which occurs under non-acidic conditions, thereby providing a complimentary approach to the oxime approach.¹³ Two examples, **5** and **3g**, have been generated from the corresponding nitrile via methyl addition and in situ trapping (Table 2). This protocol increased the overall yield of **3g** by twofold.

The literature is replete with examples of ene-acetamide asymmetric hydrogenations.⁴ For most simple aromatic substrates, one can choose from a variety of chiral ligands to achieve the highest enantioselectivity for a given substitution pattern. Our objective was to determine and demonstrate that reductions of the ene-trifluoroacetamides would provide comparable enantioselectivities with known performance ligands in this field.¹⁴ Accordingly, we selected a well-established catalyst system for this purpose. Specifically, the MeBPE ligand originally described by Burk was employed in


the demonstrations. Under standard reaction conditions, the hydrogenation of the ene-trifluoroacetamides provided high yields and enantioselectivities in each case (Table 3). In comparison to the corresponding ene-acetamides cited in the literature,⁷ the enantioselectivities derived from the reduction of the ene-trifluoroacetamides were within $\pm 1.4\%$ ee.

Recent findings in our laboratory led us to evaluate a new application of the Josiphos ligands to the asymmetric hydrogenation of enamides.¹⁵ The Josiphos derivatives **I** and **II** were found to reduce the ene-acetamide **5** to the acetamide **7** in 95.0% ee and 96.0% ee, respectively (Fig. 1).

To our knowledge, this is the first reported example of an asymmetric hydrogenation of an enamide using these ligands. A solvent screen with these ligands yielded the corresponding trifluoroacetamides with high enantioselectivity (Table 4). While ligand **I** was highly selective in both alcohol and the haloaromatic solvent 1,2-DCB, ligand **II** appeared to perform well only in 1,2-DCB. Additionally, only ligand **II** in 1,2-DCB reduced **3f** to the corresponding trifluoroamide with high enantioselectivity. Thus, subtle changes in ene-trifluoroamide structure can prompt changes in optimal ligand and solvent conditions


While pleased with the asymmetric hydrogenation results, we recognized that their utility could only be recognized and implemented for the synthesis of complex molecules if mild deprotection of the trifluoroacetamide was achieved as expected. To this end, the enantio-enriched trifluoroacetamides were treated with 2 equiv of K₂CO₃ in methanol at room temperature (Table 5). These deprotection conditions resulted in good to excellent yields of chiral amines.

Table 1. Synthesis of ene-trifluoroacetamides




Entry	R ¹	R ²	2 (%)	3 (%)
a	H	H	99	90
b	Br	H	100	93
c	CO ₂ Me	H	100	90
d	Ph	H	96	94
e	CF ₃	H	100	96
f	Br	F	100	98
g	OMOM	H	95	26

Table 2. Enamide synthesis via Blaise reaction



R ¹	R ²	R ³	MeMX	Yield	Product
Br	F	CH ₃	MeMgBr	83	5
OMOM	H	CF ₃	MeLi–LiBr	51	3g

Table 3. Asymmetric hydrogenation of ene-trifluoroacetamide with (MeBPE)Rh(COD)BF₄



Substrate	R ¹	R ²	Conversion (yield) ^a	% ee of 6	% ee of CH ₃ ^b
3a	H	H	>99 (80) ^c	96.5	95.2
3b	Br	H	>99 (91)	94.4	95.8
3c	CO ₂ Me ^d	H	>99 (95)	96.2	NDA ^e
3d	Ph	H	97 (94)	96.0	NDA ^e
3e	CF ₃	H	>99 (98)	96.8	95.6
3f	Br	F	>99 (89)	98.3	NDA ^e
3g	OMOM	H	>99 (97)	97.6	NDA ^e

^a Conversion is HPLC area percent conversion at 210 nm. Isolated yield in parentheses.

^b Percent ee of the corresponding acetamide (with 0.2% catalyst) as reported in Ref. 4p.

^c Assay yield = 95%.

^d Reaction run in MeOH.

^e NDA = no data available.




Figure 1. Asymmetric hydrogenation of ene-acetamide **5** with Josiphos ligands **I** and **II**.

Table 4. Asymmetric hydrogenation of ene-trifluoroamides with Josiphos ligands **I** and **II**

Substrate	Ligand	Solvent	% ee
3e	I	THF	95.8
3e	I	IPA	97.1
3e	II	THF	56.7
3e	II	1,2-DCB	94.2
3f	I	THF	45.4
3f	I	IPA	77.8
3f	II	THF	41.0
3f	II	1,2-DCB	91.8


Table 5. Deprotection of trifluoroacetamides to chiral amines

Substrate	R ¹	R ²	Time (h)	Yield ^a
6a	H	H	48	100
6b	Br	H	48	100
6c	CO ₂ Me	H	24	80
6d	Ph	H	72	100
6e	CF ₃	H	48	100
6f	Br	F	48	100
6g	OMOM	H	48	100
6h	OAc	H	96	84 ^b

^a Assay yield by HPLC.

^b Yield for complete hydrolysis of amide and phenolic acetate.

Further orthogonal deprotection strategies utilizing the phenolic MOM-ether demonstrated the advantages of the trifluoroacetate protecting group. From **6g**, selective



Scheme 1. Selective deprotection of trifluoroacetamide **6g**.

hydrolysis of either the MOM ether or the trifluoroacetate occurs in excellent yield (Scheme 1).

In summary, we have developed a new and practical synthesis of enantioenriched aryl amines that merges a mild protecting group with highly efficient asymmetric hydrogenation technology. The trifluoroamide functionality allowed high enantioselectivity to be achieved as well as easy deprotection under extremely mild conditions. In addition, we have discovered two new Josiphos ligands that can reduce ene-acetamides and ene-trifluoroacetamides with high enantioselectivity. The high enantioselectivity and mild deprotection conditions should render this procedure applicable to a broad range of complex synthetic targets.

Acknowledgments

We thank Mirlinda Biba for developing chiral analysis and Peter Dormer and Lisa Dimichele for assistance in NMR data.

References and notes

- (a) Kleemann, A.; Engel, J. *Pharmaceutical Substances: Synthesis, Patents, Applications*, 4th ed.; Thieme: Stuttgart, New York, 2001; (b) Blaser, H. U.; Schmidt, E. *Asymmetric Catalysis on Industrial Scale: Challenges, Approaches and Solutions*; Wiley-VCH: New York, 2004.
- (a) Jacques, J.; Collet, A.; Wilen, S. H. *Enantiomers, Racemates, and Resolutions*; John Wiley and Sons: New York, 1981; (b) Sawayama, T.; Tsukamoto, M.; Sasagawa, T.; Naruto, S.; Matsumoto, J.; Uno, H. *Chem. Pharm. Bull.* **1989**, *37*, 1382; (c) Kawai, M.; Omuri, Y.; Yamamura, H.; Butsugan, Y. *Tetrahedron Asymmetry* **1992**, *3*, 1019.
- (a) Kagan, H. B.; Langlois, N.; Dang, T. P. *J. Organomet. Chem.* **1975**, *114*, 353; (b) Sinou, D.; Kagan, H. B. *J. Organomet. Chem.* **1976**, *114*, 325.
- For leading references: (a) Bernsmann, H.; Van den Berg, M.; Hoen, R.; Minnaard, A. J.; Mehler, G.; Reetz, M. T.; De Vries, J. G.; Feringa, B. L. *J. Org. Chem.* **2005**, *70*, 943; (b) Zheng, Z.; Hi, X.-P. *Org. Lett.* **2005**, *7*, 419; (c) Cong-Dong, J.; Pagenkopf, B. L. *J. Org. Chem.* **2004**, *69*, 4177; (d) Huang, H.; Zheng, Z.; Luo, H.; Bai, C.; Hu, X.; Chen, H. *J. Org. Chem.* **2004**, *69*, 2355; (e) Evans, D. A.; Michael, F. E.; Tedrow, J. S.; Campos, K. R. *J. Am. Chem. Soc.* **2003**, *125*, 3534; (f) Tang, W.; Zhang, X. *Chem. Rev.* **2003**, *103*, 3029; (g) Reetz, M. T.; Sell, T.; Meiswinkel, A.; Mehler, G. *Angew. Chem., Int. Ed.* **2003**,

- 42, 790; (h) Hu, A.-G.; Fu, Y.; Xie, J.-H.; Zhou, H.; Wang, L.-X.; Zhou, Q.-L. *Angew. Chem., Int. Ed.* **2002**, *41*, 2348; (i) Zhang, X.; Tang, W. *Angew. Chem., Int. Ed.* **2002**, *41*, 1612; (j) Lee, S.-G.; Zhang, Y. J.; Song, C. E.; Lee, J. K.; Choi, J. H. *Angew. Chem., Int. Ed.* **2002**, *41*, 847; (k) Gridnev, I. D.; Yasutake, M.; Higashi, N.; Imamoto, T. *J. Am. Chem. Soc.* **2001**, *123*, 5268; (l) Gridnev, I. D.; Higashi, N.; Imamoto, T. *J. Am. Chem. Soc.* **2000**, *122*, 10486; (m) Yan, Y.-Y.; RajanBabu, T. V. *Org. Lett.* **2000**, *2*, 4137; (n) Zhang, Z.; Zhu, G.; Jiang, Q.; Xiao, D.; Zhang, X. *J. Org. Chem.* **1999**, *64*, 1774; (o) Zhu, G.; Zhang, X. *J. Org. Chem.* **1998**, *63*, 9590; (p) Burk, M. J.; Wang, Y. M.; Lee, J. R. *J. Am. Chem. Soc.* **1996**, *118*, 5142.
5. All exceptions are with non-phenethyl systems. For a relevant example that uses alternatives to the acetate group, including a trifluoroacetamide, see: Noyori, R.; Ohta, M.; Hsiao, Y.; Kitamura *J. Am. Chem. Soc.* **1986**, *108*, 7117.
6. *Protecting Groups*; Kocienski, P. J., Ed.; Thieme Medical Publishers: New York, 2000; p 185, Chapter 6.
7. Burk, M. J.; Casy, G.; Johnson, N. B. *J. Org. Chem.* **1998**, *63*, 6084.
8. Alternative syntheses of enamides include: (a) Brice, J. L.; Meerdink, J. E.; Stahl, S. S. *Org. Lett.* **2004**, *6*, 1845; (b) Matsubara, R.; Nakamura, Y.; Kobayashi, S. *Angew. Chem., Int. Ed.* **2004**, *43*, 1679; (c) Wallace, D. J.; Klauber, D. J.; Chen, C.-Y.; Volante, R. P. *Org. Lett.* **2003**, *5*, 4749; (d) Suen, Y. H.; Horeau, A.; Kagan, H. B. *Bull. Soc. Chim.* **1965**, *5*, 1454; (e) Mo, J.; Xu, L.; Xiao, J. *J. Am. Chem. Soc.* **2005**, *127*, 751.
9. (a) Boar, R. B.; McGhie, J. F.; Robinson, M.; Barton, D. H. R.; Horwell, D. C.; Stick, R. V. *J. Chem. Soc., Perkin Trans. 1* **1975**, 1237; (b) Barton, D. H. R.; Zard, S. Z. *J. Chem. Soc., Perkin Trans. 1* **1985**, 2191.
10. Le, J. C.-D.; Pagenkopf, B. L. *J. Org. Chem.* **2004**, *69*, 4177.
11. Li, D.; Shi, F.; Guo, S.; Deng, Y. *Tetrahedron Lett.* **2005**, *46*, 671.
12. (a) Blaise, E. E.; Hebd, C. R. *Seances Acad. Sci.* **1901**, *132*, 498; (b) Suen, Y.-H.; Horeau, A.; Kagan, H. B. *Bull. Chim. Soc. Fr.* **1965**, *5*, 1454; (c) Kagan, H. B.; Suen, Y.-H. *Bull. Soc. Chim. Fr.* **1966**, *6*, 1819.
13. (a) Van den Berg, M.; Haak, R. M.; Minnaard, A. J.; deVries, A. H. M.; Vries, J. G.; Feringa, B. L. *Adv. Synth. Catal.* **2002**, *344*, 1003; (b) Storace, L.; Anzalone, P. N.; Confalone, W. P.; Davis, J. M.; Fortunak, J. M.; Giangiordano, M.; Haley, J. J.; Kamholz, K.; Li, H.-Y.; Ma, P.; Nugent, W. A.; Parsons, R. L.; Sheeran, P. J.; Silverman, C. E.; Waltermire, R. E.; Wood, C. C. *Org. Process Res. Dev.* **2002**, *6*, 54.
14. It is assumed that optimization of enantioselectivity and catalyst loading will be conducted on case by case basis.
15. For a review of Josiphos discovery and applications, see: Blaser, H.-U.; Brieden, W.; Pugin, B.; Spindler, F.; Studer, M.; Togni, A. *Top. Catal.* **2002**, *19*, 3; For a recent example of a Josiphos ligand used for the asymmetric hydrogenation of beta-aminoamides, see: Hsiao, Y.; Rivera, N. R.; Rosner, T.; Krska, S. W.; Njolito, E.; Wang, F.; Sun, Y.; Armstrong, J. D., III; Grabowski, E. J. J.; Tillyer, R. D.; Spindler, F.; Malan, C. *J. Am. Chem. Soc.* **2004**, *125*, 1142.