

Corrigendum

**Corrigendum to “Stereoselective total synthesis
of (+)-cardiobutanolide and (+)-3-*epi*-cardiobutanolide
from diacetone D-glucose”**

[Tetrahedron Lett. 47 (2006) 4627]


Palakodety Radha Krishna* and P. V. Narsimha Reddy

D-206/B, Discovery Laboratory, Organic Chemistry Division-III, Indian Institute of Chemical Technology, Hyderabad 500 007, India

Available online 1 August 2006

The following errors are to be noted:

- (1) **Scheme 1**, Page 2: The synthetic sequence starts with the major isomer **6**, which is converted into the requisite isomer **7** by a two-step sequence and not as mentioned (from **7**). The same is described on Page 1, column 2, paragraph 2, lines 8–17.
- (2) **Scheme 1**. Reagents and conditions: (b) BnBr, NaG, THF, 0 °C–rt, 1 h (80%). Please note that it is NaH and not as printed.



Scheme 1. Reagents and conditions: (a) (i) *p*-NO₂-PhCOOH, DEAD, TPP, THF, 0 °C–rt, 6 h; (ii) K₂CO₃, MeOH, rt, 1 h, (85% over two steps); (b) BnBr, NaH, THF, 0 °C–rt, 1 h (80%); (c) 30% aq AcOH, reflux, 6 h (70%); (d) LiAlH₄, THF, 2 h, 0 °C–rt, (65%); (e) TBSCl, imidazole, CH₂Cl₂ (85%); (f) 2,2'-DMP, PPTS, CH₂Cl₂, rt, (85%); (g) TBAF, THF, rt, (90%); (h) (COCl)₂, DMSO, Et₃N, CH₂Cl₂, –78 °C; (i) EtOAc, LiHMDS, THF, –78 °C, 1 h (60% over two steps); (j) 80% aq, AcOH, 6 h; (70%); (k) 10% Pd–C/H₂, MeOH, (80%).

DOI of original article: [10.1016/j.tetlet.2006.04.146](https://doi.org/10.1016/j.tetlet.2006.04.146)

* Corresponding author. Tel.: +33 320964379; fax: +33 320964906; e-mail: prkgenius@iiict.ap.nic.in