

A copper(II)-catalyzed protocol for modified Friedländer quinoline synthesis

Chan Sik Cho,^{a,*} Wen Xiu Ren^b and Sang Chul Shim^{b,*}

^aResearch Institute of Industrial Technology, Kyungpook National University, Daegu 702-701, South Korea

^bDepartment of Applied Chemistry, Kyungpook National University, Daegu 702-701, South Korea


Received 20 June 2006; revised 13 July 2006; accepted 14 July 2006

Available online 4 August 2006

Abstract—2-Aminobenzyl alcohol reacts with an array of ketones in dioxane at 100 °C in the presence of a catalytic amount of CuCl₂ along with KOH under O₂ atmosphere to afford the corresponding quinolines in good yields. 2-Aminobenzyl alcohol is also oxidatively coupled and cyclized with various aldehydes by step-by-step procedure, an initial treatment of 2-aminobenzyl alcohol in the presence of CuCl₂ and KOH in dioxane under O₂ atmosphere and subsequent addition of aldehyde to the mixture followed by stirring under argon atmosphere, to give 3-substituted quinolines in moderate to good yields.

© 2006 Elsevier Ltd. All rights reserved.

Many synthetic methods have been developed and documented for quinolines due to their intrinsic pharmacological and biological activities.¹ Besides conventionally named routes, transition metal-catalyzed reactions for such quinoline skeletons have also been attempted as alternative methods because of the facility and efficiency of reaction and the wide availability of substrate.² As part of our continuing studies toward transition metal-catalyzed cyclization reactions, we also reported on several transition metal-catalyzed routes for quinolines via ruthenium-catalyzed alkyl or alkanol group transfer from alkylamines or alkanolamines to N-atom of anilines^{2b,3,4} (amine exchange reaction or amine scrambling reaction⁵) and palladium-catalyzed coupling and cyclization between 2-iodoaniline and propargylic alcohols.⁶ Among them, in connection with this letter, 2-aminobenzyl alcohol was found to be oxidatively coupled and cyclized with ketones⁷ (Scheme 1, route b) and secondary alcohols⁸ in the presence of a ruthenium catalyst along with a base to give quinolines.^{9,10} However, this protocol, even though superior to conventional Friedländer quinoline synthesis (Scheme 1, route a)¹¹ in a sense of price and stability of 2-aminobenzyl alcohol, led us to seek for a new elegant catalyst alternative since it also has some drawbacks requiring an expensive



Scheme 1.

catalyst and a hydrogen acceptor¹² (or the use of excess starting ketone to 2-aminobenzyl alcohol). Herein, we describe a copper-catalyzed oxidative cyclization of 2-aminobenzyl alcohol with ketones as well as aldehydes leading to quinolines.¹³

The results of several attempted oxidative cyclizations of 2-aminobenzyl alcohol (**1**) with acetophenone (**2a**) are listed in Table 1. Treatment of equimolar amounts of **1** and **2a** in dioxane in the presence of a catalytic amount of CuCl₂ (1 mol %) along with KOH at 100 °C for 5 h afforded 2-phenylquinoline (**3a**) in 65% yield without any identifiable products (run 1). In contrast to our recent report on ruthenium-catalyzed synthesis of quinolines from **1** and ketones,⁷ no direct transfer hydrogenation product 1-phenylethanol was produced. This result indicates that the present reaction proceeds irrespective of transfer hydrogenation from **1** to **2a**. In the case of ruthenium-catalyzed version, 2 equiv of **2a**

Keywords: Aldehydes; 2-Aminobenzyl alcohol; Copper catalyst; Ketones; Quinolines.

* Corresponding authors. Tel.: +82 53 950 7318; fax: +82 53 950 6594 (C.S.C.); e-mail: cscho@knu.ac.kr

Table 1. Cu-Catalyzed optimization of conditions for the reaction of **1** with **2a**

Nc1ccccc1CO + CC(=O)c1ccccc1 $\xrightarrow[\text{KOH, dioxane}]{\text{CuCl}_2/\text{O}_2 (1 \text{ atm})}$ Nc1ccc2ccccc2n1C3=CC=CC=C3

1 **2a** **3a**

Run	[2a]/[1]	KOH (mmol)	Cu (mmol)	Time (h)	Yield ^a (%)
1	1	1	CuCl ₂ (0.01)	5	65
2	1	3	CuCl ₂ (0.01)	5	70
3	1.2	3	CuCl ₂ (0.01)	5	77
4	2	3	CuCl ₂ (0.01)	5	77
5	1.2	3	CuCl ₂ (0.01)	20	79
6	1.2	3	CuCl ₂ (0.005)	5	73
7	1.2	3	CuCl ₂ (0.05)	5	79
8	1.2	3	CuCl (0.01)	5	78
9 ^b	1.2	3	CuCl ₂ (0.01)	5	51

Reaction conditions: **1** (1 mmol), dioxane (5 mL), 100 °C, O₂ (1 atm).

^a Isolated yield based on **1**.

^b The reaction was carried out under air.

to **1** was necessary for the effective formation of quinolines.⁷ This could be due to partial consumption of **2a** leading to 1-phenylethanol by ruthenium-catalyzed transfer hydrogenation of **2a** by **1**.¹² A slightly increased yield of **3a** was observed with an increased amount of KOH (run 2). The molar ratio of **2a** to **1** affected the yield of **3a**, higher molar ratio up to [**2a**]/[**1**] = 1.2 resulting in the effective formation of **3a** (runs 3 and 4). The reaction gave no significant change on the quinoline yield for a longer reaction time (run 5). The optimization for CuCl₂ amount was achieved with 1 mol % based on **1** (runs 3, 6 and 7). The reactivity toward the formation of **3a** using CuCl under the employed conditions was revealed to be as similarly effective as that using CuCl₂ (run 8). However, performing the reaction under air atmosphere resulted in a lower yield of **3a** compared with that under O₂ atmosphere (run 9).

Having established the reaction conditions, various ketones **2** were subjected to react with **1** in order to investigate the reaction scope and several representative results are summarized in Table 2. Various aryl(methyl) ketones (**2a–i**) having electron donating and withdrawing substituents on the aromatic ring were readily coupled and cyclized with **1** to give the corresponding quinolines (**3a–i**) in the range of 42–82% yields. Here again, the conventional transfer hydrogenated aryl(methyl) carbinols were not detected at all on GLC analysis.¹² The position and electronic nature of the substituent on the aromatic ring of aryl(methyl) ketones had no relevance to quinoline yield except for 4'-cyanoacetophenone (**2i**). The reaction proceeds likewise with heteroaryl(methyl) ketone **2j** to give the corresponding quinoline **3j** in similar yield. 2'-Acetonaphthone (**2k**) was also readily oxidatively coupled and cyclized with **1** to afford 2-(2-naphthyl)quinoline (**3k**) in 77% yield. Similar reaction rate and yield were observed with alkyl(aryl) ketone **2l**, which has only methylene reaction site. In the cases of alkyl(methyl) ketones (**2m** and **2n**), the corresponding quinolines were produced as a regioisomeric mixture, favoring cyclization at less-hindered methyl position over α -methylene.^{7,8,11c} With dialkyl ketone **2o** having only methylene reaction site, the

corresponding quinoline **3o** was also formed and the yield was lower than that when previously described ketones such as aryl(methyl) ketones, alkyl(aryl) ketones and alkyl(methyl) ketones were used. Cyclic ketones such as 4-phenylcyclohexanone (**2p**) and 1-tetralone (**2q**) were also reacted with **1** to give 3-phenyl-1,2,3,4-tetrahydroacridine (**3q**) and 5,6-dihydrobenzo[*c*]acridine (**3r**) in 63% and 65% yields, respectively.

We then examined a similar oxidative cyclization between **1** with aldehydes.¹³ Table 3 shows several attempted results for the reaction between **1** and octyl aldehyde (**4a**). Treatment of **1** with **4a** under similar catalytic system used in the reaction of **1** with ketones afforded 3-hexylquinoline (**3r**) in only 29% yield (run 1). However, step-by-step procedure, an initial treatment of **1** in the presence of a catalytic amount of CuCl₂ along with KOH in dioxane for 5 h under O₂ atmosphere and subsequent addition of **4a** to the mixture followed by stirring for 20 h resulted in an increased yield of **3r** (run 2). Performing the second reaction stage under air gave no significant change on the product yield (run 3). On the other hand, when the atmosphere of second reaction stage was changed from O₂ or air to argon, a considerably increased yield of **3r** was obtained (run 4). However, increasing the amount of CuCl₂ catalyst to 10 mol % rather lowered the quinoline yield (run 5).

Given suitable reaction conditions, various aldehydes **4** were subjected to react with **1** in order to investigate the reaction scope and several representative results are summarized in Table 4. From the reaction between **1** and straight aldehydes (**4a–d**), the 3-substituted quinolines (**3r–u**) were formed in the range of 48–67% yields. The product yield had no relevance to the chain length of **4a–d**. In the case of valeraldehyde (**4b**), the second stage reaction was carried out at a lower reaction temperature to obtain an allowable yield of 3-propylquinoline (**3s**). Hydrocinnamaldehyde (**4e**) having a phenyl group at position 3 reacts similarly with **1** to give 3-benzylquinoline (**3v**) in 46% yield. The reaction proceeds likewise with isovaleraldehyde (**4f**) and 3-phenylbutyraldehyde (**4g**), which have substituents such as methyl and


Table 2. Cu-Catalyzed synthesis of quinolines **3** from **1** and **2**

Ketones 2	Quinolines 3	Yield (%)
R = Ph (2a)	R = Ph (3a)	77
R = 4-MeC ₆ H ₄ (2b)	R = 4-MeC ₆ H ₄ (3b)	79
R = 3-MeC ₆ H ₄ (2c)	R = 3-MeC ₆ H ₄ (3c)	80
R = 2-MeC ₆ H ₄ (2d)	R = 2-MeC ₆ H ₄ (3d)	71
R = 4-MeOC ₆ H ₄ (2e)	R = 4-MeOC ₆ H ₄ (3e)	82
R = 2-MeOC ₆ H ₄ (2f)	R = 2-MeOC ₆ H ₄ (3f)	75
R = 4-FC ₆ H ₄ (2g)	R = 4-FC ₆ H ₄ (3g)	68
R = 3-CF ₃ C ₆ H ₄ (2h)	R = 3-CF ₃ C ₆ H ₄ (3h)	71
R = 4-CNC ₆ H ₄ (2i)	R = 4-CNC ₆ H ₄ (3i)	42
R = 2-Thienyl (2j)	R = 2-Thienyl (3j)	62
R = 2-Naphthyl (2k)	R = 2-Naphthyl (3k)	77
		62
2l	3l	
		48 ^a
2m	3m	
		44 ^b
2n	3n	
		48
2o	3o	
		63
2p	3p	
		65
2q	3q	

Reaction conditions: **1** (1 mmol), **2** (1.2 mmol), CuCl₂ (0.01 mmol), KOH (3 mmol), dioxane (5 mL), O₂ (1 atm), 100 °C, for 5 h.

^a 3-Benzyl-2-methylquinoline was also formed in 18% yield.

^b 3-Butyl-2-methylquinoline was also formed in 23% yield.

Table 3. Cu-Catalyzed oxidative cyclization of **1** with **4a** under several conditions

Run	CuCl ₂ (mmol)	Time (h) and atmosphere	Yield (%)
1	0.01	20 (O ₂)	29
2	0.01	5 (O ₂) and 20 (O ₂)	47
3	0.01	5 (O ₂) and 20 (Air)	47
4	0.01	5 (O ₂) and 20 (Ar)	67
5	0.1	5 (O ₂) and 20 (Ar)	54


Reaction conditions: **1** (1.5 mmol), KOH (3 mmol), dioxane (3 mL), 100 °C; **4a** (1 mmol), dioxane (3 mL), 100 °C.

Table 4. Cu-Catalyzed synthesis of quinolines **3** from **1** and **4**

Aldehydes 2	Quinolines 3	Yield (%)
R = Hexyl (4a)	R = Hexyl (3r)	67
R = Propyl (4b)	R = Propyl (3s)	48 ^a
R = Pentyl (4c)	R = Pentyl (3t)	48
R = Decyl (4d)	R = Decyl (3u)	50
		46
		64 ^a
		46
		41

Reaction conditions: **1** (1.5 mmol), CuCl₂ (0.01 mmol), KOH (3 mmol), dioxane (3 mL), O₂ (1 atm), 100 °C, for 5 h; **4** (1 mmol), dioxane (3 mL), Ar, 100 °C, for 20 h.

^aSecond stage (90 °C).

**Scheme 2.**

phenyl at position 3, and the corresponding 3-isopropylquinoline (**3w**) and 3-(1-phenylethyl)quinoline (**3x**) were also formed in 64% and 46% yields, respectively. Here again, in the case of **4f** the second step reaction was performed at a lower reaction temperature because of low boiling point of **4f**. The reaction of phenylacetaldehyde (**4h**), which has a phenyl substituent at position 2, with **1** also proceeds to give 3-phenylquinoline (**3y**).

As concerns the reaction pathway, it seems to proceed via the initial oxidation of **1** to 2-aminobenzaldehyde (**5**) under CuCl₂/O₂ system, which in turn triggers cross aldol condensation with **2a** to give enone **6** (Scheme 2).^{11d,14,15} This is followed by cyclodehydration to form quinoline **3a**. An alternative route for **3a** involves ketimine aldehyde **8** formation, which is formed by the condensation between **5** and **2a** as well as the oxidation of

ketimine alcohol **7** initially formed from **1** and **2a**, and subsequent intramolecular aldol-type condensation of **8**. The reaction with aldehyde **4h** for quinoline **3y** also similarly proceeds via an aldimine **9** formation followed by the intramolecular aldol reaction.

In summary, we have shown that 2-aminobenzyl alcohol can be oxidatively cyclized with ketones as well as aldehydes in the presence of a copper catalyst and a base to afford quinolines in good yields. We believe that the present reaction will work as a useful procedure for transition metal-catalyzed modified Friedländer quinoline synthesis since it proceeds not only in the presence of a cheap copper catalyst but also in the absence of a hydrogen acceptor.

Acknowledgements

The present work was supported by the Korea Research Foundation Grant funded by Korea Government (MOEHRD, Basic Research Promotion Fund) (KRF-2005-015-C00264). C.S.C. gratefully acknowledges a Research Professor Grant of Kyungpook National University (2005).

References and notes

- (a) Jones, G. In *Comprehensive Heterocyclic Chemistry*; Katritzky, A. R., Rees, C. W., Eds.; Pergamon: New York, 1984; Vol. 2, p 395; (b) Arcadi, A.; Chiarini, M.; Giuseppe, S. D.; Marinelli, F. *Synlett* **2003**, 203, and references cited therein.
- (a) Amii, H.; Kishikawa, Y.; Uneyama, K. *Org. Lett.* **2001**, 3, 1109, and references cited therein; (b) Cho, C. S.; Oh, B. H.; Kim, J. S.; Kim, T.-J.; Shim, S. C. *Chem. Commun.* **2000**, 1885, and references cited therein.
- (a) Cho, C. S.; Oh, B. H.; Shim, S. C. *Tetrahedron Lett.* **1999**, 40, 1499; (b) Cho, C. S.; Oh, B. H.; Shim, S. C. *J. Heterocycl. Chem.* **1999**, 36, 1175; (c) Cho, C. S.; Kim, J. S.; Oh, B. H.; Kim, T.-J.; Shim, S. C. *Tetrahedron* **2000**, 56, 7747; (d) Cho, C. S.; Kim, T. K.; Kim, B. T.; Kim, T.-J.; Shim, S. C. *J. Organomet. Chem.* **2002**, 650, 65; (e) Cho, C. S.; Lee, N. Y.; Kim, T.-J.; Shim, S. C. *J. Heterocycl. Chem.* **2004**, 41, 423.
- For our recent report on synthesis of indoles via similar ruthenium-catalyzed alkanol group transfer from alkanolamines to N-atom of anilines: (a) Cho, C. S.; Lim, H. K.; Shim, S. C.; Kim, T. J.; Choi, H.-J. *Chem. Commun.* **1998**, 995; (b) Cho, C. S.; Kim, J. H.; Shim, S. C. *Tetrahedron Lett.* **2000**, 41, 1811; (c) Cho, C. S.; Kim, J. H.; Kim, T.-J.; Shim, S. C. *Tetrahedron* **2001**, 57, 3321; (d) Cho, C. S.; Kim, J. H.; Choi, H.-J.; Kim, T.-J.; Shim, S. C. *Tetrahedron Lett.* **2003**, 44, 2975.
- For a review: Murahashi, S.-I. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 2443.
- (a) Cho, C. S. *J. Organomet. Chem.* **2005**, 690, 4094; (b) Cho, C. S.; Lee, N. Y.; Kim, T.-J.; Shim, S. C. *J. Heterocycl. Chem.* **2004**, 41, 409.
- Cho, C. S.; Kim, B. T.; Kim, T.-J.; Shim, S. C. *Chem. Commun.* **2001**, 2576.
- Cho, C. S.; Kim, B. T.; Choi, H.-J.; Kim, T.-J.; Shim, S. C. *Tetrahedron* **2003**, 59, 7997.
- For similar ruthenium- and iridium-catalyzed oxidative cyclization of 2-aminobenzyl alcohol with ketones leading to quinolines: (a) Motokura, K.; Mizugaki, T.; Ebitani, K.; Kaneda, K. *Tetrahedron Lett.* **2004**, 45, 6029; (b) Martínez, R.; Brand, G. J.; Ramón, D. J.; Yus, M. *Tetrahedron Lett.* **2005**, 46, 3683; (c) Taguchi, K.; Sakaguchi, S.; Ishii, Y. *Tetrahedron Lett.* **2005**, 46, 4539.
- It has recently been found that carbonyl compounds (or secondary alcohols) are coupled with primary alcohols in the presence of a ruthenium catalyst and a base: (a) Cho, C. S.; Kim, B. T.; Kim, T.-J.; Shim, S. C. *J. Org. Chem.* **2001**, 66, 9020; (b) Cho, C. S.; Kim, B. T.; Kim, T.-J.; Shim, S. C. *Tetrahedron Lett.* **2002**, 43, 7987; (c) Cho, C. S.; Kim, B. T.; Kim, H.-S.; Kim, T.-J.; Shim, S. C. *Organometallics* **2003**, 22, 3608.
- (a) Friedländer, P. *Chem. Ber.* **1882**, 15, 2572; For a review, see: (b) Cheng, C.-C.; Yan, S.-J. *Org. React.* **1982**, 28, 37; For a regioselective Friedländer quinoline synthesis, see: (c) Hsiao, Y.; Rivera, N. R.; Yasuda, N.; Hughes, D. L.; Reider, P. J. *Org. Lett.* **2001**, 3, 1101; (d) Muchowski, J. M.; Maddox, M. L. *Can. J. Chem.* **2004**, 82, 461.
- For recent reviews on transition metal-catalyzed transfer hydrogenation, see: (a) Zassinovich, G.; Mestroni, G.; Gladiali, S. *Chem. Rev.* **1992**, 92, 1051; (b) Bäckvall, J.-E.; Chowdhury, R. L.; Karlsson, U.; Wang, G. In *Perspectives in Coordination Chemistry*; Williams, A. F., Floriani, C., Merbach, A. E., Eds.; VCH: New York, 1992; pp 463–486; (c) Noyori, R.; Hashiguchi, S. *Acc. Chem. Res.* **1997**, 30, 97; (d) Naota, T.; Takaya, H.; Murahashi, S.-I. *Chem. Rev.* **1998**, 98, 2599; (e) Palmer, M.; Wills, M. *Tetrahedron: Asymmetry* **1999**, 10, 2045.
- For ruthenium-catalyzed oxidative cyclization of 2-aminobenzyl alcohol with aldehydes leading to 3-substituted quinolines Cho, C. S.; Ren, W. X.; Shim, S. C. *Bull. Korean Chem. Soc.* **2005**, 26, 2038.
- Hudlický, M. *Oxidation in Organic Chemistry*; American Chemical Society: Washington, DC, 1990; pp 114–126.
- We confirmed in a separate experiment that **1** was oxidized to **5** in 50% yield with 85% conversion of **1** under CuCl₂/O₂/KOH catalytic system, however, benzyl alcohol was not effectively oxidized to benzaldehyde (11% yield) under the same catalytic system.