

Lewis acid-promoted α -hydroxy β -dicarbonyl to α -ketol ester rearrangement

Xuechuan Hong, José M. Mejía-Oneto and Albert Padwa*

Department of Chemistry, Emory University, Atlanta, GA 30322, USA


Received 28 August 2006; revised 13 September 2006; accepted 14 September 2006

Available online 5 October 2006

Abstract—The decarbomethoxylation reaction of a substituted α -hydroxy- α -carbomethoxy pentacyclic substituted ketone, used as an advanced intermediate in the synthesis of the alkaloid aspidophytine, can be effected by heating with MgI_2 in CH_3CN . The reaction was shown to proceed by a novel α -hydroxy β -dicarbonyl to α -ketol ester rearrangement. It was possible to isolate a carbonate intermediate in 75% yield, thereby providing support for the proposed pathway.
© 2006 Elsevier Ltd. All rights reserved.

The Rh(II)-catalyzed cyclization/cycloaddition cascade of α -diazo carbonyl compounds developed in our laboratories has proven to be an extremely efficient process for the synthesis of a variety of aza polycyclic ring systems.¹ Intramolecular trapping of the carbonyl ylide dipole generated from the domino Rh(II)-reaction proceeds in a highly regio and stereoselective manner and has been used in the construction of several heterocyclic natural products.^{2–4} For example, this domino cascade was successfully applied in our laboratory toward the synthesis of the alkaloid (\pm)-lycophodine (Scheme 1).⁵ The initially formed cycloadduct **2**, obtained from the Rh(II)-cascade reaction of **1**, was treated with a Lewis acid and provided the rearranged tetracyclic lactam **3** derived from a Pictet–Spengler type cyclization of a transient *N*-acyliminium ion.⁶ Although lactam **3** contains the proper ring skeleton needed for lycophodine, it is overfunctionalized. The three-step sequence that we used to remove both the hydroxyl and ester functionalities involved (i) a Barton–McCombie deoxygenation,⁷ (ii) ester hydrolysis, and (iii) thermal decarboxylation of the resulting carboxylic acid. Compound **4** was eventually transformed into (\pm)-lycophodine.

More recently, in our successful synthesis of the alkaloid (\pm)-aspidophytine,⁸ it was also necessary to remove the hydroxyl and carbomethoxy functionalities from the advanced intermediate **7**, which was obtained by treating




Scheme 1.

cycloadduct **6** with $BF_3 \cdot OEt_2$ (Scheme 2). In this case, the three-step protocol previously used for lycophodine resulted in only very low yields of ketone **8**. During the course of preparing ketone **8** for an eventual synthesis of aspidophytine, we investigated the dealkoxycarbonylation–dehydroxylation of lactam **7** and related systems under several reaction conditions. The results that we obtained are not only useful for the future construction of molecules of the type discussed here, but also allow for the facile chemoselective removal of either (or both) the hydroxyl and/or carboalkoxy functionalities.

Dealkoxycarbonylation of activated esters via nucleophilic dealkylation⁹ represents a one-step process for

* Corresponding author. Tel.: +404 727 0283; fax: +404 727 6629; e-mail: chemap@emory.edu




Scheme 2.

the removal of the alkyl ester entity, particularly when structurally hindered. The reaction usually involves heating of the substrate in a dipolar aprotic solvent in the presence of a nucleophile. Following Krapcho's original development of this approach,¹⁰ which employed NaCl in DMSO, many other nucleophiles have been found to be applicable,⁹ including thiolates, *tert*-butoxide, thiocyanate, amines, and acetate. For malonates or β -ketoesters, this displacement is followed by rapid decarboxylation. Several accounts have also appeared describing synthetic transformations which utilize the intermediate anion generated in this reaction when run under anhydrous aprotic conditions.¹¹ Since the introduction of this procedure by Krapcho, many applications of its synthetic utility to complex molecules have been reported.⁹


On the basis of the earlier decarboalkoxylation studies, our first attempts to convert hydroxyester **7** into α -hydroxyketone **9** made use of the one-pot Krapcho protocol. This involved heating **7** at reflux in DMSO in the presence of sodium chloride which indeed furnished compound **9**, but only in 20% overall yield. After numerous attempts at improving the yield of the decarboalkoxylation reaction, the procedure that we eventually settled on for the aspidophytine synthesis involved heating **7** with MgI₂ in refluxing acetonitrile that contained a small amount of water. Gratifyingly, this method resulted in the formation of alcohol **9** in 75% isolated yield. Acetylation of **9** with acetyl chloride/NEt₃ afforded the corresponding acetate, which was subsequently reduced using SmI₂ to give **8** in 90% yield for both steps. Although a cursory analysis of the conditions used suggests that a Krapcho decarboalkoxylation reaction first occurred, the isolation of small quantities of carbonate **10** from the mixture was somewhat surprising and led us to examine the MgI₂ reaction in more detail using several model systems to probe the details of this process.

Toward this goal, we carried out experiments using α -hydroxy ketoesters **11**, **14**, and **17** in order to ascertain

the limitations of the decarboalkoxylation process employing MgI₂ and CH₃CN as solvent. The optimized reaction conditions utilize 1.5 equiv of MgI₂ in acetonitrile that contains a small amount of water at a temperature of 81 °C for 2 h. The reaction gives optimized yields of α -hydroxy ketones **12**, **15**, and **18** (Scheme 3) using substrate concentrations of ca. 0.1 M on scales up to 10 mmol. By employing this method, we were able to isolate α -hydroxy substituted cyclohexanones **12** and **15** in 84% and 81% yield, respectively. A lower yield of the hemiketal dimer **18** (42%) was obtained with the more reactive cyclopentanone system **17**. During the course of these studies, we also found that it is possible to selectively remove the hydroxyl group prior to the decarboalkoxylation reaction. Thus, treatment of the starting α -hydroxy ketoesters shown in Scheme 3 with SmI₂ in HMPA/THF at 0 °C produced the dehydroxylated products **13**, **16**, and **19** in 56%, 94%, and 97% yield, respectively. It was not necessary to convert the hydroxyl group into the corresponding acetate in order for the reduction to proceed. Most interestingly, when the ketoesters were exposed to the MgI₂/CH₃CN conditions previously employed, only recovered starting material was obtained, with no signs of any decarboalkoxylation. This result and the previous observation that small quantities of carbonate **10** could be isolated from the reaction of **7** with MgI₂ lead us to propose an alternate mechanism for the decarboalkoxylation reaction. Thus, rather than undergoing the Krapcho reaction, we believe that the formation of **9** from **7** represents a rare example of a Lewis acid-promoted α -hydroxy β -dicarbonyl to α -ketol ester rearrangement. A plausible mechanism for this process involves coordination of the MgI₂ with the more basic ester site followed by attack of the hydroxyl oxygen on the neighboring carbomethoxy group. Epoxide **22** is formed as a transient species which readily isomerizes to the magnesium enolate **23**. This is followed by ketonization to furnish carbonate **24**, which undergoes further hydrolysis with the H⁺ produced in the reaction with MgI₂ to provide the final product **25**. Indeed, heating an authentic sample of the carbonate derived from **12** with MgI₂ in acetonitrile afforded the same decarboalkoxylation product in 96% yield.^{12,13} The success of this reaction



Scheme 3.



Scheme 4.

is probably due to activation of the ester by chelation of the Mg(II) cation with both carbonyl groups. Analogous base-catalyzed rearrangements of α -hydroxy β -diketones to α -keto esters have been reported by Davis et al.¹⁴ and others.^{15–18} The pathway shown in Scheme 4 is supported by the isolation of carbonate 10 when the reaction of 7 was carried out with Cs₂CO₃ in acetonitrile at 81 °C for 1 h. Under these conditions, carbonate 10 was obtained in 75% yield and could be readily hydrolyzed to 9 when subjected to mild acidic conditions.

In summary, we have uncovered a novel Lewis acid-promoted α -hydroxy β -dicarbonyl to α -ketol ester rearrangement. This reaction can be used to give ready access to carbonate 10 which corresponds to a key intermediate en route to (\pm)-aspidoptyne. Moreover, we have investigated the decarbomethoxylation of a number of related α -hydroxy carbomethoxy substituted cycloalkanones and have developed conditions for the chemoselective removal of either (or both) the hydroxyl and/or carboalkoxy functionalities. Application of the method toward the synthesis of a number of aspidosperma alkaloids is currently underway and will be described in forthcoming publications.

Acknowledgements

We appreciate the financial support provided by the National Institutes of Health (GM 059384) and the National Science Foundation (CHE-0450779).

References and notes

- (a) Padwa, A.; Hornbuckle, S. F. *Chem. Rev.* **1991**, *91*, 263; (b) Padwa, A.; Weingarten, M. D. *Chem. Rev.* **1996**, *96*, 223.
- (a) Padwa, A. *Acc. Chem. Res.* **1991**, *24*, 22; (b) Padwa, A.; Carter, S. P.; Nimmesgern, H. *J. Org. Chem.* **1986**, *51*, 1157; (c) Padwa, A.; Carter, S. P.; Nimmesgern, H.; Stull, P. D. *J. Am. Chem. Soc.* **1988**, *110*, 2894; (d) Padwa, A.; Fryxell, G. E.; Zhi, L. *J. Org. Chem.* **1988**, *53*, 2877; (e) Padwa, A.; Chinn, R. L.; Zhi, L. *Tetrahedron Lett.* **1989**, *30*, 1491; (f) Padwa, A.; Hornbuckle, S. F.; Fryxell, G. E.;

- Stull, P. D. *J. Org. Chem.* **1989**, *54*, 817; (g) Padwa, A.; Sandanayaka, V. P.; Curtis, E. A. *J. Am. Chem. Soc.* **1994**, *116*, 2667; (h) Curtis, E. A.; Sandanayaka, V. P.; Padwa, A. *Tetrahedron Lett.* **1995**, *36*, 1989.
- (a) Padwa, A.; Precedo, L.; Semones, M. *J. Org. Chem.* **1999**, *64*, 4079; (b) Padwa, A.; Curtis, E. A.; Sandanayaka, V. P. *J. Org. Chem.* **1997**, *62*, 1317.
- (a) Kinder, F. R., Jr.; Bair, K. W. *J. Org. Chem.* **1994**, *59*, 6965; (b) Chen, B.; Ko, R. Y. Y.; Yuen, M. S. M.; Cheng, K.-F.; Chiu, P. *J. Org. Chem.* **2003**, *68*, 4195; (c) Hodgson, D. M.; Avery, T. D.; Donohue, A. C. *Org. Lett.* **2002**, *4*, 1809; (d) Dauben, W. G.; Dinges, J.; Smith, T. C. *J. Org. Chem.* **1993**, *58*, 7635.
- Padwa, A.; Brodney, M. A.; Marino, J. P., Jr.; Sheehan, S. M. *J. Org. Chem.* **1997**, *62*, 78.
- (a) Cox, E. D.; Cook, J. M. *Chem. Rev.* **1995**, *5*, 1797; (b) Claret, P. A. In *Comprehensive Organic Chemistry*; Barton, D., Ollis, W. D., Eds.; Pergamon: Oxford, 1979; Vol. 4, p 209.
- Barton, D. H. R.; McCombie, S. W. *J. Chem. Soc., Perkin Trans. 1* **1975**, 1574.
- Oneto, J. M. M.; Padwa, A. *Org. Lett.* **2006**, *8*, 3275.
- (a) Krapcho, A. P. *Synthesis* **1982**, 805, 893; (b) McMurry, J. *Org. React.* **1976**, *101*, 187; (c) Stevens, R. V.; Lee, A. W. M. *J. Am. Chem. Soc.* **1979**, *101*, 7032.
- (a) Krapcho, A. P.; Glynn, G. A.; Grenon, B. *J. Tetrahedron Lett.* **1967**, 215; (b) Krapcho, A. P.; Weimaster, J. F.; Eldridge, J. M.; Jahngen, E. G. E., Jr.; Lovey, A. J.; Stephens, W. P. *J. Org. Chem.* **1978**, *43*, 138.
- (a) Bunce, R. A.; Dowdy, E. D.; Jones, P. B.; Holt, E. M. *J. Org. Chem.* **1993**, *58*, 7143; (b) Takei, S.; Kowano, Y. *Tetrahedron Lett.* **1975**, *16*, 4389; (c) Belletire, J. L.; Walley, D. R. *Tetrahedron Lett.* **1983**, *24*, 1475; (d) Böhler, G.; Böhler, P.; Knorr, R. *Chem. Ber.* **1990**, *123*, 2167.
- It should be noted that heating an authentic sample of the carbonate derived from 12 in acetonitrile in the absence of MgI₂ resulted in the recovery of starting material. The decarbomethoxylation of the carbonate was completely suppressed when the reaction was carried out in the presence of proton sponge. By executing the reaction of 11 in MgI₂ in acetonitrile in the presence of acetic anhydride, it was possible to isolate the corresponding acetate in 78% yield.
- A typical magnesium iodide decarbomethoxylation reaction.* To a 0.9 g (1.9 mmol) sample of 7 in acetonitrile (150 mL) was added 1.1 g (4 mmol) of magnesium iodide and the mixture was heated at reflux for 4 h. The solution was allowed to cool to rt, concentrated under reduced pressure and the residue was taken up in methylene chloride (50 mL) and H₂O (50 mL). This solution was added to a saturated aqueous solution of NaHCO₃ (200 mL). The organic layer was separated and the aqueous layer was extracted twice with methylene chloride, dried over MgSO₄, and concentrated under reduced pressure to give 0.6 g (75%) of 7 as a white solid; mp 210–212 °C; IR (neat) 1794, 1720, 1609, 1494, 1263, and 1193 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.30–1.45 (m, 1H), 1.58–1.81 (m, 2H), 2.27 (d, 1H, *J* = 17.4 Hz), 2.34 (d, 1H, *J* = 17.4 Hz), 2.35–2.42 (m, 1H), 2.80 (d, 1H, *J* = 18.0 Hz), 2.93 (dt, 1H, *J* = 14.0 and 6.7 Hz), 3.14 (s, 3H), 3.27 (dd, 1H, *J* = 18.0 and 0.8 Hz), 3.68 (d, 1H, *J* = 3.6 Hz), 3.69 (s, 3H), 3.83 (s, 3H), 3.85 (d, 1H, *J* = 3.6 Hz), 4.21–4.30 (m, 1H), 4.84 (t, 1H, *J* = 3.6 Hz), 6.48 (d, 1H, *J* = 8.6 Hz), and 6.89 (d, 1H, *J* = 8.6 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 20.4, 34.1, 37.3, 40.0, 41.4, 44.7, 51.7, 53.4, 56.2, 60.4, 78.0, 98.0, 105.1, 120.4, 122.9, 135.3, 146.5, 155.5, 170.4, 172.5, and 207.8; Anal. Calcd for C₂₂H₂₄N₂O₇: C, 61.67; H, 5.65; N, 6.54. Found: C, 61.18; H, 6.21; N, 5.94.

14. (a) Davis, F. A.; Clark, C.; Kumar, A.; Chen, B.-C. *J. Org. Chem.* **1994**, 59, 1184; (b) Davis, F. A.; Liu, H.; Chen, B.-C.; Zhou, P. *Tetrahedron* **1998**, 54, 10481.
15. (a) Rubin, M. B.; Inbar, S. *Tetrahedron Lett.* **1979**, 20, 5021; (b) Rubin, M. B.; Inbar, S. *J. Org. Chem.* **1988**, 53, 3355.
16. Blatt, A. H.; Hawkins, W. L. *J. Am. Chem. Soc.* **1936**, 58, 81.
17. House, H. O.; Gannon, W. F. *J. Org. Chem.* **1958**, 23, 879.
18. Charest, M. G.; Siegel, D. R.; Myers, A. G. *J. Am. Chem. Soc.* **2005**, 127, 8292.