

Corrigendum

Corrigendum to “A highly diastereoselective approach to tetrahydrofurans via [3+2] cycloadditions of silylmethyl-substituted cyclopropanes with aldehydes and ketones”

[Tetrahedron Lett. 47 (2006) 8043]

Archana Gupta and Veejendra K. Yadav*

Department of Chemistry, Indian Institute of Technology, Kanpur 208 016, India

Available online 7 November 2006

Table 1 in the above manuscript should have read as follows:

Table 1. Screening of Lewis acids for the reaction of **1a** with furfural

Entry	Lewis acid	Equiv	dr (% yield)
1	Sn(OTf) ₂ ^{a,b}	0.15	6.3:1 (34)
2	BF ₃ ·OEt ₂ ^{b,c}	1.2	2.7:1 (23)
3	Et ₂ AlCl ^a	1.2	2.4:1 (38)
4	SnCl ₄ ^{c,d}	0.05	6.3:1 (59)
5	Sc(OTf) ₃ ^a	0.15	3.5:1 (100)

^a Reaction was carried out at rt for 3 h.

^b The yields are based on the reacted starting cyclopropane derivative.

^c Reaction was carried out at 0 °C–rt for 4 h.

^d 3-Carbomethoxy-5-*t*-butyldiphenylsilylmethyl tetrahydrofuran-2-one one was formed in 30% isolated yield.