

A convenient and regioselective oxidative bromination of electron-rich aromatic rings using potassium bromide and benzyltriphenylphosphonium peroxymonosulfate under nearly neutral reaction conditions

Hadi Adibi,^{a,*} Abdol R. Hajjipour^{b,c} and Majid Hashemi^d

^aDepartment of Medicinal Chemistry, School of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah 67168-67346, Iran

^bPharmaceutical Research Laboratory, College of Chemistry, Isfahan University of Technology, Isfahan 84156, Iran

^cDepartment of Pharmacology, University of Wisconsin, Medical School, 1300 University Avenue, Madison, Wisconsin 53706-1532, USA

^dResearch and Petroleum Engineering Center of Kermanshah (A Branch of RIPI), Kermanshah 67146-77717, Iran

Received 1 November 2006; revised 25 November 2006; accepted 8 December 2006

Available online 4 January 2007

Abstract—Regioselective oxidative bromination of electron-rich aromatic rings has been studied using potassium bromide as a bromine source in the presence of benzyltriphenylphosphonium peroxymonosulfate as oxidant under nearly neutral reaction conditions. In most cases we obtained monobrominated derivatives regioselectively and in good to high yields without the aid of strong acids. © 2006 Elsevier Ltd. All rights reserved.

Brominated aromatic compounds have gained increasing interest as versatile intermediates for the synthesis of biologically active compounds such as potent anti-tumour, antibacterial, antifungal, antiviral, and anti-oxidizing agents.¹ Conventional bromination methods typically use elemental bromine, generating toxic, and corrosive hydrogen bromide, leading to environmental pollution.² Traditional methods of aromatic bromination involve the use of nonselective hazardous acidic reagents such as mineral acids and metal halides, which can lead to separation difficulties and toxic and corrosive wastes. Examples of conventional and traditional methods include: Br₂–Lewis acids,³ NBS–H₂SO₄–CF₃CO₂H,⁴ NBS–PTSA,⁵ NBS–NaOH,⁶ NBS–SiO₂,⁷ Br₂–Al₂O₃,⁸ Br₂–Zeolite,⁹ NBS–Amberlyst,¹⁰ NBS–HZSM-5,¹¹ Clayzib,¹² *tert*-BuOOH– or H₂O₂–HBr,¹³ and HBr–DMSO.¹⁴ The replacement of such reagents with non-toxic and more selective reagents is very desirable and represents an important goal in the context of

clean synthesis. The use of potassium bromide as brominating reagent in the presence of sodium tungstate or ammonium molybdate as catalyst using hydrogen peroxide or sodium perborate as an oxidant has been developed.¹⁵ Oxone[®] (2KHSO₅·KHSO₄·K₂SO₄) is an inexpensive, water-soluble, and stable oxidizing reagent that is commercially available, however, this reagent is insoluble in organic solvents and buffering is needed due to its acidity.¹⁶ Recently, sodium bromide or potassium bromide combined with Oxone[®] has been used as an effective reagent for the bromination of activated arenes.¹⁷ Replacement of the potassium cation in potassium peroxymonosulfate with quaternary phosphonium is very desirable and increases its solubility in organic solvents.

Recently, we reported benzyltriphenylphosphonium peroxymonosulfate **BTPPMS** as a mild, inexpensive and efficient reagent in organic transformations.¹⁸ Following our continued interest in **BTPPMS** and in the course of our studies on the halogenation of organic compounds,¹⁹ we herein report highly efficient and selective bromination reactions of activated arenes employing potassium bromide as the bromine source and **BTPPMS** as the oxidant under non-aqueous conditions.

Keywords: Activated arenes; Benzyltriphenylphosphonium peroxy-monosulfate; Oxidative bromination; Potassium bromide.


* Corresponding author. Tel.: +98 831 4276482; fax: +98 831 4276493; e-mail: hadibi@kums.ac.ir

Thus, a series of electron-rich aromatics such as substituted phenols and anilines containing even numbers of electron-withdrawing substituents was subjected to bromination in acetonitrile at ambient temperature to furnish the corresponding bromoarenes. Aromatic amines were also examined as there is a possibility that they might be oxidized by this system rather than undergoing substitution.²⁰

BTPPMS is a mild, efficient, stable, and cheap reagent, which is quite soluble in dichloromethane, chloroform, acetone, and acetonitrile but insoluble in non-polar solvents such as carbon tetrachloride, *n*-hexane, and diethyl ether. It is readily prepared by the dropwise addition of an aqueous solution of Oxone[®] to an aqueous solution of benzyltriphenylphosphonium chloride in a quantitative yield at room temperature and can be stored for months without losing its potency.¹⁸ The amount of HSO₅⁻ in this reagent has been determined by iodometric titration²¹ and the measurements are consistent with almost 99% by weight of active oxidizing agent.

We have reported a selective method for the bromination of phenols, affording bromophenols in good to high yields within short reaction periods. Our method is based on the in situ oxidation of potassium bromide using **BTPPMS** in acetonitrile. Simply by adding **BTPPMS** to a solution of phenol and potassium bromide in acetonitrile, rapid and selective bromination was achieved at room temperature. The products could be separated by a straightforward workup. The method has been applied successfully to a variety of phenols. Most of the reactions were regioselective and in those examples where both *ortho*- and *para*-substitution was possible, the *para*-substituted product was the only isomer isolated. *para*-Substituted aromatics were brominated in the *ortho*-position. Introduction of an electron-withdrawing group to the aromatic ring substantially decreased the rate of ring bromination while an electron donating group increased the rate (Scheme 1).

Initially, the bromination of 2,6-dimethylphenol **1g** as a model compound using potassium bromide and **BTPPMS** in various solvents was examined. The solvents examined were dichloromethane, chloroform, and acetonitrile. The reactions were carried out by stirring **1g** with **BTPPMS** and KBr (1:1) at room temperature. Dichloromethane and chloroform were inferior solvents compared to acetonitrile. Next, the effects of the amount of the oxidant **BTPPMS** and KBr were examined with **1g** (1 mmol) in acetonitrile. The optimum molar ratio of phenol to KBr to oxidant **BTPPMS**



G = H, Electron releasing or withdrawing substituent

Scheme 1. Oxidative bromination of phenols using **BTPPMS** and potassium bromide.

(1:1:1) was found to be ideal for complete conversion. The role of **BTPPMS** was confirmed by conducting a blank experiment where the formation of bromo compound was not observed even after 24 h. The products were isolated by filtering the reaction mixture and washing with acetonitrile, quenching the filtrate with aqueous sodium thiosulfate, drying over sodium sulfate and concentrating in vacuo. The residue was purified by column chromatography over silica gel or by crystallization in an appropriate solvent. The structure was confirmed by ¹H NMR, mass spectra, and melting point and elemental analysis for crystalline products. To evaluate the utility of this procedure for large scale bromination, a 10-fold scale reaction was carried out successfully for the bromination of 2-methoxynaphthalene **1c**, 1-bromo-2-methoxynaphthalene was obtained in 89% yield within 2 h²⁶ (Table 1).































BTPPMS was also used to brominate anilines at ambient temperature. Various activated and deactivated anilines were brominated upon simple admixing with **BTPPMS** and potassium bromide. In most cases, the optimum mole ratio of aniline to KBr to oxidant **BTPPMS** was found to be 1:1:1 (Table 1). *N,N*-Dimethylaniline **1v** was selectively brominated at the *para*-position within 1 h to give the corresponding 4-bromo-*N,N*-dimethylaniline in a 80% yield. 2-Chloroaniline **1s**, 4-chloroaniline **1t**, and 4-bromo-2-methylaniline **1u** exhibited a similar reactivity leading to the corresponding brominated anilines in good to high yields with no detectable influence of the substituent. Bromination of acetanilide **1w** which is less reactive than aniline proceeded well and produced 4-bromoacetanilide in 90% yield in 4 h (Table 1). It is interesting to mention that this reaction only requires a few milliliters of acetonitrile and the end point of reaction is easily confirmed by the decolorization of the dark-brown solution. Finally, we believe that the procedure for the bromination of aromatic amines using **BTPPMS** is a highly useful method owing to its ease, mildness, and reaction under nearly neutral reaction conditions (Scheme 2).

Bromination of an aromatic compound in the presence of **BTPPMS** proceeds according to the stoichiometry of Equation 1. It is believed that the bromination proceeds via the formation of potassium hypobromous. Potassium hypobromous has a higher instability due to pronounced ionic nature and thus more reactivity towards the aromatic nucleus.

The other mechanism is proposed according to a radical pathway upon hemolytic cleavage of the O–O bond in the peroxy monosulfate anion (O₃S–O–OH) according to Equation 2. The absence of bromination of the side chain such as the ring methyl group is indicative of the electrophilic mechanism of the reaction rather than a radical pathway (Table 1). In the case of Oxone[®], a similar mechanism has been proposed in the literature.^{17c}

In conclusion, we have developed a novel system using **BTPPMS** as an interesting alternative to liquid oxidants such as hydrogen peroxide, *tert*-butylhydroperoxide, and dimethylsulfoxide for the oxidative bromination of

Table 1. Oxidative bromination of electron-rich aromatic rings using potassium bromide as a bromine source in the presence of benzyltriphenylphosphonium peroxymonosulfate as oxidant under nearly neutral reaction conditions^a

Entry	Substrate	Product	Time (h)	Yield ^b (%)	Mp or bp (°C)/Torr (lit.)
1a			5	88	65–66 (66 ²²)
1b			2	92	84–85 (84 ²²)
1c			2	89	83–84 (85 ²²)
1d			2	85	63–64 (64 ²²)
1e			6	81	32–34 (33–34 ²²)
1f			6	80	49–50 (48–49 ²²)
1g			1.5	88	78–79 (79.5 ²²)
1h			2.5	90	232/760 (231/760 ²²)
1i			2	86	86–88 (87.5 ²²)
1j			3	80	55–56 (55–57 ²³)
1k			7	80	114–115 (114 ²²)
1l			5	84	91–93 (92 ²²)
1m			7	80	114–115 (114 ²²)
1n			1.5	90	86–87 (87 ²²)
1o			2	85	63–64 (64 ²²)


(continued on next page)

Table 1 (continued)

Entry	Substrate	Product	Time (h)	Yield ^b (%)	Mp or bp (°C)/Torr (lit.)
1p			1.5	80	101–103 (100–102 ²²)
1q			1.5	85	111–113 (111.5 ²⁴)
1r			5	80	168 (168–169 ²²)
1s			2	75	71–73 (73 ²²)
1t			2	77	64–66 (65–67 ²²)
1u			1.5	88	40–41.5 (39–41 ²²)
1v			1	80	55–56 (55 ²²)
1w			4	90	167–168 (168 ²²)
1x			5	87	108–110 (109–111 ²⁵)

^a Confirmed by comparison with authentic samples.


^b Yield of isolated pure product after purification.




G = H, Electron releasing or withdrawing substituent

X = NH₂, NHCOCH₃

Scheme 2. Oxidative bromination of anilines using **BTPPMS** and potassium bromide.



Equation 1. Proposed mechanism for the oxidative bromination through a possible electrophilic pathway.



Equation 2. Proposed mechanism for the oxidative bromination through a possible radical pathway.

aromatic substrates. Nearly neutral reaction conditions, no evolution of hydrogen bromide, moderate to high yields of monobrominated products, and simple work-up are advantages of the present method. It has also been demonstrated that the method is amenable to scale-up.

Yields refer to isolated products after purification. The products were characterized by comparing their spectral and physical data (TLC, MS, melting point, combustion elemental analysis and ¹H NMR) with those of authentic samples reported in the literature. The reagent

BTPPMS was prepared according to our previously reported procedures.^{18,19}

Acknowledgements

We gratefully acknowledge the Kermanshah University of Medical Sciences Research Council for the financial support. We are also thankful of the Isfahan University of Technology (IUT), Iran, for the financial support. Further financial support from the Center of Excellency in Chemistry Research (IUT) is gratefully acknowledged.

References and notes

- (a) Christophersen, C. *Acta Chem. Scand.* **1985**, 39B, 515; (b) Muathen, H. A. *J. Org. Chem.* **1992**, 57, 2740; (c) Butler, A.; Walker, J. V. *Chem. Rev.* **1993**, 93, 1937; (d) Katritzky, A. R.; Li, J.; Stevens, C. V.; Ager, D. J. *Org. Prep. Proced. Int.* **1994**, 26, 439; (e) *Ullmann's Encyclopedia of Industrial Chemistry*, 6th ed.; Wiley-VCH: Weinheim, 1998.
- Bora, U.; Bose, G.; Chaudhuri, M. K.; Dhar, S. S.; Gopinath, R.; Khan, A. T.; Patel, B. K. *Org. Lett.* **2000**, 2, 247.
- Schmid, H. *Helv. Chim. Acta* **1946**, 29, 1144.
- Duan, J.; Zhang, L. H.; Dolbier, W. R., Jr. *Synlett* **1999**, 1245.
- Bovonsombat, P.; McNelis, E. *Synthesis* **1993**, 237.
- Auerbach, J.; Weissman, S. A.; Blacklock, T. J.; Angeles, M. R.; Hoogsteen, K. *Tetrahedron Lett.* **1993**, 34, 931.
- Konishi, H.; Aritomi, K.; Okano, T.; Kiji, J. *Bull. Chem. Soc. Jpn.* **1989**, 62, 591.
- Ranu, B. C.; Sarkar, D. C.; Chakraborty, R. *Synth. Commun.* **1992**, 22, 1095.
- (a) Vega, F.; Sasson, Y.; Huddersman, K. *Zeolites* **1993**, 13, 341; (b) Smith, K.; Bahzad, D. *J. Chem. Soc., Chem. Commun.* **1996**, 467.
- Goldberg, Y.; Alper, H. *J. Mol. Catal.* **1994**, 88, 377.
- Paul, V.; Sudalai, A.; Daniel, T.; Srinivasan, K. V. *Tetrahedron Lett.* **1994**, 35, 7055.
- Clark, J. H.; Ross, J. C.; Macquarrie, D. J.; Barlow, S. J.; Bastock, T. W. *J. Chem. Soc., Chem. Commun.* **1997**, 1203.
- Barhate, N. B.; Gajare, A. S.; Wakharkar, R. D.; Bedekar, A. V. *Tetrahedron Lett.* **1998**, 39, 6349.
- (a) Majetich, G.; Hicks, R.; Reister, S. *J. Org. Chem.* **1997**, 62, 4321; (b) Srivastava, S. K.; Chauhan, P. M. S.; Bhaduri, A. *J. Chem. Soc., Chem. Commun.* **1996**, 2679.
- (a) Choudary, B. M.; Sudha, Y.; Reddy, P. N. *Synlett* **1994**, 450; (b) Hanson, J. R.; Opakunle, A.; Petit, P. *J. Chem. Res. (S)* **1995**, 457; (c) Bandgar, B. P.; Nigal, M. N. *J. Synth. Commun.* **1998**, 28, 3225.
- For recent work using Oxone[®] as an oxidant, see: (a) Hajipour, A. R. *Indian J. Chem.* **1997**, 36B, 1069; (b) Hajipour, A. R.; Mahboubkhah, N. *Org. Prep. Proced. Int.* **1999**, 31, 112; (c) Meunier, B. *New J. Chem.* **1992**, 16, 203.
- (a) Lee, K. J.; Cho, H. K.; Song, C. E. *Bull. Korean Chem. Soc.* **2002**, 23, 773; (b) Tamhankar, B. V.; Desai, U. V.; Mane, R. B.; Wadgaonkar, P. P.; Bedekar, A. V. *Synth. Commun.* **2001**, 31, 2021; (c) Narender, N.; Srinivasu, P.; Ramakrishna, M. P.; Kulkarni, S. J.; Raghavan, K. V. *Synth. Commun.* **2002**, 32, 2313.
- (a) Hajipour, A. R.; Mallakpour, S. E.; Adibi, H. *Phosphorus, Sulfur Silicon* **2000**, 167, 71; (b) Hajipour, A. R.; Mallakpour, S. E.; Adibi, H. *Chem. Lett.* **2000**, 460; (c) Hajipour, A. R.; Mallakpour, S. E.; Mohammadpoor-Baltork, I.; Adibi, H. *Phosphorus, Sulfur Silicon* **2000**, 165, 155; (d) Hajipour, A. R.; Mallakpour, S. E.; Mohammadpoor-Baltork, I.; Adibi, H. *Synth. Commun.* **2001**, 31, 1625; (e) Hajipour, A. R.; Mallakpour, S. E.; Mohammadpoor-Baltork, I.; Adibi, H. *Synth. Commun.* **2001**, 31, 3401; (f) Hajipour, A. R.; Mallakpour, S. E.; Mohammadpoor-Baltork, I.; Adibi, H. *Monatsh. Chem.* **2003**, 134, 45; (g) Hajipour, A. R.; Mallakpour, S. E.; Adibi, H. *Chem. Lett.* **2001**, 164; (h) Hajipour, A. R.; Mallakpour, S. E.; Adibi, H. *Phosphorus, Sulfur Silicon* **2002**, 177, 2277; (i) Hajipour, A. R.; Mallakpour, S. E.; Adibi, H. *J. Org. Chem.* **2002**, 67, 8666; (j) Hajipour, A. R.; Mallakpour, S. E.; Mohammadpoor-Baltork, I.; Adibi, H. *Molecules* **2002**, 7, 674; (k) Hajipour, A. R.; Mallakpour, S. E.; Mohammadpoor-Baltork, I.; Adibi, H. *Phosphorus, Sulfur Silicon* **2002**, 177, 2805.
- Hajipour, A. R.; Adibi, H. *J. Chem. Res. (S)* **2004**, 274.
- (a) Muzart, J. *Synthesis* **1995**, 1325; (b) McKillop, A.; Sanderson, W. R. *Tetrahedron* **1995**, 51, 6145.
- Trost, B. M.; Braslau, R. *J. Org. Chem.* **1988**, 53, 532.
- Dictionary of Organic Compounds*, 6th ed.; Buckingham, J., Donaghy, S. M., Eds.; Chapman and Hall: London, 1982.
- Hajipour, A. R.; Mallakpour, S. E.; Imanieh, H.; Pourmousavi, S. A. *J. Chem. Res. (S)* **2002**, 272.
- CRC Handbook of Chemistry and Physics*, 85th ed.; Lide, D. R., Ed.; CRC Press, 2004–2005.
- Crundwell, E. *J. Chem. Soc.* **1956**, 368.
- Representative procedure for the oxidative bromination of 2-methoxynaphthalene 1c*. To a solution of BTPPMS (10 mmol, 4.66 g) and potassium bromide (10 mmol, 1.19 g) in acetonitrile (50 mL) in a 250 mL round-bottomed flask, 2-methoxynaphthalene **1c** (10 mmol, 1.58 g) was added. The reaction mixture was stirred at room temperature for 2 h. After disappearance of the starting phenol as monitored by TLC using EtOAc/cyclohexane (2:8), the mixture was filtered through a sintered glass funnel and the solid residue was washed with acetonitrile (50 mL). The organic layers were combined and washed successively with 5% aqueous sodium thiosulfate solution (100 mL), dried over Na₂SO₄ and concentrated. The crude product was crystallized from ethanol to afford 1-bromo-2-methoxynaphthalene as solid needles in a 89% yield, mp 83–84 °C (lit.²² mp 85 °C).