

1,3-Dipolar cycloaddition of 2- and 3-nitroindoles with azomethine ylides. A new approach to pyrrolo[3,4-*b*]indoles

Sujata Roy,^a Tara L. S. Kishbaugh,^a Jerry P. Jasinski^b and Gordon W. Gribble^{a,*}

^aDepartment of Chemistry, Dartmouth College, Hanover, NH 03755, USA

^bDepartment of Chemistry, Keene State College, Keene, NH 03435, USA

Received 10 October 2006; revised 19 December 2006; accepted 20 December 2006

Available online 23 December 2006

Abstract—The 1,3-dipolar cycloaddition of unstabilized azomethine ylides with 2- and 3-nitroindoles furnishes the expected hexahydropyrrolo[3,4-*b*]indole cycloadducts in good to excellent yields. The cycloadducts can be denitrated with Bu₃SnH/AIBN, and cycloadduct **5** was oxidized with MnO₂ to yield the known pyrrolo[3,4-*b*]indole **13**.


© 2007 Elsevier Ltd. All rights reserved.

Although there are many examples of the indole π bond functioning as a dienophile in Diels–Alder reactions,¹ there are fewer examples of successful 1,3-dipolar cycloaddition reactions of indole.^{2–9} Apart from the spectacular carbonyl ylide applications to the synthesis of *Aspidosperma* alkaloids by Padwa^{6,7} and Boger,⁸ most examples of 1,3-dipolar cycloaddition reactions with the indole π bond afford either low-yielding mixtures or unstable products.^{2–5}

In our ongoing interest in the synthesis and chemistry of fused indoles,¹⁰ we previously reported that 1,3-dipolar cycloaddition reactions between 2- and 3-nitroindoles and mesoionic münchenones is an efficient one-step synthesis of pyrrolo[3,4-*b*]indoles,¹¹ which can be viewed as stable synthetic analogues of indole-2,3-quinodimethane. Although there are several routes to pyrrolo[3,4-*b*]indoles,¹² one obvious approach that has apparently not been described is the 1,3-dipolar cycloaddition between 2- and 3-nitroindoles and azomethine ylides. Indeed, the 1,3-dipolar cycloaddition of azomethine ylides with alkenes is a powerful reaction since it results in the formation of a pyrrolidine ring and has been widely used for the synthesis of innumerable nitrogen heterocycles and natural products.¹³

We now report our initial results on the 1,3-dipolar cycloaddition reaction between 2- and 3-nitroindoles and unstabilized azomethine ylides. We chose the α-amino acid decarboxylative route that was discovered independently by Joucla¹⁴ and Tsuge,¹⁵ and was based on the inaugural work by Rizzi,¹⁶ for the generation of azomethine ylides derived from amino acids and formaldehyde (Scheme 1). This extremely simple method utilizes commercially available compounds and is performed under almost neutral conditions. For example, the azomethine ylide derived from sarcosine and paraformaldehyde reacts with β-nitrostyrenes to give the corresponding pyrrolidines in good yield.¹⁷


Thus, treatment of 3-nitro-1-(phenylsulfonyl)indole (**1**)¹⁸ with the azomethine ylide generated in situ from sarcosine and paraformaldehyde in refluxing toluene affords the desired hexahydropyrroloindole cycloadduct **2** in 61% yield (Scheme 2).¹⁹ Although we somewhat anticipated the loss of nitrous acid from the initial cycloadduct **2** to furnish **3** as the final product, as we experienced in similar cases,¹¹ this path is not observed in any of our reactions, and the initially formed nitro cycloadducts **2** are quite stable. No reaction occurs



Scheme 1.

Keywords: Nitroindoles; Azomethine ylides; 1,3-Dipolar cycloaddition; Pyrrolo[3,4-*b*]indoles.

* Corresponding author. Tel.: +1 603 646 3118; fax: +1 603 646 3946; e-mail: GGriddle@dartmouth.edu



Scheme 2.

between the azomethine ylide from sarcosine and paraformaldehyde and 1-(phenylsulfonyl)indole,²⁰ 3-cyano-1-(phenylsulfonyl)indole,²¹ and 1-benzyl-3-nitroindole¹⁸ under the conditions that give **2**. Consistent with our earlier münchnone cycloadditions,¹¹ the presence of an electron-withdrawing protecting group on the indole nitrogen increases the dipolarophilic reactivity of the nitroindole toward the azomethine ylide. The reaction of sarcosine/paraformaldehyde with *tert*-butyl 3-nitroindole-1-carboxylate¹⁸ results in deprotection of the Boc group under the reaction conditions and no cycloadduct is isolated.

The crystal structure of **2** (Fig. 1) confirms the expected *cis*-addition of the azomethine ylide to 3-nitroindole **2**.²²

Similarly, upon reaction with 3-nitro-1-(phenylsulfonyl)indole (**1**) the azomethine ylide generated *in situ* from *N*-benzylglycine and paraformaldehyde in refluxing toluene gives cycloadduct **5** in almost quantitative yield (Scheme 3).²³ Likewise, 1-carbethoxy-3-nitroindole (**4**)¹⁸ furnishes **6** and **7** with the appropriate azomethine ylide under the same conditions.²⁴ In contrast, the reaction of **1** with glycine and paraformaldehyde in refluxing xylene or toluene does not furnish a cycloadduct. This lack of reactivity of glycine in these azomethine cycloadditions has precedence and may simply be due to the presence of a second acidic hydrogen on glycine that prevents generation of the azomethine ylide.²⁵ In general, we find that toluene is a better solvent than xylene for these cycloaddition reactions.

To investigate this 1,3-dipolar cycloaddition reaction with 2-nitroindoles, we treated 1-(phenylsulfonyl)-2-nitroindole (**8**)²⁶ with the azomethine ylides from both





Figure 1.




Scheme 3.


sarcosine and *N*-benzylglycine, and paraformaldehyde. To our satisfaction, the desired cycloadducts **9** and **10** were isolated in 86% and 67% yield, respectively (Scheme 4).²⁷ However, no cycloadduct is obtained in the reaction of 1,2-bis(phenylsulfonyl)indole²⁸ with these azomethines, again signifying the importance of the nitro group in these cycloaddition reactions, and perhaps also indicative of a steric effect with 1,2-bis(phenylsulfonyl)indole.

To access the pyrrolo[3,4-*b*]indole ring system, we needed to eradicate the nitro group from these cycloadducts. Although initial attempts with acid, base, or heat were unproductive, we found that treatment of **3** and **5** with Bu₃SnH²⁹ gives the denitrated products **11** and **12** in excellent yields (Scheme 5).³⁰ However, thus far, these conditions do not denitrate 2-nitroindoles.


As further structure confirmation, we treated hexahydropyrrolo[3,4-*b*]indole **12** with MnO₂ to afford pyr-



Scheme 4.



Scheme 5.



Scheme 6.

rolo[3,4-*b*]indole **13** in modest yield (**Scheme 6**), which was identical to a known sample.³¹ An attempt to oxidize **12** to **13** using DDO³² was unsuccessful.

In summary, the 1,3-dipolar cycloaddition of 2- and 3-nitroindoles with the unstabilized azomethine ylides generated *in situ* from the corresponding α -amino acids and paraformaldehyde in refluxing toluene affords hexahydropyrrolo[3,4-*b*]indoless in good to excellent yields. In one case, the cycloadduct could be oxidized to the corresponding pyrrolo[3,4-*b*]indole and this method offers a potential new route to these fused indole analogues of indole-2,3-quinodimethane. Our study of other 1,3-dipolar cycloadditions of nitroindoles and efforts to optimize this new route to pyrrolo[3,4-*b*]indoless are continuing and will be reported in due course.

Acknowledgements

This work was supported by the Donors of the Petroleum Research Fund (PRF), administered by the American Chemical Society, and by Wyeth. We thank Dr. Alison Rinderspacher for the preparation of **13** for comparison purposes.

References and notes

- Scheme 6.**

rolo[3,4-*b*]indole **13** in modest yield (**Scheme 6**), which was identical to a known sample.³¹ An attempt to oxidize **12** to **13** using DDQ³² was unsuccessful.

In summary, the 1,3-dipolar cycloaddition of 2- and 3-nitroindoles with the unstabilized azomethine ylides generated *in situ* from the corresponding α -amino acids and paraformaldehyde in refluxing toluene affords hexahydropyrrolo[3,4-*b*]indoless in good to excellent yields. In one case, the cycloadduct could be oxidized to the corresponding pyrrolo[3,4-*b*]indole and this method offers a potential new route to these fused indole analogues of indole-2,3-quinodimethane. Our study of other 1,3-dipolar cycloadditions of nitroindoles and efforts to optimize this new route to pyrrolo[3,4-*b*]indoless are continuing and will be reported in due course.

Acknowledgements

This work was supported by the Donors of the Petroleum Research Fund (PRF), administered by the American Chemical Society, and by Wyeth. We thank Dr. Alison Rinderspacher for the preparation of **13** for comparison purposes.

References and notes

 - For an excellent review, see: (a) Lee, L.; Snyder, J. K. *Adv. Cycloaddit. Chem.* **1999**, *6*, 119–171; (b) Kishbaugh, T. L. S.; Gribble, G. W. *Tetrahedron Lett.* **2001**, *42*, 4783–4785, and references cited therein.
 - Ozone: (a) Witkop, B.; Graser, G. *Ann. Chem.* **1944**, *556*, 103–114; (b) Mentzer, C.; Molho, D.; Berguer, Y. *Bull. Soc. Chim. Fr.* **1950**, *555*–561.
 - Azides: (a) Bailey, A. S.; Merer, J. J. *J. Chem. Soc. (C)* **1966**, *1345*–1348; (b) Bailey, A. S.; Chum, M. C.; Wedgwood, J. J. *Tetrahedron Lett.* **1968**, *9*, 5953–5954; (c) Bailey, A. S.; Scattergood, R.; Warr, W. A.; Cameron, T. S.; Prout, C. K.; Tickle, I. *Tetrahedron Lett.* **1970**, *11*, 2979–2982; (d) Bailey, A. S.; Warr, W. A.; Allison, G. B.; Prout, C. K. *J. Chem. Soc. (C)* **1970**, *956*–964; (e) Bailey, A. S.; Buckley, A. J.; Warr, W. A. *J. Chem. Soc., Perkin Trans. I* **1972**, *1626*–1629; (f) Bailey, A. S.; Buckley, A. J.; Warr, W. A.; Wedgwood, J. J. *J. Chem. Soc., Perkin Trans. I* **1972**, *2411*–2415; (g) Harmon, R. E.; Wellman, G.; Gupta, S. K. *J. Org. Chem.* **1973**, *38*, 11–16; (h) de la Mora, M. A.; Cuevas, E.; Muchowski, J. M.; Cruz-Almanza, R. *Tetrahedron Lett.* **2001**, *42*, 5351–5353; (i) He, P.; Zhu, S.-Z. *J. Fluorine Chem.* **2005**, *126*, 113–120; (j) He, P.; Zhu, S.-Z. *J. Fluorine Chem.* **2005**, *126*, 825–830.
 - Nitrilimines: (a) Ruccia, M.; Vivona, N.; Piozzi, F.; Aversa, M. C. *Gazz. Chim. Ital.* **1969**, *99*, 588–599; (b) Ruccia, M.; Vivona, N.; Cusmano, G.; Marino, M. L.; Piozzi, F. *Tetrahedron* **1973**, *29*, 3159–3164; (c) Laude, B.; Soufiaoui, M.; Arriau, J. *J. Heterocycl. Chem.* **1977**, *14*, 1183–1190.
 - Nitrile oxides: (a) Caramella, P.; Coda Corsico, A.; Corsaro, A.; Del Monte, D.; Albini, F. M. *Tetrahedron* **1982**, *38*, 173–182; (b) Bruché, L.; Zecchi, G. *J. Org. Chem.* **1983**, *48*, 2772–2773; (c) Dehaen, W.; Hassner, A. *J. Org. Chem.* **1991**, *56*, 896–900.
 - Isomünchnones: (a) Hertzog, D. L.; Austin, D. J.; Nadler, W. R.; Padwa, A. *Tetrahedron Lett.* **1992**, *33*, 4731–4734; (b) Padwa, A.; Hertzog, D. L.; Nadler, W. R. *J. Org. Chem.* **1994**, *59*, 7072–7084.
 - Carbonyl ylides: (a) Padwa, A.; Price, A. T. *J. Org. Chem.* **1995**, *60*, 6258–6259; (b) Padwa, A.; Price, A. T. *J. Org. Chem.* **1998**, *63*, 556–565; (c) Mejía-Oneto, J. M.; Padwa, A. *Org. Lett.* **2004**, *6*, 3241–3244; (d) Padwa, A.; Lynch, S. M.; Mejía-Oneto, J. M.; Zhang, H. *J. Org. Chem.* **2005**, *70*, 2206–2218; (e) Mejía-Oneto, J. M.; Padwa, A. *Org. Lett.* **2006**, *8*, 3275–3278.
 - Carbonyl ylides: (a) Wilkie, G. D.; Elliott, G. I.; Blagg, B. S. J.; Wolkenberg, S. E.; Soenen, D. R.; Miller, M. M.; Pollack, S.; Boger, D. L. *J. Am. Chem. Soc.* **2002**, *124*, 11292–11294; (b) Yuan, Z. Q.; Ishikawa, H.; Boger, D. L. *Org. Lett.* **2005**, *7*, 741–744; (c) Choi, Y.; Ishikawa, H.; Velický, J.; Elliott, G. I.; Miller, M. M.; Boger, D. L. *Org. Lett.* **2005**, *7*, 4539–4542; (d) Elliott, G. I.; Velický, J.; Ishikawa, H.; Li, Y.; Boger, D. L. *Angew. Chem., Int. Ed.* **2006**, *45*, 620–622; (e) Ishikawa, H.; Elliott, G. I.; Velický, J.; Choi, Y.; Boger, D. L. *J. Am. Chem. Soc.* **2006**, *128*, 10596–10612.
 - Carbonyl ylides: Muthusamy, S.; Gunanathan, C.; Babu, S. A. *Tetrahedron Lett.* **2001**, *42*, 523–526.
 - (a) Gribble, G. W. *Pure Appl. Chem.* **2003**, *75*, 1417–1432; (b) Gribble, G. W. et al. *Curr. Org. Synth.* **2005**, *9*, 1493–1519.
 - (a) Gribble, G. W.; Pelkey, E. T.; Switzer, F. L. *Synlett* **1998**, 1061–1062; (b) Gribble, G. W.; Pelkey, E. T.; Simon, W. M.; Trujillo, H. A. *Tetrahedron* **2000**, *56*, 10133–10140.
 - (a) Pindur, U.; Erfanian-Abdoust, H. *Chem. Rev.* **1989**, *89*, 1681–1689; (b) Sha, C.-K. *Adv. Nitrogen Heterocycl.* **1996**, *2*, 147–178.
 - (a) Eberbach, W. *Sci. Synth.* **2004**, *27*, 441–498; (b) Nájera, C.; Sansano, J. M. *Curr. Org. Chem.* **2003**, *7*, 1105–1150; (c) *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*; Padwa, A., Pearson, W. H., Eds.; Wiley: New York, NY, 2002; For some recent examples and leading references, see: (d) Bonini, B. F.; Boschi, F.; Franchini, M. C.; Fochi, M.; Fini, F.; Mazzanti, A.; Ricci, A. *Synlett* **2006**, 543; Llamas, T.; Arrayás, R. G.; Carretero, J. C. *Org. Lett.* **2006**, *8*, 1795; Garner, P.; Kaniskan, H. Ü.; Hu, J.; Youngs, W. J.; Panzner, M. *Org. Lett.* **2006**, *8*, 3647; Grigg, R.; Sarker, M. A. B. *Tetrahedron* **2006**, *62*, 10332.
 - (a) Joucla, M.; Mortier, J. *J. Chem. Soc., Chem. Commun.* **1985**, 1566–1567; (b) Joucla, M.; Mortier, J.; Hamelin, J. *Tetrahedron Lett.* **1985**, *26*, 2775–2778.
 - (a) Tsuge, O.; Kanemasa, S.; Ohe, M.; Takenaka, S. *Chem. Lett.* **1986**, 973–976; (b) Tsuge, O.; Kanemasa, S.; Ohe, M.; Takenaka, S. *Bull. Chem. Soc. Jpn.* **1987**, *60*, 4079–4089.
 - Rizzi, G. P. *J. Org. Chem.* **1970**, *35*, 2069–2072.
 - (a) Nyerges, M.; Baláz, L.; Kádas, I.; Bitter, I.; Kövesdi, I.; Toke, L. *Tetrahedron* **1995**, *51*, 6783–6788; (b) Nyerges, M.; Bitter, I.; Kádas, I.; Tóth, G.; Toke, L. *Tetrahedron* **1995**, *51*, 6783–6788.
 - Pelkey, E. T.; Gribble, G. W. *Synthesis* **1999**, 1117–1122.
 - Compound **2**: mp 153–155 °C (dec); ^1H NMR (CDCl_3) δ 7.80 (dd, 2H, $J = 8.5$, 1.2 Hz), 7.77 (dd, 1H, $J = 8.3$, 0.7 Hz), 7.58 (tt, 1H, $J = 7.4$, 1.2 Hz), 7.41–7.46 (m, 4H), 7.14 (dt, 1H, $J = 7.6$, 1.0 Hz), 5.30 (dd, 1H, $J = 6.7$,

- 3.8 Hz), 3.39–3.41 (m, 1H), 3.19–3.29 (m, 2H), 3.12 (dd, 1H, $J = 10.4$, 3.5 Hz), 2.35 (s, 3H); ^{13}C NMR (CDCl_3) δ 143.3, 136.5, 134.4, 133.8, 132.5, 132.1, 129.7, 129.2, 127.8, 127.7, 127.2, 125.2, 125.1, 124.7, 116.2, 115.7, 98.8, 69.5, 69.2, 41.4, 40.9; LRMS (EI): m/z 359 (M^+), 313, 286, 270 (100%), 258, 171, 141, 129; HRMS (EI): calcd for $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_4\text{S}$: 359.0940, found: 359.0950.
20. Conway, S. C.; Gribble, G. W. *Heterocycles* **1990**, *30*, 627–633.
21. Janosik, T.; Lipson, A. C.; Gribble, G. W. *Org. Prep. Proc. Int.* **2004**, *36*, 289–292.
22. Crystallographic data for the structure of **3** have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication number CCDC 632312. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK [fax: +44(0)-1223-336033 or e-mail: deposit@ccdc.cam.ac.uk]. Crystal data for **3**: Intensity data were collected on a Bruker SMART APEX II CCD area detector system equipped with a graphite monochromator and a CuK α fine-focus sealed tube ($\lambda = 1.54178 \text{ \AA}$) at 100(2) K, using the ϕ - ω scan technique to a maximum θ angle of 65.08° (0.85 Å resolution). $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_4\text{S}$, $M = 359.40$, triclinic, $a = 9.2250(3) \text{ \AA}$, $b = 9.2313(3) \text{ \AA}$, $c = 10.7440(4) \text{ \AA}$, $\alpha = 78.2570(10)^\circ$, $\beta = 79.8050(10)^\circ$, $\gamma = 65.7290(10)^\circ$, $V = 812.14(5) \text{ \AA}^3$, space group $P\bar{1}$ (no. 2), $Z = 2$, $d_{\text{calcd}} = 1.470 \text{ g/cm}^3$, 7544 reflections measured, 2597 reflections [$I > 2\sigma(I)$] were used in all calculations, $R = 0.0317$, $R_w = 0.0853$. Structure solution and refinement were performed by Bruker SHELXTL.
23. Representative procedure (5): A mixture of 1-(phenylsulfonyl)-3-nitroindole (**1**) (0.15 g, 0.5 mmol), paraformaldehyde (0.09 g, 3 mmol), and *N*-benzylglycine (0.21 g, 1.25 mmol) in dry toluene (9 mL) was refluxed under nitrogen for 5 h. An additional portion of the same quantities of paraformaldehyde and *N*-benzylglycine was added and the mixture was refluxed until almost all of the nitroindole disappeared by TLC (about another 4 h). Subsequently the reaction mixture was cooled and the solvent was removed by rotary evaporation. The residue was subjected to column chromatography (initially, hexanes–dichloromethane 1:1; then hexanes–ethyl acetate 2:1) to yield the desired product (0.21 g, 95%) as a white solid: mp 141–143 °C; ^1H NMR (CDCl_3) δ 7.82 (d, 2H, $J = 8.2$ Hz), 7.80 (d, 1H, $J = 8.5$ Hz), 7.57 (t, 1H, $J = 7.6$ Hz), 7.45 (q, 3H, $J = 7.8$ Hz), 7.39 (d, 1H, $J = 7.6$ Hz), 7.28–7.34 (m, 3H), 7.21 (d, 2H, $J = 6.7$ Hz), 7.14 (t, 1H, $J = 7.6$ Hz), 5.33 (t, 1H, $J = 5.6$ Hz), 3.64 (s, 2H), 3.53 (d, 1H, $J = 10.4$ Hz), 3.39 (dd, 1H, $J = 9.9$, 7.2 Hz), 3.09 (d, 1H, $J = 10.4$ Hz), 3.01 (dd, 1H, $J = 10.2$, 4.4 Hz); ^{13}C NMR (CDCl_3) δ 143.1, 136.8, 136.6, 134.0, 132.1, 129.4, 128.7, 128.6, 128.0, 127.7, 127.4, 125.0, 124.9, 115.8, 98.6, 69.0, 63.2, 61.7, 58.4; LRMS (EI): m/z 435 (M^+), 389, 286, 270, 247, 149, 133, 129, 91 (100%); HRMS (EI): calcd for $\text{C}_{23}\text{H}_{21}\text{N}_3\text{O}_4\text{S}$: 435.1253, found: 435.1258.
24. Compound **6**: oil; ^1H NMR (CDCl_3) δ 8.01 (d, 1H, $J = 7.6$ Hz), 7.48 (d, 1H, $J = 7.6$ Hz), 7.42–7.45 (m, 1H), 7.28–7.32 (m, 3H), 7.21 (d, 2H, $J = 6.7$ Hz), 7.10 (t, 1H, $J = 7.6$ Hz), 5.52 (s, 1H), 4.31–4.40 (m, 2H), 3.53–3.70 (m, 3H), 3.17–3.35 (m, 2H), 2.82–2.96 (m, 1H), 1.35–1.46 (m, 3H); ^{13}C NMR (CDCl_3) δ 152.1, 143.7, 137.0, 131.8, 128.7, 128.6, 127.6, 126.7, 125.8, 124.2, 123.6, 115.5, 98.8, 66.9, 63.8, 62.1, 61.2, 58.5, 14.7; LRMS (EI): m/z 367 (M^+), 321, 202 (100%), 130; HRMS (EI): calcd for $\text{C}_{20}\text{H}_{21}\text{N}_3\text{O}_4$: 367.1532, found: 367.1535.
- Compound **7**: amorphous solid; ^1H NMR (CDCl_3) δ 7.97 (br d, 1H, 7 Hz), 7.51 (d, 1H, 7 Hz), 7.41 (br, 1H), 7.09 (t, 1H, 7 Hz), 5.48 (s, 1H), 4.33 (br, 2H), 2.90–3.47 (m, 4H), 2.32 (s, 3H), 1.38 (br, 3H); HRMS (EI): calcd for $\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}_4$: 291.1219, found: 291.1218.
25. Harling, J. D.; Orlek, B. S. *Tetrahedron* **1998**, *54*, 14905–14912.
26. Roy, S.; Gribble, G. W. *Tetrahedron Lett.* **2005**, *46*, 1325–1328.
27. Compound **9**: mp 130–132 °C; ^1H NMR (CDCl_3) δ 8.02–8.03 (m, 2H), 7.61–7.64 (m, 1H), 7.52–7.55 (m, 2H), 7.19–7.20 (m, 2H), 7.14 (d, 1H, $J = 7.6$ Hz), 7.02–7.05 (m, 1H), 4.13–4.16 (m, 1H), 4.10 (d, 1H, $J = 11.6$ Hz), 3.40 (d, 1H, $J = 11.6$ Hz), 3.31 (d, 1H, $J = 8.5$ Hz), 2.56 (dd, 1H, $J = 9.0$, 5.9 Hz), 2.40 (s, 3H); ^{13}C NMR (CDCl_3) δ 141.6, 139.1, 134.1, 129.6, 129.1, 128.3, 127.5, 124.8, 124.2, 112.6, 111.0, 64.7, 62.4, 56.6, 41.0; LRMS (EI): m/z 359 (M^+), 313, 270 (100%), 206, 171, 129; HRMS (EI): calcd for $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_4\text{S}$: 359.0940, found: 359.0934.
- Compound **10**: mp 119–121 °C (dec); ^1H NMR (CDCl_3) δ 7.99–8.01 (m, 2H), 7.63–7.68 (m, 1H), 7.54 (t, 2H, $J = 7.7$ Hz), 7.29–7.40 (m, 5H), 7.20–7.24 (m, 2H), 7.13 (d, 1H, $J = 6.9$ Hz), 7.03–7.08 (m, 1H), 4.18–4.25 (m, 2H), 3.75 (s, 2H), 3.54 (d, 1H, $J = 11.4$ Hz), 3.35 (t, 1H, $J = 8.6$ Hz), 2.62–2.67 (m, 1H); ^{13}C NMR (CDCl_3) δ 141.7, 134.0, 129.6, 129.2, 128.9, 128.8, 127.8, 127.7, 124.8, 124.2, 112.7, 110.9, 62.5, 60.0, 58.9, 56.1; LRMS (EI): m/z 435 (M^+), 406, 389, 297, 270, 247, 206, 155, 129, 91 (100%); HRMS (EI): calcd for $\text{C}_{23}\text{H}_{21}\text{N}_3\text{O}_4\text{S}$: 435.1253, found: 435.1258.
28. Pelkey, E. T.; Barden, T. C.; Gribble, G. W. *Tetrahedron Lett.* **1999**, *40*, 7615–7619.
29. (a) Ono, N.; Kamimura, A.; Miyake, H.; Hamamoto, I.; Kaji, A. *J. Org. Chem.* **1985**, *50*, 3692–3698; (b) Voituriez, A.; Moulinas, J.; Kouklovsky, C.; Langlois, Y. *Synthesis* **2003**, 1419–1426.
30. Compound **11**: ^1H NMR (DMSO-d_6) δ 7.80 (dd, 2H, $J = 8.6$, 1.2 Hz), 7.65 (tt, 1H, $J = 7.5$, 1.2 Hz), 7.52–7.56 (m, 2H), 7.39 (d, 1H, $J = 7.6$ Hz), 7.17 (dt, 1H, $J = 7.5$, 0.7 Hz), 7.14 (d, 1H, $J = 7.3$ Hz), 6.98 (dt, 1H, $J = 7.5$, 1.0 Hz), 4.71–4.74 (m, 1H), 3.71 (t, 1H, $J = 8.1$ Hz), 3.36 (s, 2H), 3.12 (d, 1H, $J = 10.4$ Hz), 2.76 (d, 1H, $J = 8.9$ Hz), 2.45 (dd, 1H, $J = 10.2$, 5.6 Hz), 2.37 (dd, 1H, $J = 9.0$, 7.2 Hz), 2.18 (s, 3H); ^{13}C NMR (DMSO-d_6) δ 141.9, 137.0, 134.9, 133.7, 129.5, 127.8, 126.9, 125.1, 124.0, 113.4, 66.2, 64.5, 62.7, 44.9, 40.8; LRMS (EI): m/z 314, (M^+), 173, 130 (100%); HRMS (EI): calcd for $\text{C}_{17}\text{H}_{18}\text{N}_2\text{O}_2\text{S}$: 314.1089, found: 314.1091.
- Compound **12**: mp 114–117 °C; ^1H NMR (CDCl_3) δ 7.76–7.78 (m, 2H), 7.65 (d, 1H, $J = 7.9$ Hz), 7.54 (tt, 1H, $J = 7.5$, 1.2 Hz), 7.42 (t, 2H, $J = 7.8$ Hz), 7.22–7.32 (m, 6H), 7.02 (d, 2H, $J = 4.7$ Hz), 4.62–4.66 (m, 1H), 3.71 (m, 1H), 3.64 (s, 2H), 3.07–3.09 (m, 2H), 2.89 (m, 1H), 2.62 (dd, $J = 9.3$, 4.4 Hz); ^{13}C NMR (CDCl_3) δ 142.2, 137.8, 133.3, 129.2, 129.0, 128.5, 128.4, 127.4, 127.3, 124.9, 124.4, 115.0, 65.8, 62.0, 60.3, 59.3, 44.9; LRMS (EI): m/z 390 (M^+), 318, 293, 269, 223, 202, 168, 133 (100%); HRMS (EI): calcd for $\text{C}_{23}\text{H}_{22}\text{N}_2\text{O}_2\text{S}$: 390.1402, found: 390.1405.
31. Compound **13**: mp 154–156 °C (lit.¹⁴ 156–157 °C); ^1H NMR (acetone-d_6) δ 8.03 (d, 1H, $J = 8.2$ Hz), 7.81 (dd, 2H, $J = 8.6$, 1.2 Hz), 7.54–7.58 (m, 2H), 7.39–7.43 (m, 2H), 7.35–7.38 (m, 2H), 7.28–7.32 (m, 2H), 7.23–7.26 (m, 2H), 7.19 (dt, 1H, $J = 7.6$, 0.9 Hz), 7.14 (d, 1H, $J = 1.8$ Hz), 7.09 (d, 1H, $J = 1.8$ Hz), 5.34 (s, 2H); ^{13}C NMR (acetone-d_6) δ 134.9, 129.9, 129.6, 128.6, 128.0, 127.6, 125.5, 125.0, 121.3, 116.0, 110.9, 105.0, 54.9.
32. Kishbaugh, T. L. S.; Gribble, G. W. *Synth. Commun.* **2002**, *32*, 2003–2008.