

# A Zn(II) ion selective fluorescence sensor that is not affected by Cd(II)

Julius N. Ngwendson and Anamitro Banerjee\*

Department of Chemistry, University of North Dakota, Grand Forks, ND 58202-9024, USA

Received 27 July 2007; revised 8 August 2007; accepted 9 August 2007

Available online 15 August 2007

**Abstract**—Comparison of sensors sodium-2,6-diamino-(*N,N,N',N'*-tetraacetate)-4-methylanisole **1** and sodium-2,6-diamino-(*N,N,N'*-triacetate)-4-methylanisole **2** reveal that the loss of an acetyl group in **2** leads to a more selective Zn(II) induced fluorescence enhancement and shows no response to any other metal ions including Cd(II). Structural modifications and AM1 calculations indicate that the sensor uses the three acetyl groups and the 3° amino nitrogen for binding the metal ion. AM1 calculations imply a trigonal bipyramidal coordination for Zn(II) with the solvent molecule occupying one of the axial positions.


© 2007 Elsevier Ltd. All rights reserved.

Over 300 enzymes and proteins require zinc to carry out their biological functions.<sup>1</sup> For example, the zinc sites in protein domains play structural catalytic, and cocatalytic roles.<sup>2</sup> Zinc is essential for the regulation of DNA synthesis during the proliferation and differentiation of cells.<sup>3</sup> Zinc is also suspected to play a role in neurological disorders such as Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and epileptic seizures.<sup>4</sup> Environmental impact of Zn ion in sea water due to human activity is being studied.<sup>5</sup> Zinc ion selective fluorescence sensors play an important role in studying these processes.

Several fluorescence sensors for Zn(II) ion have been reported in the literature.<sup>5a,6</sup> Unfortunately, sensors that can selectively detect Zn(II) ion without showing any response to Cd(II) are extremely rare. Selective detection of Zn(II) in the presence of Cd(II) becomes important in environmental matrix when both the ions are likely to be present.<sup>5</sup> Recently, we reported sodium-2,6-diamino-(*N,N,N',N'*-tetraacetate)-4-methylanisole **1** (Chart 1) as a Zn(II) selective fluorescence sensor that can be synthesized in three simple steps.<sup>7</sup> When the sensor is excited at 300 nm, it has an emission maximum at 386 nm. In the presence of Zn(II) ion, the sensor shows a tenfold enhancement in emission intensity. This sensor also undergoes approximately fivefold increase in the fluorescence intensity in the presence of Cd(II).

**Keywords:** Zinc sensors; Fluorescence sensors.


\* Corresponding author. Tel.: +1 701 777 3941; fax: +1 701 777 2331; e-mail: abanerjee@chem.und.edu



**Chart 1.** Structure of compounds 1–4.

In this Letter, we present compound **2** as a Zn(II) ion selective sensor that itself has minimal fluorescence, but shows approximately twenty-fivefold enhancement of the fluorescence intensity in the presence of Zn(II), while showing no response to any other metal ions including Cd(II).

Semi-empirical AM1 calculations using SPARTAN 6 of **1**–Zn(II) complex indicates that only three acetate groups of the sensor are involved in binding the metal ion (Fig. 1). This was a surprising result given the geometric




**Figure 1.** Energy minimized structure of **1**–Zn(II) complex as predicted by AM1 calculations.

constraints of involving two groups in *meta* positions in binding the metal ion. The presence of the methoxy group between them also implies that the metal ion must be held above the aromatic ring. Also, only three of the acetyl groups interact with the metal, which lies more to the side that contributes two of the acetyl groups. It appears from the calculations that the Zn(II) ion is located roughly above the 3° amino nitrogen and is coplanar with the three carboxylate oxygen atoms. When a molecule of water was added, it occupied a position directly above the metal ion without affecting the positions of other atoms. Further addition of water molecules resulted in no change around the metal ion. The additional water molecules occupied positions close to the carboxylate C=O, implying hydrogen bonding. Based on this result, we propose distorted trigonal bipyramidal coordination for the metal ion with the carboxyl groups in the equatorial positions and the amino nitrogen and a solvent molecule occupying the two axial positions.

Based on the results of the AM1 calculations, we decided to investigate the effect of the number of acetyl groups on the sensor. We synthesized compounds **2** and **3** (Chart 1) by controlling the amount of ethyl bromoacetate using the same procedure as the one used for the synthesis of **1** (see Supplementary data for the detailed procedure). An alternative structure of **1**–Zn complex could involve one iminodiacetate group and the methoxy group. To test this possibility, we synthesized anisidine derivative **4**, which should be virtually indistinguishable from **1** as a sensor if indeed only one iminodiacetate group and the methoxy group were involved in binding the metal ion.

The fluorescence spectra for compounds **2**–**4** were measured in 0.1 M HEPES buffer at pH 7 in the presence of 0.135 M NaCl using Fluorolog-3 spectrofluorometer. Compound **4** has an emission maximum at 370 nm, which is quenched in the presence of Zn(II) ion (Supplementary data). This result shows that the binding behavior of **4** is different from sensor **1**. Based on this observation, we conclude that the geometry optimization of the AM1 calculations is at least partially correct in predicting the involvement of the acetyl groups from both the amino groups. Compound **3** did not fluoresce either in the presence or absence of the metal ions. While this does not rule out Zn(II) binding of **3**, this compound cannot act as a fluorescence sensor for the metal ion. The fluorescence spectrum of compound **2** ( $\lambda_{\text{ex}} = 300$  nm, pH 7) was very interesting (Fig. 2). As expected, similar to compound **1**, Co(II), Ni(II), and Cu(II) quenched the fluorescence of sensor **2**. Also, Ca(II) and Mg(II) had no effect on the fluorescence of the sensor. Surprisingly, Cd(II) enhanced the fluorescence intensity of the sensor marginally, while Zn(II) addition resulted in nearly twenty-fivefold enhancement of the fluorescence intensity. These results do not rule out the binding of the metal ions to the sensors, but the fluorescence enhancement is observed only for Zn(II). The fluorescence enhancement of the sensor in the presence of Zn(II) was independent of pH between 6 and 7.6. However, at higher pH the fluorescence intensity dropped due to the precipitation of Zn(OH)<sub>2</sub>. UV spec-




**Figure 2.** Fluorescence spectrum of **2** in the presence of various metal ions in 0.1 M HEPES buffer at pH 7 ( $\lambda_{\text{ex}} = 300$  nm).

trum of sensor **1** or **2** did not show any detectable change in the presence of the metal ions.

Sensors **1** and **2** act as sensors due to enhanced fluorescence intensity rather than a change in the emission wavelengths. The emission profile of the sensors as well as their metal bound complexes have the same emission profile. This implies that the fluorescence quantum yield of the sensor is increased only in the presence of Zn(II) ions. While the reason for this enhancement is still not clear, it is possible that Zn(II) binding provides rigidity to the molecule thereby blocking some of the nonradiative decay pathways leading to an enhanced fluorescence intensity.

Figure 3 compares the fluorescence intensity of sensors **1** and **2** in the presence of Zn(II) and Cd(II) metal ions. When excited at 300 nm in 0.1 M HEPES buffer at pH 7 in the presence of 0.135 M NaCl, both the sensors emit with approximately the same intensity. The emission maxima for **2** and its complexes with the metal ions are blue shifted by about 10 nm. While addition of Cd(II) marginally affects the emission spectrum of **2**, a fourfold enhancement of intensity was observed for **1**.




**Figure 3.** Comparison of the fluorescence spectra of **1** and **2** in the presence of Zn(II) and Cd(II) ions in 0.1 M HEPES buffer at pH 7 ( $\lambda_{\text{ex}} = 300$  nm).

Addition of Zn(II) to the sensor solutions led to nearly tenfold enhancement of the fluorescence intensity of **1**, whereas a twenty-fivefold enhancement was observed for **2**. Following the fluorescence intensity change at 378 nm during the titration of sensor **2** with increasing concentrations of Zn(II), a  $K_d$  value of  $8.5 \pm 2.5 \mu\text{M}$  was obtained following the best fit of the quadratic equation.<sup>8</sup> This value is comparable to that of sensor **1** ( $3.9 \pm 1.8 \mu\text{M}$ ),<sup>7</sup> indicating that three acetyl groups are involved in binding the metal ion as suggested by the AM1 calculations. The  $K_d$  value for binding of **2** with other metal ions could not be determined since the fluorescence change observed for these metals was negligible. The binding constants for **3** and **4** were not determined since they did not show any sensory activity.

In biological systems, Ca(II) are present in higher concentrations than Zn(II) ions.<sup>6c,9</sup> We incubated the sensor with 10 equiv of Ca(II) at room temperature. This resulted in a marginal enhancement of the fluorescence intensity of the sensor (Fig. 4). When 1 equiv of Zn(II) was added to the mixture, a fourfold enhancement of fluorescence intensity at 375 nm was observed. This result implies that there is at least a partial displacement of Ca(II) ions by Zn(II) ions even in the presence of a large excess of the Ca(II) ions. Similar displacements of other metal ions with Zn(II) ion resulting in an enhanced fluorescence intensity was observed for sensors **1** and **2**. Results of competitive binding experiments will be reported in the future.

Proton NMR of **1**–Zn complex shows a deshielding effect and peak broadening of all the protons relative to the free sensor.<sup>7</sup> A similar deshielding effect was observed for sensor **2** when bound to Zn(II) though it was much more prominent for aromatic protons. While the deshielding effect for **1** was about 0.05 ppm, the two aromatic protons of **2** were deshielded by about 0.44 and 0.64 ppm upon complexation with Zn(II). Moreover, the peak broadening effect that was observed for **1** was nonexistent in case of **2**–Zn complex. This implies that unlike **1**–Zn complex, **2**–Zn complex is not fluxional. AM1 calculations, NMR spectra, and binding constant



**Figure 4.** Fluorescence spectra of **2** in the presence of 10 equiv of Ca(II) ions before and after the addition of 1 equiv Zn(II) in 0.1 M HEPES buffer at pH 7 ( $\lambda_{\text{ex}} = 300 \text{ nm}$ ).

studies are consistent with a 1:1 sensor–Zn complex for both **1** and **2**. While, the possibility of the formation of symmetric polynuclear species of the type  $[\text{Sensor-Zn}]_x$ , where  $x = 2$  or higher cannot be ruled out, evidence so far points to a mononuclear complex at this time: only the fluorescence intensity is enhanced when the Zn(II) ion is added, without any shift in the emission or absorption spectra; and the HRMS (ESI) of the **2**–Zn complex gives a mass of 411.0122 (calculated: 411.0146). The HRMS data points toward a mononuclear species, however, the possibility of the observed peak being a daughter ion rather than a molecular ion peak cannot be ruled out. All attempts to recrystallize the sensor–Zn complexes in an attempt to elucidate its structure by X-ray diffraction resulted in the formation of a white powder.

Of the four compounds studied, only **1** and **2** can act as sensors. The number of acetyl group in these compounds play an important role in determining the sensitivity and selectivity of the metal ion. Our results indicate that two acetyl groups must be present on at least one of the nitrogen atoms in order for the compound to fluoresce (**1**, **2**, and **4** fluoresced, while **3** did not). Also, at least three acetyl groups must be present for the compound to act as a Zn(II) ion sensor. Theoretical studies and the results presented here suggests that all four acetyl groups do not participate in binding Zn(II) at the same time, thereby introducing fluxionality in the sensor–metal complex, which results in lower sensitivity and selectivity of the sensor. Three acetyl groups in **2** lead to a more rigid sensor–metal complex leading to higher sensitivity and selectivity without compromising the binding affinity. The results imply that three acetyl groups are more effective than four acetyl groups in selective sensing of Zn(II) ions. The selectivity of Zn(II) over Cd(II) and other metal ions makes compound **2** a rare and important class of fluorescent Zn(II) ion sensor.

### Acknowledgments

Authors thank the University of North Dakota for funding this research. The authors also thank Dr. D.K. Srivastava and Dr. David Pierce for helpful suggestions, Dr. Julia Zhao for the use of fluorimeter, and Dr. Alena Kubatova for the HRMS data (the funding for the spectrometer was provided by NSF CHE-0216038).

### Supplementary data

Experimental procedures, fluorescence spectrum of **4**, determination of  $K_d$  for **2**, and spectral data for all the materials. Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.tetlet.2007.08.040](https://doi.org/10.1016/j.tetlet.2007.08.040).

### References and notes

1. Coleman, J. E. *Annu. Rev. Biochem.* **1992**, *61*, 897–946.
2. Vallee, B. L.; Falchuk, K. H. *Physiol. Rev.* **1993**, *73*, 79–118.

3. Beyersmann, D.; Haase, H. *Biometals* **2001**, *14*, 331–341.
4. (a) Cuajungco, M. P.; Lees, G. J. *Brain Res.* **1997**, *23*, 219–236; (b) Takeda, A. *Biometals* **2001**, *14*, 343–351.
5. (a) Galceran, J.; Huidobro, C.; Companys, E.; Alberti, G. *Talanta* **2007**, *71*, 1795–1803; (b) Sandberg, J.; Wallinder, I. O.; Leygraf, C.; Bozec, N. L. *J. Electrochem. Soc.* **2007**, *154*, C120–C131.
6. (a) Parkesh, R.; Lee, T. C.; Gunnlaugsson, T. *Org. Biomol. Chem.* **2007**, *5*, 310–317; (b) Gong, H.-Y.; Zheng, Q.-Y.; Zhang, X.-H.; Wang, D.-X.; Wang, M.-X. *Org. Lett.* **2006**, *8*, 4895–4898; (c) Komatsu, K.; Kikuchi, K.; Kojima, H.; Urano, Y.; Nagano, T. *J. Am. Chem. Soc.* **2005**, *127*, 10197–10204; (d) Nolan, E. M.; Jaworski, J.; Okamoto, K.-I.; Hayashi, Y.; Sheng, M.; Lippard, S. J. *J. Am. Chem. Soc.* **2005**, *127*, 16812–16823; (e) Henary, M. M.; Wu, Y.; Fahrni, C. J. *Chem. Eur. J.* **2004**, *10*, 3015–3025; (f) Pearce, D. A. *Chem. Aust.* **2001**, *68*, 14–17; (g) Reany, O.; Gunnlaugsson, T.; Parker, D. *J. Chem. Soc., Perkin Trans. 2* **2000**, 1819–1831; (h) Nolan, E. M.; Lippard, S. J. *Inorg. Chem.* **2004**, *43*, 8310–8317.
7. Ngwendson, J. N.; Amiot, C. L.; Srivastava, D. K.; Banerjee, A. *Tetrahedron Lett.* **2006**, *47*, 2327–2330.
8. Qin, L.; Srivastava, D. K. *Biochemistry* **1998**, *37*, 3499–3508.
9. Voet, D.; Voet, J. G. *Biochemistry*, 2nd ed.; John Wiley & Sons, Inc.: New York, 1995.