



## Chemoselective three-component coupling via a tandem Pd-catalyzed boron-Heck and Suzuki reactions

Justin O'Neill, Kyung Soo Yoo, Kyung Woon Jung\*

Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, Los Angeles, CA 90089-1062, United States

### ARTICLE INFO

#### Article history:

Received 18 August 2008

Revised 7 October 2008

Accepted 8 October 2008

Available online 14 October 2008

### ABSTRACT


A new approach is herein reported to prepare biaryl derivatives via a tandem Pd-catalyzed boron-Heck and Suzuki reactions. This one-pot tandem process avoided purification or addition of extra catalyst between steps. The resulting biaryl compounds can be prepared with substrate variability in good to moderate yields.

© 2008 Elsevier Ltd. All rights reserved.



The biaryl moiety has been prevalent as a key structure of the functional molecules in raw materials for LCD (liquid crystal display) and OLED (organic light emitting diode) scaffolds, as well as for biologically active compounds.<sup>1</sup> Recently, the palladium-catalyzed reactions known as Heck–Mizoroki,<sup>2</sup> Suzuki–Miyaura,<sup>3</sup> Negishi,<sup>4</sup> and Stille<sup>5</sup> reactions have been employed as powerful and versatile methods for biaryl synthesis. These reactions have offered increased utility because of their mild conditions compared to traditional biaryl synthetic methods such as the Scholl,<sup>6</sup> Gomberg–Bachmann,<sup>7</sup> or Ullmann-type reactions.<sup>8</sup>

Recently, our group reported general methods for the intermolecular oxidative palladium-catalyzed boron-Heck cross coupling between aryl- or alkenylboronic acids and olefins under very mild conditions.<sup>9</sup> In particular, we examined several examples of competitive one-pot intermolecular Suzuki, Heck, and our boron-Heck cross coupling reactions. As shown in Scheme 1, we found that under oxidative conditions, the boron-Heck compound (**5**) was

formed exclusively. These results implied that under these coupling conditions, Heck and Suzuki coupling reactions were suppressed due to the absence of base, relatively low temperature, and presence of oxygen atmosphere. For the same competition reaction under nitrogen with sodium carbonate at 100 °C, there was almost completely selective formation of the Heck product (**4**) and no trace of the boron-Heck coupling product. This suggested that we could simply use different atmospheres to sequentially employ different palladium-catalyzed cross-coupling reactions in a three-component coupling tandem reaction. Therefore, we examined the oxidative palladium-catalyzed cross-coupling reactions of 4-iodophenylboronic acid with olefins such as *tert*-butyl acrylate (Eq. 1) and 2-cyclohexen-1-one (Eq. 2), because the resulting products would become suitable substrates for ensuing Suzuki couplings.



Scheme 1.



As expected, boron Heck-type products were exclusively generated, and no other coupling products were obtained. On the basis of these results, herein we report the development of a one-pot chemo- and regioselective method for the preparation of biaryl compounds via a tandem oxidative palladium-catalyzed boron-Heck and Suzuki coupling approach.

\* Corresponding author. Tel.: +1 213 740 8768; fax: +1 213 740 6270.  
E-mail address: [kwjung@usc.edu](mailto:kwjung@usc.edu) (K. W. Jung).

**Table 1**

Conditions for the selective oxidative boron-Heck reaction with acyclic alkene, *tert*-butyl acrylate **3**<sup>a</sup>



Entry	Mole ratio					Yield <sup>d</sup> (%)
	<b>7</b>	<b>3</b>	Pd(OAc) <sub>2</sub> (mol %)	T (°C)	Time (h)	
1 <sup>b</sup>	1	1.1	5	23	16	81
2 <sup>c</sup>	1	1.1	10	23	16	85
3 <sup>c</sup>	1	2	10	23	16	92
4 <sup>c</sup>	1	2	10	50	6	87

<sup>a</sup> All reactions were carried out with **7** (1.0 mmol) and various mole ratios of *tert*-butyl acrylate in DMF (2.5 mL).

<sup>b</sup> dmphen [2,9-Dimethyl phenanthroline] (5.5 mol %).

<sup>c</sup> dmphen (11.0 mol %).

<sup>d</sup> Isolated yields.

Initial studies focused on optimization of the boron-Heck type coupling of 4-iodophenylboronic acid (**7**) and olefins such as *tert*-butyl acrylate (acyclic example) and 2-cyclohexen-1-one (cyclic example). Table 1 shows the results from the screening of various temperatures, mole ratios, and catalyst/ligand loading amounts for this boron-Heck-type coupling using *tert*-butyl acrylate. The desired product (**8**) was formed exclusively in a high yield at room temperature using 10 mol % of Pd(II) catalyst and two equivalents of *tert*-butyl acrylate (entry 3). At 50 °C, the yield of product **8** was diminished due to generation of 4-iodophenol as a side product (entry 4), despite acceleration of the reaction rate.<sup>10</sup>

Next, experiments directed toward the development of a tandem oxidative boron-Heck/Suzuki reaction were performed using


4-iodophenylboronic acid, *tert*-butyl acrylate, and phenylboronic acid as coupling partners. As shown in Table 2, the oxidative boron-Heck reaction intermediate product **8** was prepared at room temperature, and then further reacted with phenylboronic acid in a Suzuki coupling in the presence of NaOH under a N<sub>2</sub> atmosphere at 50 °C. A tandem oxidative boron-Heck/Suzuki reaction using two equivalents of *tert*-butyl acrylate as an alkene formed the biaryl product (**11**) and a disubstituted acryl product (**12**) in a 2:1 ratio (entry 1), because the Heck reaction took place in the second-step reaction. Therefore, we changed the ratio of *tert*-butyl acrylate to a molar equivalent, and obtained more of the oxidative boron-Heck/Suzuki reaction product (entry 2). However, in the coupling reaction of disubstituted olefins such as ethyl crotonate and β-methylstyrene (entries 3 and 4), only tandem oxidative boron-Heck/Suzuki reaction products (**13** and **14**) were obtained in good yields, with excellent (*E*)-stereoselectivity.

Employing 2-cyclohexen-1-one **9** as an olefin, a similar screening process for the boron-Heck reaction was carried out with 4-iodophenylboronic acid, the results of which are summarized in Table 3. At room temperature, an increase in the amount of Pd(II) complex provided a rise in yield (entries 1 and 2). Increasing the molar ratio of 2-cyclohexen-1-one to 4-iodophenylboronic acid up to 2:1 increased the yield slightly (entry 3). Elevation of the temperature to 50 °C further increased the yield, while also accelerating the reaction rate (entry 4). However, lowering the ratio of 2-cyclohexen-1-one to 4-iodophenylboronic acid at this higher temperature resulted in significant decrease in yield (entry 5). Subsequently, we determined that optimal conditions for the cross-coupling reaction were to use two equivalents of 2-cyclohexen-1-one at 50 °C in the presence of 10 mol % of the Pd(II) catalyst.

Next, we examined the Suzuki reaction of intermediate compound (**10**) with phenylboronic acid. Thus, intermediate

**Table 2**

One-pot tandem reactions; oxidative cross-coupling of **7** with various olefins<sup>a</sup> and Suzuki reaction with phenylboronic acid<sup>b</sup>




Entry	Olefin	Mole equiv	Product <sup>c</sup>
1		2	
2		1	
3		2	
4		2	

<sup>a</sup> All reactions were carried out with **7** (1.0 mmol), olefins (2.0 mmol), Pd(OAc)<sub>2</sub> (10 mol %), and dmphen (11.0 mol %) in DMF (2.5 mL).

<sup>b</sup> Phenylboronic acid (1.2 mmol) and NaOH (2.0 mmol).

<sup>c</sup> Isolated yields.

**Table 3**Conditions for the selective oxidative boron-Heck reaction with 2-cyclohexen-1-one **9**<sup>a</sup>

Entry	Mole ratio					
	7	9	Pd/ligand (mol %)	T (°C)	Time (h)	Yield <sup>d</sup> (%)
1 <sup>b</sup>	1	1.2	5	23	16	61
2 <sup>c</sup>	1	1.2	10	23	16	78
3 <sup>c</sup>	1	2	10	23	16	80
4 <sup>c</sup>	1	2	10	50	6	85
5 <sup>c</sup>	1	1.2	10	50	6	65

<sup>a</sup> All reactions were carried out with **7** (1.0 mmol) and various mole ratios of 2-cyclohexen-1-one in DMF (2.5 mL).



<sup>b</sup> dmphen [2,9-Dimethyl phenanthroline] (5.5 mol %).

<sup>c</sup> dmphen [2,9-Dimethyl phenanthroline] (11.0 mol %).

<sup>d</sup> NMR yields.

compound **10** was generated from **7**, and to the crude reaction mixture containing **10**, phenylboronic acid was added and the reaction was stirred at 50 °C under N<sub>2</sub> in the presence of NaOH. After 6 h, the cross-coupling reaction proceeded to give a 45% yield of the desired biaryl product (**15**). Although this yield was in fact the overall yield for the two-step one-pot tandem reaction, this conversion was not satisfactory compared to the Suzuki reaction of intermediate **8** under similar conditions. Subsequently, we sought optimal conditions by screening various bases and temperatures to increase the Suzuki, and consequently, the overall reaction conversion. In this reaction, the use of carbonate and phosphate bases such as Na<sub>2</sub>CO<sub>3</sub>, K<sub>2</sub>CO<sub>3</sub>, and K<sub>3</sub>PO<sub>4</sub> at 50 °C provided the biaryl coupling product **15** in low yields. Also, the use of an organic base such as Et<sub>3</sub>N reduced the yield.<sup>11</sup> However, at the elevated temperature of 90 °C in the presence of NaOH, the reaction proceeded well to afford the coupling compound in an increased yield of 67% for the two-step tandem reaction (Scheme 2).

By utilizing these optimized Suzuki reaction conditions, we examined a tandem palladium-catalyzed oxidative boron-Heck and Suzuki reaction with 4-iodophenylboronic acid, 2-cyclohexen-1-one, and various arylboronic acids as summarized in Table 4. Reactions with electron-donating substituted arylboronic acids such as 4-methoxyphenyl boronic acid, 3,4-dimethoxyphenylboronic acid, and 4-*N,N*-dimethylaminophenylboronic acid took place smoothly to provide the desired compounds **16**, **17**, and **18** in 60%, 51%, and 59% yields, respectively (entries 2–4). Also, the reactions with 4-nitrophenylboronic acid, 4-acetylphenylboronic acid, and 4-cyanophenylboronic acid possessing highly electron-withdrawing groups afforded the biaryl products **19**, **20**, and **21** in 77%, 78%, and 63% yields, respectively (entries 5–7). The coupling reaction using 2,6-dimethyl phenylboronic acid gave a 30% yield of the desired product **22** due to steric hindrance (entry 8). In addition, we examined the Suzuki coupling reaction of the alkenylboronic acid, *trans*-styrenyl boronic acid, which proceeded to afford the corresponding cross-coupling compound **23** in 48% yield. Overall, these tandem reactions offered an efficient method for the preparation of biaryl compounds in a one-pot process.

**Scheme 2.****Table 4**One-pot tandem reactions, oxidative cross-coupling of **7** with 2-cyclohexen-1-one and Suzuki reaction with various arylboronic acids<sup>a,b</sup>

Entry	Arylboronic acid	Product	Yield <sup>c</sup> (%)
1			67
2			60
3			51
4			59
5			77
6			78
7			63
8			30
9			48

<sup>a</sup> All reactions were carried out with 4-iodophenylboronic acid (1.0 mmol), 2-cyclohexen-1-one (2.0 mmol), Pd(OAc)<sub>2</sub> (10 mol%), and dmphen (11 mol%) in DMF (2.5 mL).

<sup>b</sup> Arylboronic acid (1.2 mmol) and NaOH (2.0 mmol).

<sup>c</sup> Isolated yields.

To investigate the scope and limitation of the final coupling partner, we sought to use different halo-arylboronic acids as the centerpiece component for the tandem reaction. Several different halo-arylboronic acids were used with 2-cyclohexen-1-one and phenylboronic acid as coupling partners, using the established optimal reaction conditions. The results of these reactions are summarized in Table 5. A tandem reaction of 4-iodophenylboronic acid, 2-cyclohexen-1-one, and phenylboronic acid was converted

**Table 5**The effect of various halo-arylboronic acids with 2-cyclohexen-1-one **9**<sup>a,b</sup>

Entry	Arylboronic acid	Product	Yield <sup>c</sup> (%)
1			<5%
2			45
3			52
4			32
5			36

<sup>a</sup> All reactions were carried out with Arylboronic acid (1.0 mmol), 2-cyclohexen-1-one (2.0 mmol), Pd(OAc)<sub>2</sub> (10 mol%), and dmphen (11 mol%) in DMF (2.5 mL).

<sup>b</sup> Phenylboronic acid (1.2 mmol) and NaOH (2.0 mmol).

<sup>c</sup> Isolated yields.

efficiently to biaryl product **15** in 67% yield (Table 4, entry 1), while chloro- and bromo-phenylboronic acid afforded the desired product in 15% and 45% yields, respectively (entries 1 and 2). This is because aryl bromide and iodide easily react with phenylboronic acid but chloride does not participate in coupling readily. Furthermore, 3-iodophenylboronic acid reacted with 2-cyclohexen-1-one, and then phenylboronic acid to give a 52% yield of **24** (entry 3). In the case of di-substituted arylboronic acids including 3-chloro-4-methoxyphenyl boronic acid and 5-chloro-2-methoxyphenylboronic acid, tandem reactions with 2-cyclohexen-1-one and phenylboronic acid afforded regioselective desired products **25** and **26** in 32% and 36% yields, respectively (entries 4 and 5).

In conclusion, a tandem oxidative boron-Heck/Suzuki coupling reaction was developed for the preparation of biaryls in good to moderate yields. The reaction can be performed with variation at all three coupling partners. In addition, the biaryls formed in this study were obtained without the use of long laborious purification or the further addition of more palladium catalyst between coupling reactions. Furthermore, the use of different atmospheric conditions, and consequently, different mechanisms, allows for

the longevity of palladium catalysts and opens up possibilities for other tandem oxidative palladium reactions.

**Typical procedure for the tandem reaction:** To an oven dried 10 mL round-bottomed flask equipped with a stir bar was charged 0.10 mmol Pd(OAc)<sub>2</sub>, and 0.11 mmol 2,9-dimethyl phenanthroline in 2.5 mL DMF. The reaction was stirred at room temperature for 30 minutes, at which time 2.0 mmol alkene was added, followed by 1.0 mmol halo aryl boronic acid. The reaction flask was then fitted with an oxygen balloon and stirred at room temperature for 16 h, or at 50 °C for 6 h. Following confirmation of consumption of starting material by TLC, the oxygen balloon was removed and N<sub>2</sub> was bubbled through the solution for at least 1 min. 1.2 mmol aryl boronic acid and 2.0 mmol NaOH were then added to the solution, and the solution was then stirred for up to 6 h under N<sub>2</sub> at 90 °C. The reaction mixture was then dissolved in 50 mL ethyl acetate and washed twice with 50 mL water, once with 50 mL brine, and dried over anhydrous sodium sulfate and concentrated in vacuo. The crude reaction mixture was then subjected to column chromatography using a 10:1 hexane/ethyl acetate eluent system on silica gel (230–400 mesh).

### Acknowledgment

We acknowledge generous financial support from the National Institutes of General Medical Sciences of the National Institutes of Health (RO1 GM 71495).

### Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tetlet.2008.10.039.

### References and notes

- (a) Poetsch, E. *Kontakte* **1988**, 2, 15; (b) Bemis, G. W.; Murcko, M. A. *J. Med. Chem.* **1996**, 39, 2887; (c) Pu, L. *Chem. Rev.* **1998**, 98, 2405; (d) Hajduk, P. J.; Bures, M.; Praestgaard, J.; Fesik, S. W. *J. Med. Chem.* **2000**, 43, 3443; (e) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, 103, 893; (f) Croom, K. F.; Keating, G. M. *Am. J. Cardiovasc. Drugs* **2004**, 4, 395.
- (a) Hernandez, S.; SanMartin, R.; Tellitu, I.; Dominguez, E. *Org. Lett.* **2003**, 5, 1095; (b) Alberico, D.; Scott, M.; Lautens, M. *Chem. Rev.* **2007**, 107, 174.
- (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, 95, 2457; (b) Suzuki, A. *J. Organomet. Chem.* **1999**, 576, 147; (c) Kotha, S.; Lahiri, K.; Kashinath, D. *Tetrahedron* **2002**, 58, 9633; (d) Persichini, P. *J. Curr. Org. Chem.* **2003**, 7, 1725; (e) Bellina, F.; Carpita, A.; Rossi, R. *Synthesis* **2004**, 2419.
- (a) Negishi, E.; King, A. O.; Okukado, N. *J. Org. Chem.* **1977**, 42, 1821; (b) Rottaelder, M.; Palmer, N.; Knochel, P. *Synlett* **1996**, 573; (c) Tilley, J. W.; Clader, J. W.; Zawoiski, S.; Wirkus, M.; LeMahieu, R. A.; O'Donnell, M.; Crowley, H.; Welton, A. F. *J. Med. Chem.* **1989**, 32, 1814; (d) Hoyer, T. R.; Chen, M. *J. Org. Chem.* **1996**, 61, 7940.
- (a) Kosugi, M.; Hagiwara, I.; Migita, T. *Chem. Lett.* **1983**, 839; (b) Stille, J. K. *Angew. Chem., Int. Ed. Engl.* **1986**, 98, 504; (c) Stanforth, S. P. *Tetrahedron* **1998**, 54, 263.
- (a) Kovacic, P.; Jones, M. B. *Chem. Rev.* **1987**, 87, 357; (b) March, J. *Advanced Organic Chemistry*, 4th ed.; Wiley: New York, 1992; p 539.
- (a) Gomberg, M.; Bachmann, W. E. *J. Am. Chem. Soc.* **1924**, 46, 2339; (b) March, J. *Advanced Organic Chemistry*, 4th ed.; Wiley: New York, 1992; pp 715–171.
- (a) Ullman, F.; Bielecki, J. *Chem. Ber.* **1901**, 34, 2174; (b) Hassan, J.; Sevignon, M.; Gozzi, C.; Shulz, E.; Lemaire, M. *Chem. Rev.* **2002**, 102, 1359.
- (a) Jung, Y. C.; Mishra, R. K.; Yoon, C. H.; Jung, K. W. *Org. Lett.* **2003**, 5, 2231; (b) Yoo, K. S.; Yoon, C. H.; Mishra, R. K.; Jung, Y. C.; Yi, S. W.; Jung, K. W. *J. Am. Chem. Soc.* **2006**, 128, 16384.
- Moreno-Manas, M.; Perez, M.; Pleixats, R. *J. Org. Chem.* **1996**, 61, 2346.
- Conversion yields of the tandem reaction using various bases at 50 °C for 6 h under N<sub>2</sub> for the Suzuki coupling step; Na<sub>2</sub>CO<sub>3</sub> (29%), K<sub>2</sub>CO<sub>3</sub> (24%), K<sub>3</sub>PO<sub>4</sub> (37%), and triethyl-amine (25%).