



Highly efficient Pd-catalyzed synthesis of nitriles from aldoximes

Hoo Sook Kim, Sung Hwan Kim, Jae Nyoung Kim *

Department of Chemistry and Institute of Basic Science, Chonnam National University, Gwangju 500-757, Republic of Korea

ARTICLE INFO

Article history:

Received 26 December 2008

Revised 22 January 2009

Accepted 25 January 2009

Available online 5 February 2009

Keywords:

Palladium

Nitriles

Aldoximes

ABSTRACT

An expedient method of Pd-catalyzed conversion of aldoxime into nitrile was developed. The reaction was carried out under the influence of Pd(OAc)₂/PPh₃ in refluxing CH₃CN to provide good to high yields. The use of Cs₂CO₃ (0.1–0.5 equiv) was crucial in some cases.

© 2009 Elsevier Ltd. All rights reserved.

Palladium-catalyzed reactions of oxime derivatives can be classified into a few categories.^{1–5} Allylation of oxime can be carried out by the reaction with π -allylpalladium complex.¹ Deprotection of allyl oxime to oxime can also be catalyzed by Pd under Et₃N/HCOOH conditions.² Recently, Heck-type cyclizations of oxime ether have been reported.³ Various Pd-catalyzed *ortho*-functionalizations via C–H activation have been studied recently.⁴ However, conversion of oxime or its derivatives into nitrile with the aid of Pd catalyst has not been reported, to the best of our knowledge.⁶ Although many methods for the conversion of oxime or its derivative into nitrile has been reported,⁷ the synthesis of nitrile using transition metal catalyst is somewhat limited. The use of PtCl₄(EtCN)₂,⁶ [RuCl₂(*p*-cymene)]₂,^{7a,b} (HO)ReO₃,^{7c} Ga(OTf)₃,^{7d} W–Sn hydroxide^{7e}, and Cu(OAc)₂^{7f} has been reported.

During our recent studies on the Pd-catalyzed decarboxylative protonation reaction,⁸ we imagined the possibility of allyl group migration from oxygen to carbon as shown in Scheme 1. We chose *p*-nitrobenzaloxime allyl ether (**1**) as the model substrate based on the fact that *p*-nitro group can stabilize the Pd-intermediates as in recent papers of Tunge and co-workers.⁹ Initially formed π -allylpalladium complex (**I**) could be changed to (**II**), which might be converted into nitroso compound (**III**) and produced eventually allyl oxime **4**. However, we did not observe the formation of **4** at all. Instead, we isolated *p*-nitrobenzoxime (**2a**, 26%) and cinnamyl derivative **3** (8%),¹⁰ together with recovered starting material **1** (30%).

The mechanism for the formation of **2a** from **1** could be postulated as depicted in Scheme 1: (i) an oxidative addition of *O*-allyl

bond of **1** to Pd(0) to form the π -allylpalladium complex (**I**), (ii) deprotection to oxime by trace water in the reaction mixture,^{2,8} (iii) oxidative addition of *N*–*O* bond of oxime to Pd(0),³ and (iv) the following typical β -H elimination pathway to form nitrile **2a**. Actually, the reaction of *p*-nitrobenzaloxime ethyl ether did not produce any trace amounts of **2a** under the same conditions. The results state that Pd-catalyzed deprotection of allyl ether and the following Pd-catalyzed dehydration to nitrile could be the plausible mechanism.

The interesting results made us to investigate Pd-catalyzed generation of nitrile from aldoxime (Scheme 2), and we wish to report herein the preliminary successful results. We examined the reaction of *p*-nitrobenzaldehyde oxime (**5a**) and obtained *p*-nitrobenzoxime (**2a**) in moderate yield (67%) under the influence of Pd(OAc)₂ (10 mol %)/PPh₃ (20 mol %) in toluene at refluxing temperature for 24 h (entry 1 in Table 1). Fortunately, the yield was increased to 90% when we replace the solvent to CH₃CN within short time (3 h), as shown in Scheme 2 and in Table 1 (entry 2).¹¹ The conversion was completely ineffective without Pd catalyst (entry 3). The use of other Pd catalysts including PdCl₂ (entry 4) and Pd(PPh₃)₄ (entry 5), other phosphine ligands such as P(*o*-tolyl)₃ (entry 6) or PBu₃ (entry 7) did not improve the yield of **2a**. The reaction mechanism could be regarded as shown in Scheme 2, which involve the intermediate (**IV**). As in the Pd-catalyzed β -carbon elimination of cyclobutanone *O*-benzyloximes,^{5b} β -H elimination can explain the formation of nitrile.¹² As given in Table 1, the best yield was observed under the conditions comprising Pd(OAc)₂/PPh₃ in refluxing CH₃CN (Conditions A, entry 2). This is the first successful results on Pd-catalyzed dehydration of benzaloxime into nitrile.

Encouraged by the results we examined various arylaldoxime derivatives **5b–j** and cinnamyl oxime **5k**.¹¹ When the aromatic nu-

* Corresponding author. Tel.: +82 62 530 3381; fax: +82 62 530 3389.
E-mail address: kimjn@chonnam.ac.kr (J.N. Kim).





Table 1
Optimization of conditions for the conversion of **5a** to **2a** and **5e** to **2e**

Entry	Compound	Conditions	Yield (%)
1	5a	Pd(OAc) ₂ (10 mol %), PPh ₃ (20 mol %), toluene, reflux, 24 h	67
2 ^{a,b}	5a	Pd(OAc) ₂ (10 mol %), PPh ₃ (20 mol %), CH ₃ CN, reflux, 3 h	90
3	5a	no Pd catalyst, PPh ₃ (20 mol %), CH ₃ CN, reflux, 24 h	0
4	5a	PdCl ₂ (10 mol %), PPh ₃ (20 mol %), CH ₃ CN, reflux, 24 h	85
5	5a	Pd(PPh ₃) ₄ (10 mol %), PPh ₃ (20 mol %), CH ₃ CN, reflux, 16 h	86
6	5a	Pd(OAc) ₂ (10 mol %), P(o-tol) ₃ (20 mol %), CH ₃ CN, reflux, 3 h	83
7	5a	Pd(OAc) ₂ (10 mol %), PBu ₃ (20 mol %), CH ₃ CN, reflux, 18 h	82
8	5a	Pd(OAc) ₂ (10 mol %), PPh ₃ (20 mol %), Cs ₂ CO ₃ (0.1 equiv), CH ₃ CN, 80 °C, 1 h	87
9	5e	Pd(OAc) ₂ (10 mol %), PPh ₃ (20 mol %), CH ₃ CN, reflux, 4 h	49
10 ^c	5e	Pd(OAc) ₂ (10 mol %), PPh ₃ (20 mol %), Cs ₂ CO ₃ (0.1 equiv), CH ₃ CN, reflux, 1 h	83

^a Conditions A.

^b Without PPh₃ the reaction required longer time (8 h, 84%).

^c Conditions B.













cleus has a strong electron-withdrawing substituent, the yield of oxime was good to high (entries 1–4 in Table 2). However, 2,6-dichlorobenzaldoxime (**5e**) showed sluggish reactivity under the conditions (entry 9 in Table 1). Thus, we used Cs₂CO₃ (0.1 equiv) as in entry 10 in Table 1 (vide supra) and could improve the yield (Conditions B) although the role of Cs₂CO₃ is not clear at this stage. Thus, we used variable amounts of Cs₂CO₃ (0.1–0.5 equiv) for the substrates which showed sluggish reactivity under the conditions A such as **5e–h**, and **E-5j**. The results are summarized in Table 2.

The reaction of **5g** required long reaction time for the completion (entry 7). The reaction of cinnamyl oxime (**5k**) also showed same reactivity to produce **2k** in high yield (entry 12). It is interesting to note that both reactions of *E*- and *Z*-form of 2,4,6-trimethylbenzaldoxime (**5j**) produced nitrile **2j** in similar yields

(entries 10 and 11). From the results, we have to reconsider our proposed mechanism (vide supra, Scheme 2) in part. The possibility involving the Pd-catalyzed H₂O transfer from oxime to nitrile, as shown in Scheme 3, could not be excluded at this stage. This type of mechanism was suggested by Maffioli et al. very recently in their PdCl₂-catalyzed reversible dehydration of amides.¹³ Thus, the clarification of the final mechanism deserved further studies.^{14–16}

In summary, we disclosed the first successful Pd-catalyzed dehydration of aldoximes into nitriles. The plausible mechanism involved an oxidative addition of *N*-O bond to Pd(0) and the following typical β-H elimination pathway for most of the *E*-oximes. Irrespective of the mechanism as well as the configuration of oxime, Pd-catalyzed dehydration of aldoximes occurred very easily under mild conditions.

Table 2
Pd-catalyzed dehydration of aldoximes.


Entry	Aldoxime ^a	Conditions	Nitrile ^b (%)
1		A(3 h)	2a (90)
2		A(3 h)	2b (81)
3		A(2 h)	2c (87)
4		A(1 h)	2d (92)
5		B (1 h) ^c	2e (83)
6		B (8 h) ^d	2f (88)
7		B (80 h) ^d	2g (90)
8		B (8 h) ^d	2h (88)
9		A(7 h)	2i (87)
10		B(12 h) ^c	2j (87)
11		A(1 h)	2j (83)
12		A(3 h)	2k (91)

^a The configuration of oximes **5a–k** is *E*-form (>95%) except **Z-5j** (entry 11).

^b Isolated yield.

^c Cs₂CO₃ (0.1 equiv) was used.

^d Cs₂CO₃ (0.5 equiv) was used.



Scheme 3.

Acknowledgments

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, KRF-2008-313-C00487). Spectroscopic data was obtained from the Korea Basic Science Institute, Gwangju branch.

References and notes

- For the Pd-assisted allylation of oximes, see: (a) Miyabe, H.; Yoshida, K.; Reddy, V. K.; Matsumura, A.; Takemoto, Y. *J. Org. Chem.* **2005**, *70*, 5630–5635; (b) Miyabe, H.; Matsumura, A.; Yoshida, K.; Yamauchi, M.; Takemoto, Y. *Synlett* **2004**, 2123–2126.
- For deprotection of allyl group, see: Yamada, T.; Goto, K.; Mitsuda, Y.; Tsuji, J. *Tetrahedron Lett.* **1987**, *28*, 4557–4560.
- For Heck type cyclizations, see: (a) Ichikawa, J.; Nadano, R.; Ito, N. *Chem. Commun.* **2006**, 4425–4427; (b) Ohno, H.; Aso, A.; Kadoh, Y.; Fujii, N.; Tanaka, T. *Angew. Chem., Int. Ed.* **2007**, *46*, 6325–6328; (c) Zhu, J.-L.; Chan, Y.-H. *Synlett* **2008**, 1250–1254; For the Pd-catalyzed amino-Heck reactions of oxime derivatives, see: (d) Narasaka, K. *Pure Appl. Chem.* **2003**, *75*, 19–28; (e) Kitamura, M.; Zaman, S.; Narasaka, K. *Synlett* **2001**, 974–976; (f) Kitamura, M.; Narasaka, K. *Chem. Rec.* **2002**, *2*, 268–277. and further references cited therein.
- For C–H activations, see: (a) Desai, L. V.; Hull, K. L.; Sanford, M. S. *J. Am. Chem. Soc.* **2004**, *126*, 9542–9543; (b) Thu, H.-Y.; Yu, W.-Y.; Che, C.-M. *J. Am. Chem. Soc.* **2006**, *128*, 9048–9049; (c) Desai, L. V.; Malik, H. A.; Sanford, M. S. *Org. Lett.* **2006**, *8*, 1141–1144; (d) Thirunavukkarasu, V. S.; Parthasarathy, K.; Cheng, C.-H. *Angew. Chem., Int. Ed.* **2008**, *47*, 9462–9466.
- For the miscellaneous reactions of oximes, see: (a) Jiang, D.; Peng, J.; Chen, Y. *Org. Lett.* **2008**, *10*, 1695–1698; (b) Nishimura, T.; Nishiguchi, Y.; Maeda, Y.; Uemura, S. *J. Org. Chem.* **2004**, *69*, 5342–5347; (c) Grigg, R.; Markandu, J. *Tetrahedron Lett.* **1991**, *32*, 279–282; (d) Suzuki, O.; Hashiguchi, Y.; Inoue, S.; Sato, K. *Chem. Lett.* **1988**, 291–294; (e) Miyabe, H.; Yamaoka, Y.; Naito, T.; Takemoto, Y. *J. Org. Chem.* **2004**, *69*, 1415–1418.
- For ligand-mediated dehydration of aldoxime with nitrile complex trans-[PtCl₂(EtCN)₂], see: Makarycheva-Mikhailova, A. V.; Bokach, N. A.; Haukka, M.; Kukushkin, V. Y. *Inorg. Chim. Acta* **2003**, *356*, 382–386.
- For examples, on the synthesis of nitriles from oximes, see: (a) Yang, S. H.; Chang, S. *Org. Lett.* **2001**, *3*, 4209–4211; (b) Choi, E.; Lee, C.; Na, Y.; Chang, S. *Org. Lett.* **2002**, *4*, 2369–2371; (c) Ishihara, K.; Furuya, Y.; Yamamoto, H. *Angew. Chem., Int. Ed.* **2002**, *41*, 2983–2986; (d) Yan, P.; Batamack, P.; Prakash, G. K. S.; Olah, G. A. *Catal. Lett.* **2005**, *101*, 141–143; (e) Yamaguchi, K.; Fujiwara, H.; Ogasawara, Y.; Kotani, M.; Mizuno, N. *Angew. Chem., Int. Ed.* **2007**, *46*, 3922–3925; (f) Attanasi, O.; Palma, P.; Serra-Zanetti, F. *Synthesis* **1983**, 741–742; (g) Maeyama, K.; Kobayashi, M.; Kato, H.; Yonezawa, N. *Synth. Commun.* **2002**, *32*, 2519–2525; (h) Konwar, D.; Boruah, R. C.; Sandhu, J. S. *Tetrahedron Lett.* **1990**, *31*, 1063–1064; (i) Hart-Davis, J.; Battioni, P.; Boucher, J.-L.; Mansuy, D. *J. Am. Chem. Soc.* **1998**, *120*, 12524–12530; (j) Supana, P.; Liaskopoulos, T.; Tsoungas, P. G.; Varvounis, G. *Synlett* **2007**, 2671–2674. and further references cited therein.
- For our recent contribution on Pd-mediated decarboxylative protonation and allylation, see: Gowrisankar, S.; Kim, K. H.; Kim, S. H.; Kim, J. N. *Tetrahedron Lett.* **2008**, *49*, 6241–6244. and further references on Pd-mediated protonation and allylation reactions were cited therein.
- For the Pd-mediated decarboxylative allylation and related reactions involving nitro arene or pyridine moiety, see: (a) Waetzig, S. R.; Tunge, J. A. *J. Am. Chem. Soc.* **2007**, *129*, 14860–14861; (b) Waetzig, S. R.; Tunge, J. A. *J. Am. Chem. Soc.* **2007**, *129*, 4138–4139.
- Formation of cinnamyl compound **3** was very unusual. The result in CH₃CN was almost similar to that of the reaction in toluene. The phenyl moiety must be derived from PPh₃; however, detailed mechanism and scope of this novel finding needs further study. For the use of tetraarylphosphonium halides as arylating reagents in Pd-catalyzed Heck and cross-coupling reactions, see: Hwang, L. K.; Na, Y.; Lee, J.; Do, Y.; Chang, S. *Angew. Chem., Int. Ed.* **2005**, *44*, 6166–6169.
- Typical procedure for the conversion of *p*-nitrobenzaldehyde oxime (**5a**) into *p*-nitrobenzonitrile (**2a**): A stirred mixture of *p*-nitrobenzaldehyde oxime (83 mg, 0.5 mmol), Pd(OAc)₂ (11 mg, 10 mol %), and PPh₃ (26 mg, 20 mol %) in CH₃CN (1.5 mL) was heated to reflux for 3 h. The reaction mixture was filtered through a Celite pad, washed with EtOAc, and the solvent was removed. After column chromatographic purification process (hexanes/EtOAc, 8:1), analytically pure *p*-nitrobenzonitrile was isolated, 67 mg (90%) as a white solid.
- For the reactions involving L-Pd-OH intermediate, see: (a) Hosokawa, T.; Sugafuji, T.; Yamanaka, T.; Murahashi, S.-I. *J. Organomet. Chem.* **1994**, *470*, 253–255; (b) Kim, S. H.; Lee, H. S.; Kim, S. H.; Kim, J. N. *Tetrahedron Lett.* **2008**, *49*, 5863–5866.
- For the reversible dehydration of primary amides with PdCl₂ in aqueous acetonitrile, see: Maffioli, S. I.; Marzorati, E.; Marazzi, A. *Org. Lett.* **2005**, *7*, 5237–5239.
- For the possibility of anti-β-hydride elimination, see: Schwarz, I.; Braun, M. *Chem. Eur. J.* **1999**, *5*, 2300–2305. and further references cited therein.
- The reaction of acetophenone oxime under the same reaction conditions (CH₃CN, 0.1 equiv of Cs₂CO₃, reflux, 36 h) showed no reaction.
- During the evaluation process of this Letter, we carried out the reaction of phenylacetaldehyde oxime (*E/Z* = 5:4 mixture), a typical aliphatic oxime, and obtained benzyl cyanide in 56% under the same conditions given in Table 2 (conditions B with 0.5 equiv of Cs₂CO₃, 3 h).