



## Palladium-catalyzed carbonylative cyclization of 2-bromobenzaldehyde with primary amines leading to isoindolin-1-ones

Chan Sik Cho\*, Wen Xiu Ren

Department of Applied Chemistry, Kyungpook National University, Daegu 702-701, South Korea

### ARTICLE INFO

#### Article history:

Received 20 January 2009

Revised 13 February 2009

Accepted 16 February 2009

Available online 21 February 2009

#### Keywords:

2-Bromobenzaldehyde

Carbonylative cyclization

Isoindolin-1-ones

Palladium catalyst

Primary amines

### ABSTRACT


2-Bromobenzaldehyde is carbonylative cyclized with primary amines under carbon monoxide pressure in DMF at 100 °C in the presence of a catalytic amount of a palladium catalyst to give the corresponding isoindolin-1-ones in moderate to high yields.

© 2009 Elsevier Ltd. All rights reserved.

Palladium-catalyzed annulation provides a useful and convenient synthetic tool for the construction of skeletons of carbo- and heterocyclic compounds, which play an important role as a basic unit for the design of many pharmacologically and biologically active compounds.<sup>1,2</sup> Among them, palladium-catalyzed carbonylation with carbon monoxide followed by cyclization (carbonylative cyclization) has been recognized as an easy route for heterocyclic compounds containing carbonyl group such as lactones and lactams.<sup>3</sup> In connection with this report, several synthetic methods for isoindolin-1-ones have also been exemplified with such a palladium-catalyzed carbonylative cyclization.<sup>4</sup> For example, it is reported by us that 2-(2-bromophenyl)-2-oxazolines are carbonylative cyclized with aliphatic alcohols in the presence of a palladium catalyst to give tricyclic isoindolinones.<sup>5</sup> 2-Bromobenzaldehyde was also found to be coupled and cyclized with ethanalamines and ethylenediamines under carbon monoxide pressure in the presence of a palladium catalyst to afford tricyclic isoindolin-1-ones.<sup>6,7</sup> It is also reported that isoindolin-1-ones can be synthesized by the palladium-catalyzed three-component cyclative carbonylation–amination cascade.<sup>8</sup> It was disclosed by Miura that aryl iodides undergo a novel palladium-catalyzed cross-carbonylation reaction with *N*-benzylideneamines to give isoindolin-1-ones.<sup>9</sup> Under these circumstances, during the course of our ongoing studies on palladium-catalyzed cyclization reactions,<sup>10,11</sup> herein this report describes a palladium-catalyzed carbonylative cyclization of 2-bromobenzaldehyde with primary amines leading to isoindolin-1-ones.<sup>12,13</sup>

The results of several attempted carbonylative cyclizations of 2-bromobenzaldehyde (**1**) with aniline (**2a**) for the optimization of conditions are listed in Table 1. Treatment of **1** with equimolar amount of **2a** under carbon monoxide (10 atm) in DMF in the presence of PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> afforded *N*-phenylisoindolin-1-one (**3a**) in 25%

**Table 1**  
Optimization of conditions for the reaction of **1** with **2a**<sup>a</sup>



Run	[1]/[2a]	Palladium catalyst	Bases	Temp (°C)	Yield (%)
1	1.0	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	—	100	25
2	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	—	100	48
3	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	—	130	30
4 <sup>b</sup>	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	—	100	64
5 <sup>b</sup>	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	—	80	52
6 <sup>b</sup>	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	HCO <sub>2</sub> Na	100	66
7 <sup>b</sup>	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	Et <sub>3</sub> N	100	6
8 <sup>b</sup>	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	K <sub>2</sub> CO <sub>3</sub>	100	46
9 <sup>b</sup>	1.2	PdCl <sub>2</sub> (PPh <sub>3</sub> ) <sub>2</sub>	NaHCO <sub>3</sub>	100	27
10 <sup>b</sup>	1.2	PdCl <sub>2</sub> /2PPh <sub>3</sub>	—	100	73
11 <sup>b</sup>	1.2	Pd(OAc) <sub>2</sub> /2PPh <sub>3</sub>	—	100	58
12 <sup>b</sup>	1.2	PdCl <sub>2</sub> (PhCN) <sub>2</sub> /2PPh <sub>3</sub>	—	100	81

<sup>a</sup> Reaction conditions: **2a** (0.5 mmol), palladium catalyst (0.02 mmol), base (2 mmol), CO (10 atm), DMF (5 mL), for 24 h.

<sup>b</sup> DMF (10 mL).

\* Corresponding author. Tel.: +82 53 950 5586; fax: +82 53 950 6594.  
E-mail address: cscho@knu.ac.kr (C.S. Cho).


**Table 2**  
Palladium-catalyzed synthesis of isoindolin-1-ones **3**<sup>a</sup>

Run	Primary amines <b>2</b>	Isoindolin-1-ones <b>3</b>	Isolated yield (%)
1	<b>2a</b>	<b>3a</b>	81
2	<b>2b</b>	<b>3b</b>	86
3	<b>2c</b>	<b>3c</b>	57
4	<b>2d</b>	<b>3d</b>	80
5	<b>2e</b>	<b>3e</b>	81
6	<b>2f</b>	<b>3f</b>	41
7	<b>2g</b>	<b>3g</b>	77
8	<b>2h</b>	<b>3h</b>	72
9	<b>2i</b>	<b>3i</b>	76
10	<b>2j</b>	<b>3j</b>	30
11	<b>2k</b>	<b>3k</b>	78
12	<b>2l</b>	<b>3l</b>	54
13	<b>2m</b>	<b>3m</b>	61
14	<b>2n</b>	<b>3n</b>	49
15	<b>2o</b>	<b>3o</b>	54


<sup>a</sup> Reaction conditions: **1** (0.6 mmol), **2** (0.5 mmol), PdCl<sub>2</sub>(PhCN)<sub>2</sub> (0.02 mmol), DMF (10 mL), CO (10 atm), 100 °C, for 24 h.

isolated yield (run 1). The yield of **3a** was considerably affected by the molar ratio of **1** to **2a**, a slightly higher molar ratio, [1]/[2a] = 1.2, resulting in the effective formation of **3a** (run 2). Lower and higher reaction temperatures resulted in lower yields of **3a** (runs 3 and 5). Higher yield of **3a** was observed with dilution of the reaction mixture (run 4). It is known that sodium formate works as a hydride source in palladium-catalyzed hydrogenolysis of organohalides and intramolecular cyclization.<sup>14,15</sup> However, the reaction gave no significant change with further addition of sodium formate (run 6). Further addition of several bases such as Et<sub>3</sub>N, K<sub>2</sub>CO<sub>3</sub>, and NaHCO<sub>3</sub> to the reaction system rather resulted in a decreased yield of **3a** (runs 7–9). The catalytic systems using PdCl<sub>2</sub> and Pd(OAc)<sub>2</sub> combined with PPh<sub>3</sub> were revealed to be as effective as that using PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (runs 10 and 11). As a result, the best result was accomplished by the catalytic system using PdCl<sub>2</sub>(PhCN)<sub>2</sub> combined with PPh<sub>3</sub> as shown in run 12 of Table 1.

After the reaction conditions have been established, various aromatic and aliphatic primary amines **2** were subjected to the reaction with **1** in order to investigate the reaction scope, and several representative results are summarized in Table 2 (Scheme 1). With aromatic primary amines (**2a–k**), the corresponding isoindolin-1-ones (**3a–k**) were formed in the range of 30–86% isolated yields without any identifiable side products (runs 1–11). The product yield was not significantly affected by the electronic nature of the substituent on the aromatic ring of **2a–k**, whereas the position of that had a considerable relevance to the product yield. With *meta*-substituted aromatic primary amines (**2c**, **2f**, and **2j**),



**Scheme 1.**




**Scheme 2.**

the product yield is generally lower than those with primary aromatic amines having *ortho*- and *para*-substituents. Lower reaction rates and yields were observed with 1-aminonaphthalene (**2l**) and heteroaromatic primary amine **2m** when compared to those of primary aromatic amines having *ortho*- and *para*-substituents (runs 12 and 13). Aliphatic primary amines such as benzylamine (**2n**) and octylamine (**2o**) were also reacted with **1** to give *N*-benzylisoindolin-1-one (**3n**) and *N*-octylisoindolin-1-one (**3o**) in 49% and 54% yields, respectively (runs 14 and 15).

As to the reaction pathway, although it is not yet fully understood, this seems to proceed via an initial formation of imine **4** by the condensation between **1** and **2** (Scheme 2). Oxidative addition of a carbon–bromide bond of **4** to palladium(0) produces an arylpalladium(II) complex **5**, where carbon monoxide coordination to palladium and then aryl migration from palladium to the carbon of carbon monoxide occurs to give an arylpalladium(II) intermediate **6**. This is followed by an intramolecular acylpalladation to carbon–nitrogen double bond to give an alkylpalladium(II) intermediate **7**. Subsequent hydrogenolysis of **7** with molecular hydrogen gives isoindolin-1-one **3**. Molecular hydrogen seems to be produced by the reaction of DMF and/or CO with H<sub>2</sub>O generated in the initial condensation stage (Scheme 3). It is known that several transition metals catalyze hydrogen evolution from aqueous DMF solutions under relatively mild conditions.<sup>16,17</sup> The conversion of CO and H<sub>2</sub>O into CO<sub>2</sub> and H<sub>2</sub> is well known as water gas shift reaction.<sup>18</sup> We confirmed in a separate experiment that treatment of **1** with **2a** and further addition of D<sub>2</sub>O (0.5 mL) under the employed conditions afforded ca. 78% deuterated **3a** in 69% yield.<sup>19</sup> This result clearly indicates that H<sub>2</sub>O works as a hydrogen source in the hydrogenolysis of **7**.<sup>20,21</sup>

**General experimental procedure:** To a 50 mL stainless steel autoclave were added 2-bromobenzaldehyde **1** (0.6 mmol), primary amine **2** (0.5 mmol), PdCl<sub>2</sub>(PhCN)<sub>2</sub> (0.02 mmol), PPh<sub>3</sub> (0.04 mmol), and DMF (10 mL). After the system was flushed and then pressurized with carbon monoxide to 10 atm, the mixture was stirred at 100 °C for 24 h. The reaction mixture was passed through a short silica gel column (ethyl acetate–hexane) to eliminate inorganic salts. Removal of the solvent left a crude mixture, which was



**Scheme 3.**

separated by thin layer chromatography (silica gel, ethyl acetate–hexane mixture) to give isoindolin-1-ones **3**.<sup>22</sup>

In summary, we have shown that 2-bromobenzaldehyde undergoes carbonylative cyclization with aromatic and aliphatic primary amines under carbon monoxide pressure in the presence of a palladium catalyst to give isoindolin-1-ones in good yields. The present reaction is a straightforward methodology for the synthesis of isoindolin-1-ones from readily available starting compounds. The mechanistic rationale and further elaborated synthetic application for N-heterocycles using this protocol are currently under investigation.

### Acknowledgment

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2008-331-C00176).

### References and notes

1. Tsuji, J. *Palladium Reagents and Catalysis*; Wiley: Chichester, 1995.
2. *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; Wiley: New York, 2002.
3. Colquhoun, H. M.; Thompson, D. J.; Twigg, M. V. *Carbonylation: Direct Synthesis of Carbonyl Compounds*; Plenum Press: New York, 1991. p 173.
4. For physiological activities of isoindolin-1-ones, see: Takahashi, I.; Kawakami, T.; Hirano, E.; Yokota, H.; Kitajima, H. *Synlett* **1996**, 353. and references cited therein.
5. Cho, C. S.; Lee, J. W.; Lee, D. Y.; Shim, S. C.; Kim, T. J. *Chem. Commun.* **1996**, 2115.
6. Cho, C. S.; Chu, D. Y.; Lee, D. Y.; Shim, S. C.; Kim, T.-J.; Lim, W. T.; Heo, N. H. *Synth. Commun.* **1997**, 27, 4141.
7. Cho, C. S.; Jiang, L. H.; Shim, S. C. *Synth. Commun.* **1998**, 28, 849.
8. Grigg, R.; Zhang, L.; Collard, S.; Keep, A. *Tetrahedron Lett.* **2003**, 44, 6979.
9. Okuro, K.; Inokawa, N.; Miura, M.; Nomura, M. *J. Chem. Res. (S)* **1994**, 372.
10. (a) Cho, C. S.; Lim, D. K.; Zhang, J. Q.; Kim, T.-J.; Shim, S. C. *Tetrahedron Lett.* **2004**, 45, 5653; (b) Cho, C. S.; Patel, D. B.; Shim, S. C. *Tetrahedron* **2005**, 61, 9490.
11. (a) Cho, C. S.; Shim, H. S.; Choi, H.-J.; Kim, T.-J.; Shim, S. C.; Kim, M. C. *Tetrahedron Lett.* **2000**, 41, 3891; (b) Cho, C. S.; Lim, D. K.; Heo, N. H.; Kim, T.-J.; Shim, S. C. *Chem. Commun.* **2004**, 104; (c) Cho, C. S. *J. Organomet. Chem.* **2005**, 690, 4094; (d) Cho, C. S.; Patel, D. B. *Tetrahedron* **2006**, 62, 6388; (e) Cho, C. S.; Patel, D. B. *J. Mol. Catal. A: Chem.* **2006**, 260, 105; (f) Cho, C. S.; Kim, J. U. *Tetrahedron Lett.* **2007**, 48, 3775; (g) Cho, C. S.; Ren, W. X. *J. Organomet. Chem.* **2007**, 692, 4182.
12. For our reports on palladium-catalyzed carbonylative cyclizations, see: Cho, C. S.; Shim, H. S. *Tetrahedron Lett.* **2006**, 47, 3835. and references cited therein.
13. For 2-bromobenzaldehydes as a useful annulation counterpart, see: (a) Larock, R. C.; Doty, M. J. *J. Org. Chem.* **1993**, 58, 4579; (b) Gevorgyan, V.; Quan, L. G.; Yamamoto, Y. *Tetrahedron Lett.* **1999**, 40, 4089; (c) Dyker, G.; Grundt, P. *Tetrahedron Lett.* **1996**, 37, 619; (d) Terao, Y.; Satoh, T.; Miura, M.; Nomura, M. *Tetrahedron* **2000**, 56, 1315.
14. Pri-Bar, I.; Buchman, O. *J. Org. Chem.* **1986**, 51, 734.
15. Larock, R. C.; Babu, S. *Tetrahedron Lett.* **1987**, 28, 5291.
16. Schreiner, S.; Yu, J. Y.; Vaska, L. *J. Chem. Soc., Chem. Commun.* **1988**, 602.
17. Yu, J. Y.; Schreiner, S.; Vaska, L. *Inorg. Chim. Acta* **1990**, 170, 145.
18. Schumacher, N.; Boisen, A.; Dahl, S.; Gokhale, A. A.; Kandoi, S.; Grabow, L. C.; Dumesic, J. A.; Mavrikakis, M.; Chorkendorff, I. *J. Catal.* **2005**, 229, 265.
19. Deuterated **3a** was characterized with <sup>13</sup>C NMR spectrum [ $\delta$  50.58 (t, <sup>1</sup>J<sub>C-D</sub> = 21.8 Hz)] and the distribution was determined from the comparison of the peak area of a clearly separated signal with benzylic signal in <sup>1</sup>H NMR spectrum. An excess H<sub>2</sub>O seems to reduce the product yield by imine hydrolysis.
20. Cho, C. S.; Kim, J. U.; Choi, H.-J. *J. Organomet. Chem.* **2008**, 693, 3677.
21. A reviewer suggested that the transformation of **7** to **3** may also involve a hydridopalladium(II) **8** formation by the action of CO (on coordination sphere) and H<sub>2</sub>O without forming H<sub>2</sub>.
22. Selected spectroscopic data. Compound **3a**: Solid (hexane–chloroform); mp 162–163 °C; (lit.<sup>23</sup> 163–163.5 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  4.85 (s, 2H), 7.17 (t, *J* = 7.3 Hz, 1H), 7.40–7.44 (m, 2H), 7.48–7.52 (m, 2H), 7.57–7.61 (m, 1H), 7.85–7.88 (m, 2H), 7.92 (d, *J* = 7.6 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  51.15, 119.89, 123.04, 124.56, 124.90, 128.80, 129.58, 132.49, 133.64, 139.90, 140.54, 167.94. Compound **3b**: Solid (hexane–chloroform); mp 98 °C; (lit.<sup>23</sup> 98–98.5 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  2.26 (s, 3H), 4.72 (s, 2H), 7.22–7.34 (m, 4H), 7.51 (t, *J* = 7.6 Hz, 2H), 7.58–7.62 (m, 1H), 7.94 (d, *J* = 7.3 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  18.34, 53.15, 122.95, 124.33, 126.96, 127.55, 128.30, 128.38, 131.32, 131.80, 132.51, 136.48, 137.09, 141.69, 167.75. Compound **3n**: Solid (hexane–chloroform); mp 88–89 °C; (lit.<sup>24</sup> 87–89 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  4.25 (s, 2H), 4.79 (s, 2H), 7.26–7.38 (m, 6H), 7.43–7.52 (m, 2H), 7.89 (d, *J* = 7.1 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  46.48, 49.55, 122.89, 123.96, 127.79, 128.15, 128.26, 128.91, 131.49, 132.69, 137.11, 141.34, 168.62.
23. Yamamoto, I.; Tabo, Y.; Gotoh, H. *Tetrahedron Lett.* **1971**, 2295.
24. Wada, Y.; Nagasaki, H.; Tokuda, M.; Orito, K. *J. Org. Chem.* **2007**, 72, 2008.