

We have now determined that the reactivity of the PIDA/ammonium bromides is significantly enhanced in water, and the aqueous oxidation of an aliphatic alcohol **1a** to the corresponding ketone **2a** could smoothly occur at room temperature (Eq. 1). In this oxidation, the effect of water is remarkable, and the oxidation of **1a** did not effectively proceed in organic solvents.

Table 1
Aqueous oxidation with the combination of PIDA and Et₄N⁺Br^{-a}

Entry	Substrate	Product	Yield ^b (%), time (h)
1			76, 13
2			72, 4
3			90, 4
4			90, 6
5			89, ^c 5
6			81, 6
7			77, ^c 4
8			84, 4
9			78, 8 75, ^c 8
10			88, 3
11			75, 12
12 ^d			66, ^c 9

^a Reactions were performed using PIDA (1.2 equiv) and Et₄N⁺Br⁻ (1.0 equiv) in H₂O (0.1 M) at room temperature unless otherwise noted.

^b GC yield.

^c Isolated yield after purification.

^d 2.5 equiv of PIDA was used.


As the ammonium bromides, we chose Et₄N⁺Br⁻ because of the several following reasons: it is not only readily available and low-cost, but also easily removable from the product mixture after the reactions. Indeed, the separation of Et₄N⁺Br⁻ from the product **2a** could be easily attained by a simple workup (AcOEt or CH₂Cl₂ extraction),¹¹ taking advantages of the unique high partition coefficient of Et₄N⁺Br⁻ to water at ambient temperature.¹²

To confirm the scope of the aqueous system, we next examined the reactions using several types of alcohols, and the results are summarized in Table 1. The aliphatic secondary alcohols **1a–f** generally afforded the corresponding ketones **2a–f** in good to excellent yields (entries 1–6). The yields of **2c–f** were higher than the results in cyclohexanol **1a** and a sterically hindered alcohol **1b**. In an ester **1e**, no hydrolysis of **1e** and the product **2e** was observed (entry 5). Furthermore, benzyl alcohols **1g–k** also gave the aryl ketones **2g–k** in good yields (entries 7–11). Due to the mild reaction nature of the present system, a nitrogen atom containing substrate **1k** was applicable for the reactions (entry 11). Unfortunately, when we tried to perform the oxidation of primary alcohols under the same reaction conditions, the formation of small amounts of ester products was detected by GC analysis as a result of the oxidative condensation between two molecules of the starting primary alcohols.¹³ An intramolecular fashion of this type of condensation rapidly occurred in the diol **1l** to exclusively produce the corresponding five-membered lactone **2l** in a good yield (entry 12).

The important synthetic merit of the present method is featured in the case of the oxidation of an optically active alcohol **1m** (Scheme 2). The chiral alcohol **1m** is a challenging substrate, which produces an enolizable α -aryl ketone **2m** as a result of the oxidation. With the [PhIO]_n/Et₄N⁺Br⁻ combination,¹⁴ the oxidation of an alcohol **1m** could proceed in a good yield and notably, the optical activity of the ketone **2** was well maintained throughout the reaction. We have clarified the racemization-free evidence by confirming the no in situ H/D exchange of deuterium at the acidic α -position of the resulting α -aryl ketone **2m** based on ¹H NMR and GC–Mass analyses. In contrast, the use of KBr⁶ instead of Et₄N⁺Br⁻ was found to somewhat reduce the ee value of **2m** (85% yield and 73% ee), indicating that even the weak Lewis acidity of the potassium ion is enough to cause the undesired enolization process of the α -aryl ketone **2m** in water. The racemization-free oxidation of **1m** to **2m** was also difficult using the metal-catalyzed aqueous oxidation strategy such as Na₂WO₄/[(C₈H₁₇)₃NMe]⁺H₂O₄⁻ in 30% H₂O₂ aq.^{3a,3b}

In the present iodine(III)-mediated oxidation method, iodobenzene (PhI) was typically co-produced together with the desired oxidation products **2**. This forces the removal of the organic contaminant from the products **2** by tedious chromatographic workup. To make the purification step simpler, we therefore planned to apply a recyclable hypervalent iodine(III) reagent as a useful alternative to PIDA or [PhIO]_n. The use of 1,3,5,7-tetrakis[4-(diacetoxyiodo)phenyl]adamantane **3** (Fig. 1), a recyclable iodine(III) reagent developed by us,⁹ met this purpose, and ketones **2** were obtained by a simpler experimental operation (see below). Typically, the recyclable reagent **3** could afford the ketones **2** in better yields than PIDA, except for the alcohols **1c** and **1j** (Table 2).

In all the experiments listed in Table 2, the reagent **3** could easily be removed from the reaction mixtures by simple filtration,



Scheme 2. Racemization-free oxidation of an optically active alcohol **1m**.




Figure 1. Adamantane-type recyclable hypervalent iodine(III) reagent.

Table 2
Reactions using a recyclable iodine(III) reagent **3**^a

Entry	Substrate	Product	Yield (%), time (h)
1	1a	2a	99, 4
2	1c	2c	88, 18
3 ^b	1e	2e	99, 2
4	1g	2g	99, 4
5	1h	2h	91, 5
6	1j	2j	82, 3
7 ^{c,d}	1l	2l	97, 24

^a Reactions were performed using **3** (1.1 × 1/4 equiv) and Et₄N⁺Br⁻ (0.5 equiv) in H₂O (0.1 M) at room temperature. The reagent **3** was recovered as its reduced form **4** in over 95% yields by filtration.

^b 0.2 equiv of Et₄N⁺Br⁻ was used.

^c 2.0 × 1/4 equiv of **3** was used.

^d 1.0 equiv of Et₄N⁺Br⁻ was used.

utilizing the insolubility of the co-produced tetraiodide **4** in water and methanol. Thus, after completion of the reactions, saturated NaHCO₃ aq and methanol were added to the reaction mixtures. The resulting precipitates involving **4** were collected by filtration and the residues were washed several times with small portions of methanol. With this operation, the reagent **4** could be removed from the mixtures in at least 95% recovered yields, which could be reoxidized to the initial form **3** in nearly quantitative yields by the treatment with *m*-chloroperbenzoic acid (*m*CPBA).⁹ On the other hand, the crude products **2** could be obtained from the filtrates by extraction with AcOEt or CH₂Cl₂ and by evaporation of the solvents. Isolation of the pure **2** could be easily achieved by short column chromatography on silica gel.

Another merit of choosing Et₄N⁺Br⁻ instead of KBr⁶ is the fact that the former bromide source exhibits superior performance when using the recyclable reagent **3**. Et₄N⁺Br⁻ is not only an effective bromide source, but also works as a phase transfer catalyst permitting the high reactivity of the water-insoluble reagent **3** in water. For example, the oxidation of **1a** with the recyclable reagent **3** and KBr (0.5 equiv) required a longer reaction time to reach the high conversion of the alcohol **1a** (24 h, 76% yield) compared to that using the same amount of Et₄N⁺Br⁻ (Table 2, entry 1). This result also implies that the use of Et₄N⁺Br⁻ enables the utilization of a variety of alternative reagents, and will potentially contribute to the development of new and green aqueous-phase oxidations.

In summary, we have demonstrated the facile and green aqueous oxidation of alcohols **1** to ketones **2** using the combination of hypervalent iodine(III) reagents and Et₄N⁺Br⁻. The present system showed a broad generality of the substrates **1** including the aliphatic alcohols. The choice of Et₄N⁺Br⁻ is indispensable for the successful reaction progress and suppression of the undesired

racemization process of the enolizable α-aryl ketone product **2m**. In addition, effective utilization of the recyclable hypervalent iodine(III) reagent **3** is possible when using Et₄N⁺Br⁻ as the bromide source, enhancing the practicability of the aqueous method. Further investigations on the scope of the functional group compatibility and availability of chiral compounds will lead to application of the method to more complex molecules such as natural products.

Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research (S and A) and for Young Scientists (B), and a grant for Scientific Research on Priority Areas 'Advanced Molecular Transformations of Carbon Resources' from the Ministry of Education, Culture, Sports, Science, and Technology, Japan. T.D. also acknowledges support from the Industrial Technology Research Grant Program from the New Energy and Industrial Technology Development Organization (NEDO) of Japan. N.T. thanks research fellowship of J.S.P.S. for Young Scientists.

References and notes

- (a) *Comprehensive Organic Functional Group Transformations*; Katritzky, A. R., Meth-Cohn, O., Rees, C. W., Pattenden, G., Moody, C. J., Eds.; Elsevier Science: Oxford, 1995; Vols. 3 and 5, (b) March, J. *Advanced Organic Chemistry*, 4th ed.; John Wiley & Sons: New York, 1992. Chapter 19; (c) Caron, S.; Dugger, R. W.; Ruggeri, S. G.; Ragan, J. A.; Ripin, D. H. B. *Chem. Rev.* **2006**, *106*, 2943.
- (a) Lindstrom, U. M. *Chem. Rev.* **2002**, *102*, 2751; (b) Klijn, J. E.; Engberts, J. B. F. *N. Nature* **2005**, *435*, 746; (c) Li, C.-J.; Chen, L. *Chem. Soc. Rev.* **2006**, *35*, 68.
- Selected recent papers: (a) Sato, K.; Aoki, M.; Takagi, J.; Noyori, R. *J. Am. Chem. Soc.* **1997**, *119*, 12386; (b) Sato, K.; Aoki, M.; Takagi, J.; Zimmermann, K.; Noyori, R. *Bull. Chem. Soc. Jpn.* **1999**, *72*, 2287; (c) Brink, G.-J.; Arends, I. W. C. E.; Sheldon, R. A. *Science* **2000**, *287*, 1636; (d) Figiel, P. J.; Leskela, M.; Repo, T. *Adv. Synth. Catal.* **2007**, *349*, 1173; (e) Kon, Y.; Yazawa, H.; Usui, Y.; Sato, K. *Chem. Asian J.* **2008**, *3*, 1642; (f) Tervekar, V. N.; Jadhav, N. C. *Synth. Commun.* **2008**, *38*, 3107.
- Recent reviews, see: (a) Stang, P. J.; Zhdankin, V. V. *Chem. Rev.* **1996**, *96*, 1123; (b) Kita, Y.; Takada, T.; Tohma, H. *Pure Appl. Chem.* **1996**, *68*, 627; (c) Kirschning, A. *Eur. J. Org. Chem.* **1998**, 2267; (d) Zhdankin, V. V.; Stang, P. J. *Chem. Rev.* **2002**, *102*, 2523; (e) *Hypervalent Iodine Chemistry*; Wirth, T., Ed.; Springer: Berlin, Heidelberg, 2003; (f) Zhdankin, V. V.; Stang, P. J. *Chem. Rev.* **2008**, *108*, 5299.
- (a) Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277; (b) Frigerio, M.; Santagostino, M.; Sputore, S.; Palmisano, G. *J. Org. Chem.* **1995**, *60*, 7272; (c) Tohma, H.; Kita, Y. *Adv. Synth. Catal.* **2004**, *346*, 111.
- (a) Tohma, H.; Takizawa, S.; Maegawa, T.; Kita, Y. *Angew. Chem., Int. Ed.* **2000**, *39*, 1306; (b) Tohma, H.; Maegawa, T.; Kita, Y. *Adv. Synth. Catal.* **2002**, *344*, 328.
- (a) Qian, W.; Jin, E.; Bao, W.; Zhang, Y. *Tetrahedron* **2006**, *62*, 556; (b) Su, W.; Wang, H.; Xia, C.; Li, J.; Zhao, P. *Angew. Chem., Int. Ed.* **2003**, *42*, 1042.
- Mu, R.; Liu, Z.; Yang, Z.; Lui, Z.; Wu, L.; Lui, Z.-L. *Adv. Synth. Catal.* **2005**, *347*, 1333.
- (a) Tohma, H.; Maruyama, A.; Maeda, A.; Maegawa, T.; Dohi, T.; Shiro, M.; Morita, T.; Kita, Y. *Angew. Chem., Int. Ed.* **2004**, *43*, 3595; (b) Dohi, T.; Maruyama, A.; Yoshimura, M.; Morimoto, K.; Tohma, H.; Shiro, M.; Kita, Y. *Chem. Commun.* **2005**, 2205.
- Brunjes, M.; Sourkouni-Argirusi, G.; Kirschning, A. *Adv. Synth. Catal.* **2003**, *345*, 635.
- Typical experimental procedure in Table 1: To a suspension of **1e** (33.3 mg, 0.150 mmol) in H₂O (1.50 mL) were added PIDA (58.0 mg, 0.180 mmol) and Et₄N⁺Br⁻ (31.5 mg, 0.150 mmol). The mixture was then stirred for 5 h at room temperature. AcOEt and solid sodium thiosulfate were successively added to the reaction mixture. After stirring for 5 min, the organic layer was separated, dried with Na₂SO₄, and evaporated to remove the solvent. Purification of the residue by column chromatography on silica gel gave **2e** (29.5 mg, 0.134 mmol) in 89% yield.
- It is known that the concentration of Et₄N⁺Br⁻ in the same volume of CH₂Cl₂ and water is in the ratio of 1/10,000: Pradines, V.; Despoux, S.; Claparols, C.; Martins, N.; Micheau, J.-C.; Lavabre, D.; Pimienta, V. *J. Phys. Org. Chem.* **2006**, *19*, 350.
- Direct oxidative esterification of alcohols, see: Tohma, H.; Maegawa, T.; Kita, Y. *Synlett* **2003**, 723, and references cited therein.
- The use of PIDA caused partial racemization of **2m** (3 h, 85% yield and 83% ee), probably due to the in situ released acetic acid.