



## Enantioselective synthesis of $\beta$ -lactams via the IndaBox–Cu(II)-catalyzed Kinugasa reaction

Takao Saito\*, Tomohiro Kikuchi, Hiroaki Tanabe, Junichi Yahiro, Takashi Otani\*

Department of Chemistry, Faculty of Science, Tokyo University of Science, Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan

### ARTICLE INFO

#### Article history:

Received 20 April 2009

Revised 8 June 2009

Accepted 11 June 2009

Available online 13 June 2009

#### Keywords:

Kinugasa reaction

$\beta$ -Lactam

Bis(oxazoline)

Enantioselective catalysis

Nitrono cycloaddition

### ABSTRACT

The enantioselective Kinugasa reaction of nitrones with terminal alkynes in the presence of 20 mol % of IndaBox–Cu(OTf)<sub>2</sub> and di-*sec*-butylamine (1.5 equiv) produced  $\beta$ -lactams with the highest level of enantiomeric excesses among the catalytic enantioselective Kinugasa reactions reported so far.

© 2009 Published by Elsevier Ltd.

$\beta$ -Lactams<sup>1</sup> are an important class of compounds because of their utility as antibiotics such as penicillins, cephalosporins, monobactams, and carbapenems. In addition, chiral  $\beta$ -lactams also serve as versatile building blocks for stereocontrollable synthesis of complex organic compounds. Among a number of synthetic approaches to chiral  $\beta$ -lactams,<sup>1,2</sup> the chiral ligand–Cu(I or II)-catalyzed cycloaddition of nitrones with terminal alkynes (known as the enantioselective Kinugasa reaction) has recently received much attention.<sup>3</sup>

In 1972, Kinugasa and Hashimoto first reported the reaction between copper(I) phenylacetylide and nitrones in dry pyridine to produce  $\beta$ -lactams.<sup>4</sup> In 1995, Miura et al. described the first catalytic version of the Kinugasa reaction using bis-diphenylphosphinoalkane catalysts, and they also pioneered the enantioselective Kinugasa reaction using 10–100 mol % of CuI–bis(oxazoline) **1a–c** catalysts in the reaction of phenylacetylene (**7a**) with diphenylnitrono **8a** to give *cis*- $\beta$ -lactam (*cis*-**9a**) with 40–68% enantiomeric excesses (ees) and 30% diastereomeric excess (de) (Fig. 1).<sup>5</sup> Fu and coauthors reported the C<sub>2</sub>-symmetric bis(azaferrocene) **2**–CuCl-catalyzed Kinugasa reaction, in which they succeeded in achieving excellent *cis*:*trans* diastereoselectivity (71:29 to >95:5 *drs*) and enantioselectivity (67–93% ees).<sup>6</sup> Tang and co-workers demonstrated the pseudo C<sub>3</sub>-symmetric tris(oxazoline) **3**–Cu(II)-catalyzed highly enantioselective (45–85% ees) and diastereoselective (*cis*:*trans* = 97:3–67:33, 20:80–11:89) Kinugasa reaction.<sup>7</sup> They stated that the Cu(II) catalytic system is effective

even under air atmosphere and without rigorously dry reaction conditions. Shintani and Fu described the first examples of an intramolecular enantioselective Kinugasa reaction using a P,N-mixed ligand **4**–CuBr complex to attain 85–91% ees.<sup>8</sup> Guiry et al. also employed HETPHOX **5**–Cu(I) as a P,N-mixed ligand complex in the Kinugasa reaction; however, the enantioselectivities of the predominant *cis*- $\beta$ -lactams were only 4–48% ees with *cis*-selectivity (up to 88% de).<sup>9</sup> L-Proline-mediated Kinugasa reaction for synthesis of exomethylene  $\beta$ -lactams with up to 15% ee was reported by Basak et al.<sup>10</sup> Thus, only a very limited number of the enantioselective Kinugasa reactions have appeared so far,<sup>11</sup> and development of catalytic highly enantioselective Kinugasa reactions is still a challenging task.

It is of value to find a potential ability of a simple and readily available chiral ligand–metal complex inducing high enantioselectivity in the Kinugasa reaction. In this context, IndaBox **6** would be a candidate ligand for the chiral Lewis acid-catalyzed Kinugasa reaction.<sup>12</sup> IndaBox **6** has received increasing attention among C<sub>2</sub>-symmetric chiral bis(oxazoline) ligands.<sup>13</sup> We have previously found that the IndaBox **6a**–Cu(II) complex catalyst effectively induces high enantioselectivities in a hetero-Diels–Alder reaction of 1-thiabutadienes<sup>14</sup> and 1,3-dipolar cycloaddition of nitrones with 3-alkenyl-2-oxazolidinones.<sup>15</sup> These findings suggested to us that the IndaBox **6**–Cu(II)-catalytic system would be applicable to a Kinugasa reaction that must be carried out under Lewis base conditions.<sup>12</sup> We report here the catalytic, highly enantioselective Kinugasa reaction using IndaBox **6** as the homochiral ligand.

In initial studies, we selected the reaction of phenylacetylene (**7a**) with C,N-diphenylnitrono **8a** as the model reaction using

\* Corresponding authors. Fax: +81 03 5261 4631 (T.S.).

E-mail address: [tsaito@rs.kagu.tus.ac.jp](mailto:tsaito@rs.kagu.tus.ac.jp) (T. Saito).




Figure 1. Chiral ligands used in the Kinugasa reaction.

**6**-Cu(I or II) chiral catalyst (*Y* mol %) and an amine (*Z* equiv) under the conditions (solvent, temperature, and time) listed in Table 1 (Scheme 1). First, in order to evaluate the IndaBox ligands **6a–c**, the model reaction between **7a** and **8a** was carried out in the presence of 20 mol % **6a–c**-Cu(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O and 1.0 equiv dicyclohexylamine (Cy<sub>2</sub>NH) in dichloromethane at room temperature (20 °C) (entries 1–3).<sup>7</sup> β-Lactam **9a** was obtained (42%, 49%, and 51% yields) in cis:trans ratios of 85:15–86:14 with 78%, 67%, and 69% ees of the cis-isomer, respectively. Thus, IndaBox **6a** gave the best enantioselectivity. We next examined **6a**-Cu(I) salt complexes in the Kinugasa reaction (entries 4–7) because Fu et al. achieved high enantioselectivity (77% ee for **9a**) using the CuCl–**6a** complex.<sup>6</sup> Copper(I) halides gave high cis-selectivities (95:5–86:14) but moderate enantioselectivities (53–23% ees) (entries 4–6), with no β-lactam (**9a**) being formed with CuOTf (entry 7). The Cu(OTf)<sub>2</sub>-**6a** catalyst was found to be the best choice among the copper salts examined, giving 84:16 cis:trans selectivity and 82% ee (entry 8).

Next, screening of amines as a base was performed. Primary amines RNH<sub>2</sub> (R = *n*-Pr, Cy), secondary amines R<sub>2</sub>NH (R = Et, *i*-Pr, *n*-Bu, *n*-Oct, piperidine, and morpholine), and triethylamine showed low to moderate enantioselectivities (15–71% ees) and cis:trans selectivities (86–61:14–39). Gratifyingly, di-*sec*-butylamine (*s*-Bu<sub>2</sub>NH) gave high enantio- and diastereoselectivities

Table 1


Screening of the copper catalysts and amines and optimization of the reaction conditions

Entry	Ligand	Cu(I) or Cu(II)	<i>Y</i> (mol %)	Amine	<i>Z</i> (Equiv)	Solvent	Temp (°C)	Time (h)	Yield <sup>a</sup> (%)	Cis:trans <sup>b</sup>	Cis ee <sup>c</sup> (%)
1	<b>6a</b>	Cu(ClO <sub>4</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	14	42	85:15	78
2	<b>6b</b>	Cu(ClO <sub>4</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	21	49	85:15	67
3	<b>6c</b>	Cu(ClO <sub>4</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	22	51	86:14	69
4	<b>6a</b>	CuCl	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	17	58	95:5	53
5	<b>6a</b>	CuBr	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	6	55	92:8	35
6	<b>6a</b>	CuI	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	8	50	86:14	23
7	<b>6a</b>	CuOTf	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	8	0	—	—
8	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	Cy <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	15	32	84:16	82
9	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.0	CH <sub>2</sub> Cl <sub>2</sub>	20	21	40	84:16	82
10	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.5	CH <sub>2</sub> Cl <sub>2</sub>	20	14	32	84:16	85
11	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.5	PhMe	20	28	40	77:23	68
12	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.5	THF	20	16	60	78:22	75
13	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.5	MeCN	20	5	61	91:9	18
14	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.5	<i>i</i> PrOAc	20	14	39	84:16	87
15	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.5	<i>i</i> -PrOAc	40	1	30	83:17	83
16	<b>6a</b>	Cu(OTf) <sub>2</sub>	20	<i>s</i> -Bu <sub>2</sub> NH	1.5	<i>i</i> -PrOAc	5	38	47	85:15	90
17	<b>6a</b>	Cu(OTf) <sub>2</sub>	15	<i>s</i> -Bu <sub>2</sub> NH	1.5	<i>i</i> -PrOAc	5	46	47	82:18	88
18	<b>6a</b>	Cu(OTf) <sub>2</sub>	10	<i>s</i> -Bu <sub>2</sub> NH	1.5	<i>i</i> -PrOAc	5	65	41	84:16	85

<sup>a</sup> Total yield of cis- and trans-isomers.

<sup>b</sup> Determined by <sup>1</sup>H NMR.

<sup>c</sup> Determined by chiral HPLC.



Scheme 1.

Table 2

Catalytic enantioselective Kinugasa reaction: variation of the nitrone *N*-aryl substituent R

Entry	<b>8</b>	R	Time (h)	Adduct	Yield <sup>a</sup> (%)	Cis:trans <sup>b</sup>	Cis ee <sup>c</sup> (%)
1	<b>8a</b>	H	38	<b>9a</b>	47	85:15	90
2	<b>8b</b>	Me	136	<b>9b</b>	41	81:19	89
3	<b>8c</b>	MeO	135	<b>9c</b>	40	79:21	93
4	<b>8d</b>	Cl	106	<b>9d</b>	49	86:14	79
5	<b>8e</b>	EtO <sub>2</sub> C	138	<b>9e</b>	72	83:17	81

<sup>a</sup> Total yield of cis- and trans-isomers.


<sup>b</sup> Determined by <sup>1</sup>H NMR.

<sup>c</sup> Determined by chiral HPLC.

(82% ee and cis:trans 84:16, entry 9), which are comparable with the results (Cy<sub>2</sub>NH) in entry 8. When the reaction using 1.5 equiv amounts of *s*-Bu<sub>2</sub>NH was carried out, a high level of enantioselectivity (85% ee) and diastereoselectivity (cis:trans 84:16) was obtained (entry 10), but the use of an excess amount (2.0 equiv) of the amine resulted in reduction of both yield and enantioselectivity (21%, 69% ee).


Optimization of the solvents (entries 10–14) led us to find somewhat higher enantioselectivity (87% ee) with good diastereoselectivity (cis:trans 84:16) when using isopropyl acetate (entry 14). The reaction at a higher temperature (40 °C) shortened the reaction time (to 1 h) with slight reduction of both the stereoselectivities (83% ee, cis:trans 83:17) (entry 15). Finally, when the reaction at 5 °C in *i*-PrOAc was performed using a 20 mol % Cu(OTf)<sub>2</sub>-**6a** catalyst and 1.5 equiv *s*-Bu<sub>2</sub>NH, the highest enantioselectivity (90% ee) of cis-isomer **9a** was attained in a cis:trans ratio of 85:15 (entry 16). The amounts of the catalyst could be reduced to 15 mol % or 10 mol % while maintaining a good enantiomeric excess of 88% ee (entry 17) or 85% ee (entry 18), respectively.

Under the optimized reaction conditions (Table 1, entry 16), the reaction of phenylacetylene (**7a**) with nitrones **8** with variation of

**Scheme 2.****Table 3**

Catalytic enantioselective Kinugasa reaction: variation of the nitron C-aryl substituent R

Entry	<b>8</b>	R	Time (h)	Adduct	Yield <sup>a</sup> (%)	Cis:trans <sup>b</sup>	Cis ee <sup>c</sup> (%)
1	<b>8a</b>	H	38	<b>9a</b>	47	85:15	90
2	<b>8f</b>	Me	48	<b>9f</b>	50	80:20	92
3	<b>8g</b>	MeO	40	<b>9g</b>	64	84:16	87
4	<b>8h</b>	Br	48	<b>9h</b>	45	87:13	81
5	<b>8i</b>	CF <sub>3</sub>	48	<b>9i</b>	41	84:16	92
6	<b>8j</b>	NO <sub>2</sub>	37	<b>9j</b>	47	84:16	94

<sup>a</sup> Total yield of cis- and trans-isomers.<sup>b</sup> Determined by <sup>1</sup>H NMR.<sup>c</sup> Determined by chiral HPLC.**Scheme 3.**

*N*-aryl substituents R was examined (Scheme 2). The results are shown in Table 2. The electron-withdrawing *N*-aryl substituent (R = Cl, EtO<sub>2</sub>C) lowered the enantioselectivity to 79% and 81% ee (entries 4 and 5) compared with the phenyl and *p*-tolyl groups (R = H, Me entry 1 and 2, 90% and 89% ees), whereas the elec-

tron-donating *p*-MeOC<sub>6</sub>H<sub>4</sub> substituent showed the highest 93% ee (entry 3). Because the reaction of **7a** with nitrones **8** bearing an alkyl substituent such as methyl or benzyl group failed, an *N*-aryl substituent seems to be necessary for the reaction, which is compatible with Fu's and Tang's results.<sup>6,7</sup>

The nitron C-aryl substituent effects were also examined (Scheme 3, Table 3). In these cases, the strongly electron-withdrawing substituents (R = CF<sub>3</sub>, NO<sub>2</sub>) showed the highest level of enantioselectivity (entries 5 and 6, 92% ee and 94% ee), in contrast to slight lowering by the *p*-MeO group (87% ee, entry 3). The *p*-bromo group was as low as 81% ee.

Finally, the scope with respect to the alkyne component **7** was probed (Scheme 4, Table 4). The electron-donating *p*-tolyl and *p*-MeOC<sub>6</sub>H<sub>4</sub> substituents showed the highest level of enantioselectivity (93% and 94% ees, entries 2 and 3). Although, in the case of the *p*-BrC<sub>6</sub>H<sub>4</sub> and *p*-CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub> substituents, the cis-isomer was the major product (76:24 and 72:28, entries 4 and 5), the much more electron-withdrawing substituent (R = *p*-NO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>) clearly affects the cis:trans diastereoselectivity (18:82, entry 6). In contrast, trans-isomer was formed as the major β-lactam while maintaining a high level of enantioselectivity. Because of the fact that separated *cis*-**9o** with 86% ee was completely transformed into *trans*-**9o** maintaining 86% ee after addition of an amine (*s*-Bu<sub>2</sub>NH) or left under the same reaction conditions, it is likely that the initially formed *cis*-isomer bearing an electron-withdrawing substituent (R) epimerizes highly selectively at the acidic 3-CH, α to the lactam carbonyl group, to the thermodynamically more stable *trans*-isomer while maintaining high enantioselectivity.<sup>5,16</sup> Similarly, the reaction of ethyl propiolate (**7g**) produced *trans*-isomer **9p** only (54%) but, in this case, with low enantioselectivity (32% ee, entry 7). The reaction of aliphatic acetylene, 1-cyclohexenylacetylene (**7h**), gave **9q** (31%) in a cis:trans ratio of 83:17 with high enantioselectivity (83% ee, entry 8).

Although in the present reaction chemical yields and diastereoselectivities were not improved, the highest level of enantioselectivities were attained using the Cu(OTf)<sub>2</sub> complex with the simple and readily available C<sub>2</sub>-symmetric IndaBox ligand (**6a**) (commercially and cheaply available) and the optimal amine under the optimized reaction conditions.

**Table 4**

Catalytic enantioselective Kinugasa reaction: variation of the alkyne substituent R

Entry	<b>7</b>	R	Time (h)	Adduct	Yield <sup>a</sup> (%)	Cis:trans <sup>b</sup>	Cis ee <sup>c</sup> (%)
1	<b>7a</b>	Ph	38	<b>9a</b>	47	85:15	90
2	<b>7b</b>	<i>p</i> -MeC <sub>6</sub> H <sub>4</sub>	21	<b>9k</b>	38	63:37	93
3	<b>7c</b>	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	90	<b>9l</b>	40	83:17	94
4	<b>7d</b>	<i>p</i> -BrC <sub>6</sub> H <sub>4</sub>	336	<b>9m</b>	38	76:24	88
5	<b>7e</b>	<i>p</i> -CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	25	<b>9n</b>	55	72:28	91
6	<b>7f</b>	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	64	<b>9o</b>	51	18:82	86 <sup>d</sup>
7	<b>7g</b>	EtO <sub>2</sub> C	0.1	<b>9p</b>	54	Trans only	32 <sup>d</sup>
8	<b>7h</b>	1-Cyclohexenyl	36	<b>9q</b>	31	83:17	83

<sup>a</sup> Total yield of cis- and trans-isomers.<sup>b</sup> Determined by <sup>1</sup>H NMR.<sup>c</sup> Determined by chiral HPLC.<sup>d</sup> ee for trans-isomer.


## Supplementary data

Supplementary data (experimental procedure, full entries data of Table 1,  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data, compounds characterization data, and chiral HPLC copies of compounds **9**) associated with this article can be found, in the online version, at [doi:10.1016/j.tetlet.2009.06.050](https://doi.org/10.1016/j.tetlet.2009.06.050).

## References and notes

- (a) *Comprehensive Heterocyclic Chemistry II*; Katrizky, A. R., Rees, C. W., Scriven, E. F. V., Eds.; Pergamon: New York, 1996; Vol. 1B., Chapters 1.18–1.20 (b) *Synthesis of  $\beta$ -Lactam Antibiotics, Chemistry, Biocatalysis and Process Integration*; Bruggink, A., Ed.; Kluwer: Dordrecht, Netherlands, 2001; (c) Ojima, I.; Delaloge, F. *Chem. Soc. Rev.* **1997**, 26, 377–386.
- (a) France, S.; Weatherwax, A.; Taggi, A. E.; Lectka, T. *Acc. Chem. Res.* **2004**, 37, 592; (b) Hodous, B. L.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, 124, 1578–1579; (c) Taggi, A. E.; Hafez, A. M.; Wack, H.; Young, B.; Drury, W. J., III; Lectka, T. *J. Am. Chem. Soc.* **2000**, 122, 7831–7832; (d) Fujieda, H.; Kanai, M.; Kambara, T.; Iida, A.; Tomioka, K. *J. Am. Chem. Soc.* **1997**, 119, 2060–2061; (e) Anada, M.; Watanabe, N.; Hashimoto, S.-I. *Chem. Commun.* **1998**, 1517; (f) Calet, S.; Urso, F.; Alper, H. *J. Am. Chem. Soc.* **1989**, 111, 931–934.
- Reviews: (a) Marco-Contelles, J. *Angew. Chem., Int. Ed.* **2004**, 43, 2198–2200; (b) Pal, R.; Ghosh, S. C.; Chandra, K.; Basak, A. *Synlett* **2007**, 2321–2330.
- Kinugasa, M.; Hashimoto, S. *J. Chem. Soc., Chem. Commun.* **1972**, 466–467.
- Miura, M.; Enna, M.; Okuro, K.; Nomura, M. *J. Org. Chem.* **1995**, 60, 4999–5004.
- Lo, M. M.-C.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, 124, 4572–4573.
- (a) Ye, M.-C.; Zhou, J.; Huang, Z.-Z.; Tang, Y. *Chem. Commun.* **2003**, 2554–2555; (b) Ye, M.-C.; Zhou, J.; Huang, Z.-Z.; Tang, Y. *J. Org. Chem.* **2006**, 71, 3576–3582.
- Shintani, R.; Fu, G. C. *Angew. Chem., Int. Ed.* **2003**, 42, 4082–4085.
- Coyne, A. G.; Mueller-Bunz, H.; Guiry, P. J. *Tetrahedron: Asymmetry* **2007**, 18, 199–207.
- Basak, A.; Ghosh, S. C. *Synlett* **2004**, 1637.
- Stanley, L. M.; Sibi, M. P. *Chem. Rev.* **2008**, 108, 2887–2902.
- A referee pointed out that the reaction would proceed through Cu(I) acetylides as a key intermediate, rather than a Lewis acid-catalyzed reaction in the presence of a Lewis base. However, we believe that the following catalytic cycle is involved in this Kinugasa reaction. Cu(II) triflate and chiral ligand **6a** produce (TfO)<sub>2</sub>Cu(II)-**6a** complex. This complex actually works as the *chiral Lewis acid* in the catalytic cycle to give the copper acetylide-**6a**-complex (**X**) by the reaction with acetylide formed by the action of an amine (base) as shown in the scheme. The complex **X** is indeed the key species as the *chiral*

dipolarophile reacting with a nitron to finally give a  $\beta$ -lactam together with the reformed chiral Lewis acid catalyst after stereo-controlled protonation.



- Sibi et al. showed utility of IndaBox **6c**. (a) Sibi, M. P.; Ma, Z.; Jasperse, C. P. *J. Am. Chem. Soc.* **2004**, 126, 718–719; (b) Sibi, M. P.; Ma, Z.; Itoh, K.; Prabakaran, N.; Jasperse, C. P. *Org. Lett.* **2005**, 7, 2349–2352; (c) Sibi, M. P.; Liu, M. *Org. Lett.* **2000**, 2, 3393–3396; For reviews, see: (d) Ghosh, A. K.; Mathivanan, P.; Cappiello, J. *Tetrahedron: Asymmetry* **1998**, 9, 1; (e) Johnson, J. S.; Evans, D. A. *Acc. Chem. Res.* **2000**, 33, 325–335; (f) Thorhaug, J.; Roberson, M.; Hazell, R. G.; Jørgensen, K. A. *Chem. Eur. J.* **2002**, 8, 1888–1898; (g) Desimoni, G.; Faita, G.; Quadrelli, P. *Chem. Rev.* **2003**, 103, 3119–3154.
- Saito, T.; Takekawa, K.; Takahashi, T. *Chem. Commun.* **1999**, 1001–1002.
- Saito, T.; Yamada, T.; Miyazaki, S.; Otani, T. *Tetrahedron Lett.* **2004**, 45, 9581–9584.
- (a) Ding, L. K.; Irwin, W. J. *J. Chem. Soc., Perkin Trans. 1* **1976**, 2382; (b) Bose, A. K.; Narayanan, C. S.; Manhas, M. S. *Chem. Commun.* **1970**, 975–976; (c) Wagle, D. R.; Garai, C.; Chang, J.; Monteleone, M. G.; Kurys, B. E.; Strohmeyer, T. W.; Hegde, V. R.; Manhas, M. S.; Bose, A. K. *J. Org. Chem.* **1988**, 53, 4227–4236; (d) Alcaide, B.; Almendros, P.; Salgado, N. R.; Rodriguez-Vicente, A. *J. Org. Chem.* **2000**, 65, 4453–4455; (e) Alcaide, B.; Aly, M. F.; Rodriguez, C.; Rodriguez-Vicente, A. *J. Org. Chem.* **2000**, 65, 3453–3459; (f) Chiba, T.; Nakai, T. *Tetrahedron Lett.* **1985**, 26, 4647–4648.