



Palladium-catalyzed carboxylative cyclization of α -allenyl amines in dense carbon dioxide

Yoshihito Kayaki, Naoko Mori, Takao Ikariya *

Department of Applied Chemistry, Graduate School of Science and Engineering, Tokyo Institute of Technology, 2-12-1-E4-1 O-okayama, Meguro-ku, Tokyo 152-8552, Japan

ARTICLE INFO

Article history:

Received 12 August 2009

Revised 31 August 2009

Accepted 3 September 2009

Available online 6 September 2009

Keywords:

Allenes

Carbon dioxide fixation

Homogeneous catalysis

Palladium

Urethane

ABSTRACT


Carboxylative transformation of 2,3-allenic amines into 5-vinyl-1,3-oxazolidin-2-ones was promoted by palladium catalysts under a pressurized CO₂ condition. The presence of a stereogenic center adjacent to the allene group resulted in the formation of the corresponding cyclic urethane as a single trans-diastereomer.

© 2009 Elsevier Ltd. All rights reserved.

Transformation of carbon dioxide (CO₂) into synthetically useful compounds is of great interest from the viewpoints of utilization of naturally abundant carbon resources.¹ The CO₂ molecule having a Lewis acidic character can react with primary and secondary amines to form carbamic acids or ammonium carbamates, however, the resulting carbamate moieties revert easily to the parent amines by releasing CO₂ under atmospheric or reduced pressure.² As principal methods for transformation of the thermodynamically less stable carbamic acid families into robust functionalized molecules, urethane synthesis from CO₂ and amines has been broadly reported.^{3,4} On the basis of our spectroscopic studies on the facile formation of *N,N*-dialkylcarbamic acids from secondary amines in supercritical CO₂ (scCO₂), we have recently demonstrated the inter- and intra-molecular additions of the carbamic acids generated in scCO₂ to carbon–carbon triple bonds as well as ring-opening copolymerization of aziridines with scCO₂ affording stimuli-responsive poly(urethane-amine)s (Scheme 1).^{5,6} On the other hand, the addition of carbamic acid derivatives to carbon–carbon double bonds has scarcely been investigated, except for the stoichiometric transformation of ammonium carbamates and cyclic diolefins bound to a Pd center,⁷ the three-component reaction of CO₂, secondary amines, and vinyl ethers,⁸ and the iodine-mediated cyclization of allylic and homoallylic amines with CO₂.⁹ With regard to various metal-catalyzed nucleophilic cyclization of allenic compounds,¹⁰ aminoallenes are potential candidates for substrates

in catalytic CO₂ fixation reactions. Here, we disclose the intramolecular addition of amines having two cumulated double bonds with Pd catalysts in dense CO₂, affording cyclic urethanes.

The carboxylative cyclization of aminoallenes with CO₂ was conducted in a batch autoclave (50 mL) containing palladium catalysts with a substrate/catalyst ratio of 150. Table 1 summarizes the results of a test reaction of *N*-methyl-2,3-butadienylamine (**1a**) and CO₂ under supercritical conditions of 8.0 MPa and 50 °C for 15 h (Eq. 1). Although **1a** was quantitatively recovered after the reaction in the absence of metal catalysts (entry 1), palladium acetate catalyzed the CO₂ incorporation to give 3-methyl-5-



Scheme 1. Urethane synthesis via formation of carbamic acids.

* Corresponding author. Tel.: +81 3 5734 2636; fax: +81 3 5734 2637.

E-mail address: tikariya@apc.titech.ac.jp (T. Ikariya).


Table 1
Carboxylative cyclization of **1a**^a

Entry	Catalyst	Solvent	Yield of 2a ^b (%)
1	—	—	0
2	Pd(OCOCH ₃) ₂	—	41
3	PdCl ₂ [P(OC ₂ H ₅) ₃] ₂	—	20
4	Pd(OCOCH ₃) ₂	Toluene	46
5	PdCl ₂ [P(OC ₂ H ₅) ₃] ₂	Toluene	30
6	PdCl ₂ (CH ₃ CN) ₂	Toluene	37
7	Pd(dba) ₂	Toluene	37
8	PdCl ₂ [P(OC ₆ H ₅) ₃] ₂	Toluene	27
9	Pd[P(OC ₆ H ₅) ₃] ₄	Toluene	33
10	PdCl ₂ [P(C ₆ H ₅) ₃] ₂	Toluene	0
11	Pd[P(C ₆ H ₅) ₃] ₄	Toluene	0

^a Reaction conditions: The reaction was carried out using **1a** (1.5 mmol) with a substrate/catalyst ratio of 150 at 50 °C for 15 h under CO₂ (8.0 MPa).

^b Determined by ¹H NMR, using hexamethyldisilane as an internal standard.

vinyl-2-oxazolidinone (**2a**) in 41% yield as a 5-*exo* cyclization product (entry 2). A CO₂-soluble P(OC₂H₅)₃-Pd complex, which has been demonstrated as a suitable catalyst precursor for homogeneous reactions in scCO₂,¹¹ afforded a lower yield of **2a** (20%; entry 3). When toluene (1.5 mL) was added as a cosolvent to both the catalyst systems, the yields of **2a** were improved to 46% and 30%, respectively (entries 4 and 5), implying that the reaction may proceed in a liquid phase. In fact, visual inspection using a high-pressure view cell containing **1a** and toluene revealed that phase separation from the CO₂ phase occurred under the reaction conditions. Among the catalysts examined, both Pd(0) and Pd(II) precursors worked equally well. Pd complexes having P(OC₆H₅)₃ were also applicable to the urethane formation, whereas the use of the P(C₆H₅)₃ analogues resulted in a loss of catalytic activity (entries 8–11). Consequently, the following reactions were performed in the presence of toluene with Pd(OCOCH₃)₂ as the comparably effective catalyst.



By varying the CO₂ pressure at constant temperature of 50 °C, the maximum **2a** yield of 65% was obtained under the condition of 11.5 MPa (*d*_{CO₂} = 0.5 g mL⁻¹) for 15 h, as shown in Figure 1. The positive pressure effect can be explained by the enhancement of CO₂ uptake by amines. The reaction under higher pressure of CO₂ resulted in a drop in the yield, possibly due to a reduction of the




Figure 1. CO₂ pressure effect on yields of **2a** (●) and **3a** (○) for the carboxylative cyclization of **1a**. Reaction conditions: **1a** (1.5 mmol), Pd(OCOCH₃)₂ (0.01 mmol), toluene (1.5 mL) at 50 °C for 15 h.

liquid reaction phase in the high density region. Notably, a careful ¹H NMR analysis of each crude reaction mixture showed that a slight amount (<10%) of 1-methyl-3-pyrroline (**3a**) was formed simultaneously as a 5-*endo-trig* cyclization¹² product regardless of the CO₂ pressure.

Figure 2 shows the temperature dependence on the yields of **2a** at a constant density of CO₂ (0.5 g mL⁻¹). Although the catalytic activity was improved significantly by changing the temperature from 30 to 50 °C, a further increase in the temperature to 100 °C resulted in a lowering of the ratio of **2a/3a** to 30/22. It can be ascribed to the easy release of CO₂ from carbamic acid upon heating, as confirmed by our preceding NMR experiments.^{5c} Under the optimized conditions of 11.5 MPa and 50 °C, the use of other cosolvents such as CH₂Cl₂ and THF had an insignificant effect on the yield of **2a**.

The scope of the carboxylative cyclization under the standard conditions is summarized in Table 2. While secondary 2,3-butadienylamines (**1a–d**) were converted into the desired products with a yield range of 44–65%, the primary amine (**1e**) provided a poor result in line with our previous studies (entries 1–5).^{5b} It is noteworthy that the reaction of **1f** bearing a stereogenic center adjacent to the allene group proceeded smoothly to give the desired urethane as a single trans-diastereomer (Eq. 2). The relative stereochemistry was determined by a ³J_{HH} coupling constant of 7.8 Hz between the hydrogens attached to the C-4 and C-5 carbons on the oxazolidinone ring in the ¹H NMR spectrum, in addition to an NOE experiment. Similar diastereoselective reactions have been reported for the related 5-*exo* cyclization of α-substituted allenic substrates promoted by Pd catalysts.¹³





Figure 2. Reaction temperature dependence in the formation of **2a** (●) and **3a** (○). Reaction conditions: **1a** (1.5 mmol), Pd(OCOCH₃)₂ (0.01 mmol), CO₂ (*d* = 0.5 g mL⁻¹), toluene (1.5 mL) for 15 h.

Table 2
Carboxylative cyclization of **1a**^a

Entry	CH ₂ =C=CHCH ₂ NHR	Yield ^{b,c} (%)
1	R = CH ₃ , 1a	65
2	R = C ₂ H ₅ , 1b	44(39)
3	R = C ₃ H ₇ , 1c	54(49)
4	R = CH ₂ C ₆ H ₅ , 1d	53
5	R = H, 1e	2


^a Reaction conditions: The reaction was carried out using **1** (1.5 mmol) with a substrate/catalyst ratio of 150 in toluene (1.5 mL) at 50 °C for 15 h under CO₂ (11.5 MPa).

^b Determined by ¹H NMR.

^c Isolated yields were in parentheses.

A possible catalytic mechanism of the urethane formation is shown in Scheme 2. The catalytic reaction is likely initiated by the oxidative addition of carbamic acid generated from the allenic amine and CO₂, which is responsible for the formation of hydridopalladium species. The allene moiety inserts into the Pd–H bond to give an η³-allylpalladium intermediate, and the following reductive elimination would furnish the cyclization product.¹⁴ A similar mechanistic sequence has been expected for the Pd-catalyzed intramolecular hydroamination of allenes in which the presence of acetic acid enhances the rates and yields.¹⁵ In the light of the reaction conducted in dense CO₂ favoring the carbamic acid formation from secondary amines, the hypopalladation mechanism seems to be feasible, while an alternative mechanism involving nucleophilic attack on the allene bound to a Pd(II) center¹⁶ may be operative in the case of the 5-*endo* cyclization producing **3a**.¹⁷ Although 1,3-dienes may also be accessible to the η³-allylpalladium analogues by insertion into palladium hydrides, an attempt to extend the reaction to *N*-methyl-2,4-pentadienylamine was unsuccessful.

In summary, we have first accomplished the Pd-catalyzed cyclic urethane synthesis via intramolecular addition of carbamic acids across the carbon–carbon double bond in allenic substrates. The dense CO₂ conditions proved to be crucial for smooth CO₂ fixation. Although further improvement of the catalyst performance is needed for the reaction, the present direct addition system provides a straightforward method without forming salt waste, which is thus of great promise for achieving environmentally benign processes.



Scheme 2. A plausible mechanism for the palladium-catalyzed carboxylative cyclization of allenic amines.

Acknowledgments

This work is supported by a Grant-in-Aid for Scientific Research on Priority Areas (No. 18065007; ‘Chemistry of Concerto Catalysis’) and partially supported by GCOE program from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

References and notes

- (a) Gibson, D. H. *Chem. Rev.* **1996**, *96*, 2063–2096; (b) Sakakura, T.; Choi, J.-C.; Yasuda, H. *Chem. Rev.* **2007**, *107*, 2365–2387; (c) Aresta, M.; Dibenedetto, A. *Dalton Trans.* **2007**, 2975–2992; (d) *Green Chemistry Using Liquid and Supercritical Carbon Dioxide*; DeSimone, J. M., Tumas, W., Eds.; Oxford University Press: New York, 2003.
- (a) Wright, H. B.; Moore, M. B. *J. Am. Chem. Soc.* **1948**, *70*, 3865–3866; (b) Dell’Amico, D. B.; Calderazzo, F.; Labella, L.; Marchetti, F.; Pampaloni, G. *Chem. Rev.* **2003**, *103*, 3857–3898; (c) Aresta, M.; Ballivet-Tkatchenko, D.; Dell’Amico, D. B.; Bonnet, M. C.; Boschi, D.; Calderazzo, F.; Faure, R.; Labella, L.; Marchetti, F. *Chem. Commun.* **2000**, 1099–1100; (d) George, M.; Weiss, R. G. *J. Am. Chem. Soc.* **2001**, *123*, 10393–10394; (e) Hampe, E. M.; Rudkevich, D. M. *Chem. Commun.* **2002**, 1450–1451; (f) Hampe, E. M.; Rudkevich, D. M. *Tetrahedron* **2003**, *59*, 9619–9625; (g) Masuda, K.; Ito, Y.; Horiguchi, M.; Fujita, H. *Tetrahedron* **2005**, *61*, 213–229; (h) Dijkstra, Z. J.; Doornbos, A. R.; Weyten, H.; Ernsting, J. M.; Elsevier, C. J.; Keurentjes, J. T. F. *J. Supercrit. Fluids* **2007**, *41*, 109–114.
- Chaturvedi, D.; Ray, S. *Curr. Org. Chem.* **2007**, *11*, 987–998.
- (a) Dimroth, P.; Pasedach, H. DE Pat 1164411, 1964; (b) Sasaki, Y.; Dixneuf, P. H. *J. Chem. Soc., Chem. Commun.* **1986**, 790–791; (c) Mahé, R.; Dixneuf, P. H.; Lécolier, S. *Tetrahedron Lett.* **1986**, *27*, 6333–6336; (d) Sasaki, Y.; Dixneuf, P. H. *J. Org. Chem.* **1987**, *52*, 314–315; (e) Sasaki, Y.; Dixneuf, P. H. *J. Org. Chem.* **1987**, *52*, 4389–4391; (f) Mitsudo, T.; Hori, Y.; Yamakawa, Y.; Watanabe, Y. *Tetrahedron Lett.* **1987**, *28*, 4417–4418; (g) Mahe, R.; Sasaki, Y.; Bruneau, C.; Dixneuf, P. H. *J. Org. Chem.* **1989**, *54*, 1518–1523; (h) Costa, M.; Chiusoli, G. P.; Rizzardi, M. *J. Chem. Soc., Chem. Commun.* **1996**, 1699–1700; (i) Costa, M.; Chiusoli, G. P.; Taffurelli, D.; Dalmonego, G. *J. Chem. Soc., Perkin Trans. 1* **1998**, 1541–1546; (j) Rohr, M.; Geyer, C.; Wandeler, R.; Schneider, M. S.; Murphy, E. F.; Baiker, A. B. *Green Chem.* **2001**, *3*, 123–125; (k) Kim, T. J.; Kwon, K. H.; Kwon, S. C.; Baeg, J. O.; Shim, S. C.; Lee, D. H. *J. Organomet. Chem.* **1990**, *389*, 205–217; (l) Shi, M.; Shen, Y. *M. J. Org. Chem.* **2002**, *67*, 16–21; (m) Jiang, J.-L.; Hua, R. *Tetrahedron Lett.* **2006**, *47*, 953–955.
- (a) Kayaki, Y.; Suzuki, T.; Noguchi, Y.; Sakurai, S.; Imanari, M.; Ikariya, T. *Chem. Lett.* **2002**, *31*, 424–425; (b) Kayaki, Y.; Yamamoto, M.; Suzuki, T.; Ikariya, T. *Green Chem.* **2006**, *8*, 1019–1021; (c) Kayaki, Y.; Suzuki, T.; Ikariya, T. *Chem. Asian J.* **2008**, *3*, 1865–1870.
- (a) Ihata, O.; Kayaki, Y.; Ikariya, T. *Angew. Chem., Int. Ed.* **2004**, *43*, 717–719; (b) Ihata, O.; Kayaki, Y.; Ikariya, T. *Chem. Commun.* **2005**, 2268–2270; (c) Ihata, O.; Kayaki, Y.; Ikariya, T. *Macromolecules* **2005**, *38*, 6429–6434.
- McGhee, W. D.; Riley, D. P. *Organometallics* **1992**, *11*, 900–907.
- Yoshida, Y.; Inoue, S. *Chem. Lett.* **1977**, *6*, 1375–1376.
- Toda, T.; Kitagawa, Y. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 334–335.
- (a) Zimmer, R.; Dinesh, C. U.; Nandan, E.; Khan, F. A. *Chem. Rev.* **2000**, *100*, 3067–3125; (b) Bates, R. W.; Satcharoen, V. *Chem. Soc. Rev.* **2002**, *31*, 12–21; (c) Ma, S. *Chem. Rev.* **2005**, *105*, 2829–2871.
- Kayaki, Y.; Noguchi, Y.; Iwasa, S.; Ikariya, T.; Noyori, R. *Chem. Commun.* **1999**, 1235–1236.
- Related reports: (a) Claesson, A.; Sahlberg, C.; Luthman, K. *Acta Chem. Scand., Ser. B* **1979**, *33*, 309–310; (b) Kimura, M.; Harayama, H.; Tanaka, S.; Tamaru, Y. *J. Chem. Soc., Chem. Commun.* **1994**, 2531–2533.
- (a) Kimura, M.; Fugami, K.; Tanaka, S.; Tamaru, Y. *J. Org. Chem.* **1992**, *57*, 6377–6379; (b) Gallagher, T.; Davies, I. W.; Jones, S. W.; Lathbury, D.; Mahon, M. F.; Molloy, K. C.; Shaw, R. W.; Vernon, P. J. *Chem. Soc., Perkin Trans. 1* **1992**, 433–440.
- A facile reduction of Pd(II) to Pd(0) in the presence of amine has been well documented: Trzeczal, A. M.; Ciunik, Z.; Ziolkowski, J. *J. Organometallics* **2002**, *21*, 132–137 and referenced cited therein.
- Meguro, M.; Yamamoto, Y. *Tetrahedron Lett.* **1998**, *39*, 5421–5424.
- Hashmi, A. S. K.; Ruppert, T. L.; Knöfel, T.; Bats, J. W. *J. Org. Chem.* **1997**, *62*, 7295–7304.
- The cyclic urethane product **2a** containing an allylic carbon–oxygen bond proved to be stable even in the presence of Pd catalysts such as Pd(OAc)₂, Pd(dba)₂, and Pd[P(OC₆H₅)₃]₄ under the reaction conditions that will preclude the possibility of catalytic access from **2a** to **3a**.