



Stable solid-supported leucoanthocyanidin variants for flavanoid biosynthesis elucidation

Denis Deffieux^{a,b,*}, Sophie Gaudrel-Grosay^{a,b}, Axelle Grelard^b, Céline Chalumeau^{a,b}, Stéphane Quideau^{a,b,*}

^a Université de Bordeaux, Institut des Sciences Moléculaires (CNRS-UMR 5255), 351 cours de la libération, 33405 Talence Cedex, France

^b Institut Européen de Chimie et Biologie, 2 rue Robert Escarpit, 33607 Pessac Cedex, France

ARTICLE INFO

Article history:

Received 20 July 2009

Revised 7 September 2009

Accepted 9 September 2009

Available online 12 September 2009

Keywords:

Flavanoids

Leucoanthocyanidins

Anthocyanidins

Proanthocyanidins

Biosynthesis

ABSTRACT

Remarkable progress toward the complete elucidation of the biosynthesis of flavanoids has been accomplished during the last decade, but the final steps presumably involving the transformation of leucoanthocyanidins, which are highly unstable when free in aqueous solution, into both anthocyanidins and proanthocyanidins still remain to be fully understood. Herein is described the synthesis of stabilized solid-supported leucoanthocyanidin variants that should serve as valuable tools for in vitro studies aimed at investigating the metabolism of these seemingly fleeting common precursors of two main classes of flavanoids.

© 2009 Elsevier Ltd. All rights reserved.


The biosynthesis of flavo/flavanoids is probably one of the most extensively studied plant metabolic pathways.¹ Most of the genes encoding the enzymes that are responsible for the catalysis of each individual step have been isolated from different plant sources.² However, the last steps leading to the formation of flavylum-based anthocyani(di)ns and flavan-3-ol-derived proanthocyanidins remain unclear and are still under debate due in part to the lack of in vitro evidence.³ The putative biosynthesis pathway leading to these two important classes of flavanoids⁴ is depicted in Scheme 1. Leucoanthocyanidins [i.e., (2R,3S,4S)-flavan-2,3-trans-3,4-cis-diols] are presumed to be the common precursors of both anthocyani(di)ns and proanthocyanidins. Results from in vitro studies have shown that Leucoanthocyanidin DiOxygenase (LDOX, also referred to as ANS for ANthocyanidin Synthase), a 2-oxoglutarate-dependent enzyme, is involved in the conversion of colorless leucoanthocyanidins into colored anthocyanidins 3-O-glucosides.⁵ These anthocyanins can then undergo methylation of their ring-B hydroxyl groups.

However, the mechanistic details of the conversion of leucoanthocyanidins into anthocyanidins have not yet been fully elucidated due to the lack of structural information on the intermediate(s) entailed in this biotransformation.⁶ Even more speculative are the final stages of the biosynthesis of proanthocyanidins.⁷ These so-called condensed tannins would result from the polymerization of flavan-3-ols and/or flavan-3,4-diols (Scheme 1). At least two additional enzymes would be involved in the process leading to flavan-

3-ols from leucoanthocyanidins. Thus, (2R,3R)-cis-flavan-3-ols such as (–)-epicatechin indirectly derive from leucoanthocyanidins through the action of an ANthocyanidin Reductase (ANR) on anthocyanidins,^{3a} whereas a LeucoAnthocyanidin Reductase (LAR) directly converts leucoanthocyanidins into (2R,3S)-trans-flavan-3-ols such as (+)-catechin.⁸

Several questions about the (bio)polymerization leading to proanthocyanidins still remain unanswered such as those concerning the exact nature of the starter and extension units, as well as whether or not a dedicated enzyme (i.e., a 'proanthocyanidin synthase') is involved in the process.^{7,9} The keystone position of leucoanthocyanidins in this biosynthetic scheme obviously makes them substrates of choice for in vitro experiments aimed at addressing all of the above issues remaining obscure in these final steps of the biogenesis of anthocyani(di)ns and proanthocyanidins. Unfortunately, such experiments are far from being easy to conduct because of the notable instability of these flavan-3,4-cis-diol compounds, which are prone to polymerize even under slightly acidic conditions.^{1a,10} In aqueous solutions, highly reactive (protonated) *para*- and/or *ortho*-quinone methides resulting from a facile departure of the benzylic hydroxyl group at position 4 are probably responsible for the rapid formation of polymeric materials in an uncontrolled manner. This is also probably the main reason why 3,4-cis leucoanthocyanidins that have been previously generated by chemical synthesis were characterized under their protected methyl ether forms.¹¹ The very few enzymatic studies that have been carried out to date with these compounds in fact report the use of unprotected (2R,3S,4R)-diastereomers of

* Corresponding authors. Tel.: +33 540 00 30 10; fax: +33 540 00 22 15 (S.Q.).
E-mail address: s.quideau@iecb.u-bordeaux.fr (S. Quideau).



Scheme 1. Putative last steps of the biosynthesis of anthocyanins and proanthocyanidins.

leucoanthocyanidins that are claimed to be easily converted into the desired natural isomers under the conditions used for the enzymatic assays.⁵ However, the discrete chemical nature of the materials thus involved in these assays remains questionable.^{10b}


We thus decided to engage in efforts aimed at the elaboration of leucoanthocyanidin variants of enhanced stability yet still amenable to enzymatic studies *in vitro*. Three flavan-3,4-*cis*-diol derivatives (**7a–c**) convertible into some of the most abundant anthocyanins found in berries [i.e., malvidin 3-*O*-glucoside ($R_1 = R_2 = \text{OMe}$), peonidin 3-*O*-glucoside ($R_1 = \text{OMe}$, $R_2 = \text{H}$), and pelargonidin 3-*O*-glucoside ($R_1 = R_2 = \text{H}$)]¹² were synthesized and grafted onto a biocompatible solid support (*vide infra*). The premise behind this approach was that

thus immobilized at an appropriate mutual distance on an insoluble cross-linked polymer, the selected leucoanthocyanidin variants should find themselves in conditions approaching those of high (infinite) dilution, hence minimizing the possibilities of intermolecular reactions.¹³


A synthesis of solid-supported leucoanthocyanidins relying on an early grafting step using the polymer support as a protecting group could have constituted a more elegant strategy, but the difficulty of structurally characterizing polymer-bound intermediates by means of spectroscopical methods led us to favor a more practical approach based on the synthesis of the desired leucoanthocyanidin molecules prior to their loading onto the polymer support. The synthesis of the racemic protected leucoanthocyanidins **7a–c** was thus first accomplished (Scheme 2). In each of the three cases, the phenolic hydroxyl function at the 4'-position of ring-B had to be orthogonally protected in anticipation of its exploitation for the attachment to the polymer support. In addition, the protecting groups for all of the other hydroxyl functions had to be chosen while keeping in mind the necessity of removing them under mild conditions without perturbing the fragile leucoanthocyanidin molecules still attached to the polymer support. To meet these requirements, silyl groups were selected to protect the hydroxyl functions at positions 3, 4, 5, and 7, whereas a photolabile 6-nitroveratryl or, alternatively, a classic benzyl group was used for the orthogonal protection of the phenolic 4'-hydroxyl function.

The syntheses started with the partial silylation of commercially available 2,4,6-trihydroxyacetophenone using triisopropylsilyl triflate (TIPSOTf) and triethylamine (Et_3N) in CH_2Cl_2 at low temperature to furnish the phenolic ketone **1** in high yield. The phenolic hydroxyl group of commercial syringaldehyde was protected via a Williamson reaction using 4,5-dimethoxy-2-nitrobenzyl bromide to furnish **2a** in 97% yield (see Supplementary data). The other two ring-B precursors, that is, 4-benzyloxy-3-methoxybenzaldehyde (**2b**) and 4-benzyloxybenzaldehyde (**2c**), are both commercially available. The phenolic ketone **1** was then submitted to a Claisen–Schmidt aldol condensation reaction with each of the three aldehydes **2a–c** to deliver the expected chalcone products **3a–c** in moderate to good yields.^{11c,14} The chalcones **3a–c** were then cyclized to give rise directly to the flavones **4a–c** according to the Clark–Lewis method using sodium borohydride in 2-methoxyethanol (Scheme 2).^{11c,15,17a}

The following *cis*-dihydroxylation of the flavones **4a–c** was performed using a catalytic amount of osmium tetroxide in the presence of *N*-methylmorpholine *N*-oxide (NMO)¹⁶ to furnish the expected racemic 2,3-*trans*-3,4-*cis*-leucoanthocyanidins **5a–c** in



Scheme 2. Synthesis of silylated leucomalvidin (**7a**), leucopaeonidin (**7b**), and leucopelargonidin (**7c**).



Scheme 3. Grafting of leucoanthocyanidins onto a functionalized PEGA resin. (a) $\text{R}_1, \text{R}_2 = \text{OMe}$, (b): $\text{R}_1 = \text{OMe}$, $\text{R}_2 = \text{H}$; (c): $\text{R}_1, \text{R}_2 = \text{H}$.

good yields.¹⁷ The 2,3-trans-3,4-cis configuration was confirmed by ^1H NMR analysis, which indicated for each compound a $J_{2,3}$ coupling constant of about 10 Hz and a $J_{3,4}$ of 3 to 4 Hz. Next, the *cis*-3,4-diol units of **5a-c** were protected with a di-*tert*-butylsilyl group (Scheme 2). Previous attempts to independently protect

these vicinal hydroxyl groups with triisopropylsilyl group failed, as only one silyl group could be introduced using TIPSOTf in the presence of Et_3N . The ultimate step consisted in the deprotection of the 4'-hydroxyl group. Thus, the flavan-3,4-diol **6a** was subjected to irradiation using a high-pressure mercury lamp ($\lambda > 300$ nm) to furnish the silyl-protected leucomalvidin **7a** in 11% overall yield. The corresponding leucopaeonidin **7b** and leucopelargonidin **7c** were obtained after hydrogenolysis of **6b** and **6c**, respectively, in 32% and 15% yields over the six steps of their synthesis.

The instability of free leucoanthocyanidins was here verified by monitoring via ^1H NMR analysis the *quasi* immediate and steady degradation of a sample of leucopaeonidin **7b** in CDCl_3 undergoing desilylation with a solution of TBAF in THF at room temperature (see Supplementary data).

With adequately protected leucoanthocyanidins **7a-c** in hand, we turned our attention to the selection of a resin on which to graft them with the following criteria in mind. Such a resin should be not only appropriately functionalized to enable the attachment of **7a-c** via their free phenolic hydroxyl group, but also made of a polymeric material capable of swelling extensively in aqueous media for allowing access of proteins (enzymes) into the resin beads during biochemical assays. Moreover, the loading onto the resin beads should not be too high to avoid the risk of intermolecular reactions between leucoanthocyanidins once deprotected. The polyethyleneglycol-polyacrylamide (PEGA) resin with a potential maximum loading of 0.4 mmol/g meets these criteria.¹⁸

Succinic anhydride was used to install a functionalized linker compatible with the attachment of **7a-c** (Scheme 3). This requisite adaptation of the PEGA resin was conveniently achieved in the




Figure 1. (Top) ^1H NMR spectrum of silylated leucopaeonidin **7b** in CDCl_3 ; (middle) HR-MAS ^1H NMR spectrum of PEGA-supported silylated leucopaeonidin **9b** in CDCl_3 ; and (bottom) HR-MAS ^1H NMR spectrum of PEGA-supported acetylated leucopaeonidin **11b** in CDCl_3 .

presence of Hunig's base in DMF.¹⁹ Activation by microwaves permitted to drastically reduce the reaction time from about 5 days to only 15 min at 50 °C (see Supplementary data). Then, esterification of the resulting carboxylic acid **8** to the free phenolic function of **7a–c** was accomplished in a standard fashion using EDCI/DMAP as the condensation reagents (Scheme 3). To evaluate the extent of the loading onto the resin, an aliquot of the PEGA-supported silylated leucoanthocyanidin **9b** was treated with sodium methoxide in a CH₂Cl₂/MeOH (2:1) mixture. The quantity of **7b** thus released and estimated by HPLC analysis indicated a loading onto the resin of about 50% (i.e., 0.2 mmol/g).

The PEGA-supported silylated leucoanthocyanidins **9a–c** were characterized by high-resolution magic angle spinning (HR-MAS) NMR analysis. Comparison of their HR-MAS ¹H NMR spectra with the standard ¹H NMR spectra of the silylated leucoanthocyanidins **7a–c** in solution indicated an excellent level of concordance between the diagnostic signals of these pairs of spectra (see Fig. 1 and Supplementary data). The absence of methoxy signal(s) in the spectra of **9a** and **9b** (see Fig. 1 for **9b**, middle spectrum) is a consequence of the presaturation applied at 3.6 ppm to suppress the overwhelming ethylene signals from the PEG chains of the resin.²⁰

The desilylation of **9a–c** caused us some difficulties, as the use of TBAF in THF or cesium fluoride in DMF consistently led to cleavage of the ester linkage to the resin.²¹ We thus had to rely on the use of Et₃N–3HF in THF at room temperature (Scheme 3), but the corresponding HR-MAS ¹H NMR spectra run in either CDCl₃ or a CDCl₃/DMSO (1:1) solvent mixture exhibited such a low resolution that confirmation of the efficiency of this desilylation step leading to **10a–c** could not be made by this means. Nevertheless, good quality spectra were obtained in CDCl₃ after peracetylation leading to the PEGA-supported leucoanthocyanidin derivatives **11a–c** (see Fig. 1 for **11b**, bottom spectrum). The coupled proton spin system of their cycle C (i.e., H₂, H₃, and H₄) could clearly be assigned by 2D COSY HR-MAS NMR experiments (see Fig. 2 for **11b**). These satisfactory HR-MAS NMR analyses of **11a–c** not only confirmed the success of the preceding desilylation step (Scheme 3), but also provided a sound indication that the resulting PEGA-supported

leucoanthocyanidins **10a–c** are thus granted with a much higher stability than their free counterparts, since they survive successive exposures to desilylation, HR-MAS NMR analysis, and acetylation conditions (see Supplementary data).²²

In summary, the synthesis of three leucoanthocyanidin derivatives immobilized on a biocompatible PEGA resin via their 4'-hydroxyl group was achieved. These materials constitute stabilized variants of these otherwise fleeting flavan-3,4-diol species that should serve, in spite of their racemic nature, as valuable tools for in vitro enzymatic studies aimed at elucidating the final steps of the biosynthesis of both anthocyanidins and proanthocyanidins. Affinity and biochemical reactivity tests with LDOX are in progress and the results will be reported in due course.

Acknowledgments

We wish to thank the Conseil Interprofessionnel du Vin de Bordeaux for their generous financial support, including research assistantships for Sophie Gaudrel-Grosay and Céline Chalumeau. This project was undertaken within the framework of the association of our research team with the Institut des Sciences de la Vigne et du Vin, Bordeaux-Aquitaine, France.

Supplementary data

Supplementary data (experimental procedures, characterization data, and NMR spectra for all major compounds) associated with this article can be found, in the online version, at doi:10.1016/j.tetlet.2009.09.045.

References and notes

- (a) Forkmann, G.; Heller, W. In *Comprehensive Natural Products Chemistry*; Sankawa, U., Ed.; Pergamon: Oxford, 1999; Vol. 1, pp 713–748; (b) Davies, K. M.; Schwinn, K. E. In *Flavonoids: Chemistry, Biochemistry and Applications*; Andersen, Ø. M., Markham, K. R., Eds.; CRC-Taylor & Francis: Boca Raton, FL, 2006; pp 143–218.
- Holton, T. A.; Cornish, E. C. *The Plant Cell* **1995**, *7*, 1071–1083.
- (a) Xie, D.-Y.; Sharma, S. B.; Paiva, N. L.; Ferreira, D.; Dixon, R. A. *Science* **2003**, *299*, 396–399; (b) Xie, D.-Y.; Dixon, R. A. *Phytochemistry* **2005**, *66*, 2127–2144.
- (a) Williams, C. A.; Grayer, R. J. *Nat. Prod. Rep.* **2004**, *21*, 539–573; (b) Ferreira, D.; Nel, R. J. J.; Bekker, R. In *Comprehensive Natural Products Chemistry*; Pinto, B. M., Ed.; Pergamon: Oxford, 1999; Vol. 3, pp 747–797.
- (a) Saito, K.; Kobayashi, M.; Gong, Z.; Tanaka, Y.; Yamazaki, M. *Plant J.* **1999**, *17*, 181–189; (b) Turnbull, J. J.; Sobey, W. J.; Aplin, R. T.; Hassan, A.; Firmin, J. L.; Schofield, C. J.; Prescott, A. G. *Chem. Commun.* **2000**, 2473–2474; (c) Nakajima, J.; Tanaka, Y.; Yamazaki, M.; Saito, K. *J. Biol. Chem.* **2001**, *276*, 25797–25803.
- (a) Wellmann, F.; Griesser, M.; Schwab, W.; Martens, S.; Eisenreich, W.; Matern, U.; Lukacin, R. *FEBS Lett.* **2006**, *580*, 1642–1648; (b) Turnbull, J. J.; Nakajima, J.; Welford, R. W. D.; Yamazaki, M.; Saito, K.; Schofield, C. J. *J. Biol. Chem.* **2004**, *279*, 1206–1216.
- Dixon, R. A.; Xie, D.-Y.; Sharma, S. B. *New Phytol.* **2005**, *165*, 9–28.
- Tanner, G. J.; Francki, K. T.; Abrahams, S.; Watson, J. M.; Larkin, P. J.; Ashton, A. R. *J. Biol. Chem.* **2003**, *278*, 31647–31656.
- (a) Tian, L.; Pang, Y.; Dixon, R. A. *Phytochem. Rev.* **2008**, *7*, 445–465; (b) He, F.; Pan, Q.-H.; Shi, Y.; Chang-Qing, D. *Molecules* **2008**, *13*, 2674–2703.
- (a) Halbwirth, H.; Kahl, S.; Jäger, W.; Reznicek, G.; Forkmann, G.; Stich, K. *Plant Sci.* **2006**, *170*, 587–595; (b) Stafford, H. A.; Lester, H. H.; Porter, L. J. *Phytochemistry* **1985**, *24*, 333–338.
- (a) Porter, L. J.; Foo, L. Y. *Phytochemistry* **1982**, *21*, 2947–2952; (b) Steenkamp, J. A.; Ferreira, D.; Roux, D. G. *Tetrahedron Lett.* **1985**, *26*, 3045–3048; (c) Marais, J. P. J.; Ferreira, D.; Slade, D. *Phytochemistry* **2005**, *66*, 2145–2176. and references cited therein.
- (a) Andersen, Ø.; Jordheim, M. In *Flavonoids: Chemistry, Biochemistry and Applications*; Andersen, Ø. M., Markham, K. R., Eds.; CRC-Taylor & Francis: Boca Raton, FL, 2006; pp 471–551; (b) Munõz, S.; Mestres, M.; Busto, O.; Guasch, J. *Anal. Chim. Acta* **2008**, *628*, 104–110.
- Patchornik, A.; Kraus, M. A. J. *Am. Chem. Soc.* **1970**, *92*, 7587–7589.
- (a) Claisen, L.; Claparède, A. *Ber. Dtsch. Chem. Ges.* **1881**, *14*, 2460–2468; (b) Schmidt, J. G. *Ber. Dtsch. Chem. Ges.* **1881**, *14*, 1459–1461.
- (a) Clark-Lewis, J. W.; Jemison, R. W.; Skingle, D. C.; Williams, L. R. *Chem. Ind.* **1967**, 1455–1456; (b) Clark-Lewis, J. W.; Skingle, D. C. *Aust. J. Chem.* **1967**, *20*, 2169–2190.
- For a review on osmium tetroxide-mediated cis-hydroxylation of olefins, see: Schröder, M. *Chem. Rev.* **1980**, *80*, 187–213.




Figure 2. Diagnostic portion of ¹H–¹H HR-MAS COSY map of PEGA-supported acetylated leucoponidin **11b** in CDCl₃.

17. (a) Kawamoto, H.; Nakatsubo, F.; Murakami, K. *J. Wood Chem. Technol.* **1989**, *9*, 35–52; (b) Nay, B.; Arnaudinaud, V.; Peyrat, J.-F.; Nuhrich, A.; Deffieux, G.; Méryllon, J.-M.; Vercauteren, J. *Eur. J. Org. Chem.* **2000**, 1279–1283.
18. (a) Meldal, M. *Tetrahedron Lett.* **1992**, *33*, 3077–3080; (b) Auzanneau, F.-I.; Meldal, M.; Bock, K. *J. Pept. Sci.* **1995**, *1*, 31–44.
19. Fruchart, J.-S.; Grass-Masse, H.; Melnyk, O. *Tetrahedron Lett.* **1999**, *40*, 6225–6228.
20. Keifer, P. A. *J. Org. Chem.* **1996**, *61*, 1558–1559.
21. Curtis, A. D. M.; Humphries, M. J.; Ramsden, C. A. *ARKIVOC* **2000**, *iii*, 218–227.
22. However, for long-term storage, these PEGA-supported leucoanthocyanidins were kept under their silylated forms **9a–c** in the freezer at -18°C . After several weeks of storage under these conditions, HR-MAS ^1H NMR analysis also indicated no degradation of these materials.