



## New Gilman-type lithium cuprate from a copper(II) salt: synthesis and deprotonative cupration of aromatics

Tan Tai Nguyen, Floris Chevallier, Viatcheslav Jouikov\*, Florence Mongin\*

Chimie et Photonique Moléculaires, UMR 6510 CNRS, Université de Rennes 1, Bâtiment 10A, Case 1003, Campus Scientifique de Beaulieu, 35042 Rennes Cedex, France

### ARTICLE INFO

#### Article history:

Received 28 July 2009

Revised 11 September 2009

Accepted 18 September 2009

Available online 24 September 2009

#### Keywords:

Lithium

Copper

Metalation

Aromatic compound

Heterocycle

### ABSTRACT

Deprotonative cupration of aromatics including heterocycles (anisole, 1,4-dimethoxybenzene, thiophene, furan, 2-fluoropyridine, 2-chloropyridine, 2-bromopyridine, and 2,4-dimethoxypyrimidine) was realized in tetrahydrofuran at room temperature using the Gilman-type amido-cuprate  $(\text{TMP})_2\text{CuLi}$  in situ prepared from  $\text{CuCl}_2\cdot\text{TMEDA}$  through successive addition of 1 equiv of butyllithium and 2 equiv of LiTMP. The intermediate lithium (hetero)arylcuprates were evidenced by trapping with iodine, allyl bromide, methyl iodide, and benzoyl chlorides, the latter giving the best results. Symmetrical dimers were also prepared from lithium azine and diazine cuprates using nitrobenzene as an oxidative agent.

© 2009 Elsevier Ltd. All rights reserved.

Lithium bases have been largely employed for the deprotonation of aromatic rings.<sup>1</sup> Even if less nucleophilic hindered lithium dialkylamides are more suitable for the metalation of aromatics bearing reactive functions or sensitive  $\pi$ -deficient heterocycles, low reaction temperatures are required due to the high reactivity of the corresponding (hetero)aryllithiums.

The use of bimetallic bases in order to get more efficient and/or more chemoselective reactions is a challenging field. LIC–KOR (LIC = butyllithium, KOR = potassium *tert*-butoxide) first described by Schlosser<sup>2</sup> and Lochmann,<sup>3</sup> and  $\text{BuLi-LiDMAE}$  (DMAE = 2-dimethylaminoethoxide) introduced by Caubère<sup>4</sup> and developed further by Gros and Fort in the pyridine series<sup>5</sup> are well-known examples of synergic (or superbasic) mixtures of organolithiums and alkali metal alkoxides. By combining soft organometallic compounds with alkali (or alkaline-earth metal) additives, bases have been more recently prepared and used to generate functionalized aromatic compounds including heterocycles.<sup>6</sup> Examples are  $\text{R}_2\text{Zn}(\text{TMP})\text{Li}\cdot(\text{TMEDA})$  ( $\text{R} = \text{tBu}, \text{Bu}$ ; TMP = 2,2,6,6-tetramethyl piperidino, TMEDA = *N,N,N',N'*-tetramethylethylenediamine) (described by the groups of Kondo, Uchiyama, Mulvey, and Hevia),<sup>7</sup>  $(\text{TMP})_2\text{Zn}\cdot 2\text{MgCl}_2\cdot 2\text{LiCl}$ <sup>8</sup> and  $\text{TMPZnCl}\cdot\text{LiCl}$ <sup>9</sup> (Knochel),  $\text{tBu}_3\text{Al}(\text{TMP})\text{Li}$  (Uchiyama and Mulvey),<sup>10</sup>  $\text{Al}(\text{TMP})_3\cdot 3\text{LiCl}$  (Knochel),<sup>11</sup>  $(\text{Me}_3\text{SiCH}_2)_2\text{Mn}(\text{TMP})\text{Li}\cdot\text{TMEDA}$  (Mulvey),<sup>12</sup> and  $\text{MeCu}(\text{TMP})(\text{CN})\text{Li}_2$  (Uchiyama and Wheatley).<sup>13</sup>


We recently accomplished the deprotonation of a large range of aromatics including heterocycles using a newly developed lithium–cadmium base,  $(\text{TMP})_3\text{CdLi}$ , in situ prepared from  $\text{CdCl}_2\cdot\text{TMEDA}$  and 3 equivalents of LiTMP.<sup>14</sup> We here describe the synthesis of putative  $(\text{TMP})_2\text{CuLi}$  using the same approach, and its use to deprotonate aromatic substrates.

Wheatley and Uchiyama documented in 2007 the first direct metalation using a lithium cuprate.<sup>13</sup> The authors showed that Gilman-type amidocuprates prepared from CuI were less efficient bases than Lipshutz-type amidocuprates prepared from CuCN.  $\text{MeCu}(\text{TMP})(\text{CN})\text{Li}_2$  was identified as the best base when used at the rate of 2 equiv in tetrahydrofuran (THF) at 0 °C.


Our approach was based on the in situ generation of putative  $(\text{TMP})_2\text{CuLi}$  from  $\text{CuCl}_2\cdot\text{TMEDA}$ <sup>15</sup> by (i) reduction of Cu(II) to Cu(I) and (ii) formation of the lithium cuprate by addition of 2 equiv of LiTMP. The reaction of copper(II) chloride with lithium bis(trimethylsilyl)amide at the reflux temperature of THF being known to produce the corresponding copper(I) amide,<sup>16</sup> the generation of  $(\text{TMP})_2\text{CuLi}$  starting from  $\text{CuCl}_2\cdot\text{TMEDA}$  and using 3 equiv of LiTMP was first considered. Electron paramagnetic resonance (EPR) response of Cu(II) was used in order to check the validity of the reduction step using LiTMP. To this purpose, the EPR spectrum of a THF solution prepared from  $\text{CuCl}_2\cdot\text{TMEDA}$  and LiTMP ( $2\cdot 10^{-3}$  M each) was collected, and was compared with a spectrum recorded from a THF solution of  $\text{CuCl}_2\cdot\text{TMEDA}$  ( $2\cdot 10^{-3}$  M). It was observed that the reduction to Cu(I) was not quantitative, even after 20 h at room temperature, and that TEMPO was formed, probably by reaction between LiTMP and dissolved oxygen<sup>17</sup> (Fig. 1).<sup>18</sup> It was then decided to attempt the use of butyllithium for the

\* Corresponding authors. Fax: +33 (0) 2 2323 6955 (F.M.).

E-mail addresses: [viatcheslav.jouikov@univ-rennes1.fr](mailto:viatcheslav.jouikov@univ-rennes1.fr) (V. Jouikov), [florence.mongin@univ-rennes1.fr](mailto:florence.mongin@univ-rennes1.fr) (F. Mongin).




**Figure 1.** EPR spectra of (i) a THF solution prepared from  $\text{CuCl}_2\cdot\text{TMEDA}$  and  $\text{LiTMP}$  ( $2.10^{-3}$  M each), (ii) the same solution 20 h later, and (iii) a THF solution of TEMPO ( $g = 2.007$ ,  $a_N = 15.54$  G).



**Scheme 1.** Synthesis of  $(\text{TMP})_2\text{CuLi}$  from  $\text{CuCl}_2\cdot\text{TMEDA}$  as copper(II) source.

**Table 1**  
Deproto-cupration of **1a,b** followed by trapping with electrophiles




Entry	Substrate	Electrophile	Product, yield (%)
1	<b>1a</b>	$\text{I}_2$	<b>2a</b> , 20
2	<b>1b</b>	$\text{I}_2$	<b>2b</b> , 19
3	<b>1b</b>	$\text{CH}_2=\text{CHCH}_2\text{Br}$	<b>3b</b> , 41
4	<b>1b</b>	$\text{C}_6\text{H}_5\text{COCl}$	<b>4b</b> , 53
5	<b>1b</b>	$4\text{-ClC}_6\text{H}_4\text{COCl}$	<b>5b</b> , 57

reduction of  $\text{CuCl}_2\cdot\text{TMEDA}$  at  $0^\circ\text{C}$ , and the EPR spectrum of a THF solution prepared from  $\text{CuCl}_2\cdot\text{TMEDA}$  and  $\text{BuLi}$  ( $2.10^{-3}$  M each) was recorded. The spectrum showing a complete reduction of  $\text{Cu(II)}$ , we prepared the mixed lithium–copper base as depicted in


**Scheme 1**, and recorded again the spectra of the solution. The only observable signal being attributed to  $\text{TEMPO}$ ,<sup>19</sup> the reactivity of such prepared  $(\text{TMP})_2\text{CuLi}$  was studied toward various aromatics.

With anisole (**1a**) and the 4-methoxy derivative **1b** as substrates, the metalation reactions performed in THF at room temperature by using 1 equiv of in situ prepared  $(\text{TMP})_2\text{CuLi}$ , followed by subsequent trapping with elemental iodine after 2 h, proceeded in low yields (compounds **2a**<sup>20</sup> and **2b**,<sup>21</sup> Table 1, entries 1 and 2). Concomitant formation of 2,2'-dimethoxybiphenyl<sup>22</sup> and *N*-(2-methoxyphenyl)-2,2,6,6-tetramethylpiperidine<sup>23</sup> was observed from




**Scheme 2.** Deproto-cupration of **6a,b** followed by trapping with benzoyl chlorides.

**Table 2**  
Deproto-cupration of **9a–c** followed by trapping with electrophiles or oxidation



Entry	Substrate	Electrophile or oxidative agent	Product, yield %
1	<b>9a</b>	$\text{C}_6\text{H}_5\text{COCl}$	<b>10a</b> , 85
2	<b>9b</b>	$\text{C}_6\text{H}_5\text{COCl}$	<b>10b</b> , 78
3	<b>9c</b>	$\text{C}_6\text{H}_5\text{COCl}$	<b>10c</b> , 20
4	<b>9a</b>	$4\text{-ClC}_6\text{H}_4\text{COCl}$	<b>11a</b> , 83
5	<b>9a</b>	$\text{I}_2$	<b>12a</b> , 31
6	<b>9a</b>	$\text{C}_6\text{H}_5\text{NO}_2$	<b>13a</b> , 84



**Scheme 3.** Deprotonation of **14** followed by trapping with electrophiles or oxidation. (See above-mentioned references for further information.)

anisole (**1a**), probably through reactions occurring during the trapping step with iodine. Indeed, it was noted that carrying out the reaction using water as electrophile instead of iodine resulted in recovered anisole. Whereas the use of ethyl acrylate and enones such as 2-cyclohexen-1-one and 2-cyclopenten-1-one did not allow any trapping products, reactions proved successful when performed with allyl bromide and benzoyl chlorides at the reflux temperature of THF (entries 3–5) to afford the allylated derivative **3b**,<sup>24</sup> and the benzophenones **4b**<sup>25</sup> and **5b**<sup>26</sup> in yields ranging from 41% to 57%.

We next demonstrated that the cuprate base was suitable for the metalation of both  $\pi$ -excessive and  $\pi$ -deficient aromatic heterocycles. Thiophene (**6a**) and furan (**6b**) proved to react under the same reaction conditions to furnish after trapping with benzoyl chlorides the expected ketones **7a**,<sup>20</sup> **8a**,<sup>27</sup> and **7b**<sup>27</sup> (Scheme 2).

2-Halopyridines **9a–c** were similarly successively treated with the amido-cuprate and benzoyl chlorides. Whereas a low 20% yield (compound **10c**)<sup>28</sup> was obtained starting from 2-bromopyridine (**9c**) and using benzoyl chloride, side reactions were not observed with its lighter isomers **9a,b**, and the expected ketones **10a**,<sup>29</sup> **10b**,<sup>30</sup> and **11a**<sup>31</sup> were isolated in good yields. Using iodine as an electrophile in the case of 2-fluoropyridine (**9a**) resulted in the formation of the iodide **12a**<sup>20</sup> in a moderate 31% yield due to the concomitant formation of the 3,3'-dimer **13a**.<sup>32</sup> The latter was obtained in a high 84% yield using nitrobenzene,<sup>33</sup> which proved to be in this reaction a better oxidant than iodine (Table 2).

2,4-Dimethoxyypyrimidine (**14**) was finally similarly involved in the reaction to afford after subsequent trapping with benzoyl chlorides, allyl bromide and iodomethane the regioselectively functionalized derivatives **15–18** in yields ranging from 19% to 52%;<sup>34</sup> the corresponding 5,5'-dimer **19** also formed using nitrobenzene as an oxidative agent (Scheme 3).

In summary, deprotonative cupration of aromatics was realized using a Gilman-type amido-cuprate in situ prepared from stable  $\text{CuCl}_2$ -TMEDA.<sup>37</sup> The intermediate lithium arylcuprates were notably evidenced by trapping with benzoyl chlorides in satisfying yields.

Further development will notably concern the impact of TEMPO on reaction yields. Indeed, it is known from the literature that TEMPO is an excellent ligand for alkali metals, and its presence in TMP-containing complexes can modify the availability of the amido groups.<sup>38</sup>


## Acknowledgment

We gratefully acknowledge the financial support of MESR of France (to T.T.N.).

## References and notes

- (a) Gschwend, H. W.; Rodriguez, H. R. *Org. React.* **1979**, *26*, 1–360; (b) Beak, P.; Snieckus, V. *Acc. Chem. Res.* **1982**, *15*, 306–312; (c) Snieckus, V. *Chem. Rev.* **1990**, *90*, 879–933; (d) Gant, T. G.; Meyers, A. I. *Tetrahedron* **1994**, *50*, 2297–2360; (e) Schlosser, M. In *Organometallics in Synthesis*; Schlosser, M., Ed., 2nd ed.; Wiley, 2002. Chapter 1.
- Schlosser, M. *Pure Appl. Chem.* **1988**, *60*, 1627–1634.
- Lochmann, L. *Eur. J. Inorg. Chem.* **2000**, 1115–1126.
- Caubère, P. *Chem. Rev.* **1993**, *93*, 2317–2334.
- Gros, P.; Fort, Y. *Eur. J. Org. Chem.* **2002**, 3375–3383.
- For reviews, see: (a) Mulvey, R. E. *Organometallics* **2006**, *25*, 1060–1075; (b) Mulvey, R. E.; Mongin, F.; Uchiyama, M.; Kondo, Y. *Angew. Chem., Int. Ed.* **2007**, *46*, 3802–3824; (c) Mulvey, R. E. *Acc. Chem. Res.* **2009**, *42*, 743–755.
- (a) Kondo, Y.; Shilai, M.; Uchiyama, M.; Sakamoto, T. *J. Am. Chem. Soc.* **1999**, *121*, 3539–3540; (b) Uchiyama, M.; Miyoshi, T.; Kajihara, Y.; Sakamoto, T.; Otani, Y.; Ohwada, T.; Kondo, Y. *J. Am. Chem. Soc.* **2002**, *124*, 8514–8515; (c) Barley, H. R. L.; Clegg, W.; Dale, S. H.; Hevia, E.; Honeyman, G. W.; Kennedy, A. R.; Mulvey, R. E. *Angew. Chem., Int. Ed.* **2005**, *44*, 6018–6021; (d) Clegg, W.; Dale, S. H.; Hevia, E.; Honeyman, G. W.; Mulvey, R. E. *Angew. Chem., Int. Ed.* **2006**, *45*, 2370–2374; (e) Clegg, W.; Dale, S. H.; Harrington, R. W.; Hevia, E.; Honeyman, G. W.; Mulvey, R. E. *Angew. Chem., Int. Ed.* **2006**, *45*, 2374–2377; (f) Clegg, W.; Dale, S. H.; Drummond, A. M.; Hevia, E.; Honeyman, G. W.; Mulvey, R. E. *J. Am. Chem. Soc.* **2006**, *128*, 7434–7435; (g) Uchiyama, M.; Kobayashi, Y.; Furuyama, T.; Nakamura, S.; Kajihara, Y.; Miyoshi, T.; Sakamoto, T.; Kondo, Y.; Morokuma, K. *J. Am. Chem. Soc.* **2008**, *130*, 472–480; (h) Clegg, W.; Conway, B.; Hevia, E.; McCall, M. D.; Russo, L.; Mulvey, R. E. *J. Am. Chem. Soc.* **2009**, *131*, 2375–2384.
- (a) Wunderlich, S. H.; Knochel, P. *Angew. Chem., Int. Ed.* **2007**, *46*, 7685–7688; (b) Wunderlich, S.; Knochel, P. *Chem. Commun.* **2008**, 6387–6389; (c) Wunderlich, S. H.; Knochel, P. *Org. Lett.* **2008**, *10*, 4705–4707; (d) Mosrin, M.; Knochel, P. *Chem. Eur. J.* **2009**, *15*, 1468–1477.
- Mosrin, M.; Knochel, P. *Org. Lett.* **2009**, *11*, 1837–1840.
- (a) Uchiyama, M.; Naka, H.; Matsumoto, Y.; Ohwada, T. *J. Am. Chem. Soc.* **2004**, *126*, 10526–10527; (b) Garcia-Alvarez, J.; Graham, D. V.; Kennedy, A. R.; Mulvey, R. E.; Weatherstone, S. *Chem. Commun.* **2006**, 3208–3210; (c) Garcia-Alvarez, J.; Hevia, E.; Kennedy, A. R.; Klett, J.; Mulvey, R. E. *Chem. Commun.* **2007**, 2402–2404; (d) Conway, B.; Hevia, E.; Garcia-Alvarez, J.; Graham, D. V.; Kennedy, A. R.; Mulvey, R. E. *Chem. Commun.* **2007**, 5241–5243; (e) Naka, H.; Uchiyama, M.; Matsumoto, Y.; Wheatley, A. E. H.; McPartlin, M.; Morey, J. V.; Kondo, Y. *J. Am. Chem. Soc.* **2007**, *129*, 1921–1930; (f) Naka, H.; Morey, J. V.; Haywood, J.; Eisler, D. J.; McPartlin, M.; Garcia, F.; Kudo, H.; Kondo, Y.; Uchiyama, M.; Wheatley, A. E. H. *J. Am. Chem. Soc.* **2008**, *130*, 16193–16200.
- Wunderlich, S. H.; Knochel, P. *Angew. Chem., Int. Ed.* **2009**, *48*, 1501–1504.
- Garcia-Alvarez, J.; Kennedy, A. R.; Klett, J.; Mulvey, R. E. *Angew. Chem., Int. Ed.* **2007**, *46*, 1105–1108.

13. Usui, S.; Hashimoto, Y.; Morey, J. V.; Wheatley, A. E. H.; Uchiyama, M. *J. Am. Chem. Soc.* **2007**, *129*, 15102–15103.
14. (a) L'Helgoual'ch, J.-M.; Bentabed-Ababsa, G.; Chevallier, F.; Yonehara, M.; Uchiyama, M.; Derdour, A.; Mongin, F. *Chem. Commun.* **2008**, 5375–5377; (b) Snégaroff, K.; L'Helgoual'ch, J.-M.; Bentabed-Ababsa, G.; Nguyen, T. T.; Chevallier, F.; Yonehara, M.; Uchiyama, M.; Derdour, A.; Mongin, F. *Chem. Eur. J.* **2009**, *15*, 10280–10290; See also: (c) L'Helgoual'ch, J.-M.; Bentabed-Ababsa, G.; Chevallier, F.; Derdour, A.; Mongin, F. *Synthesis* **2008**, 4033–4035; (d) Bentabed-Ababsa, G.; Blanco, F.; Derdour, A.; Mongin, F.; Trécourt, F.; Quéguiner, G.; Ballesteros, R.; Abarca, B. *J. Org. Chem.* **2009**, *74*, 163–169.
15. We chose CuCl<sub>2</sub> as copper salt source instead of CuCl because of its higher air stability. In addition, the presence of TMEDA makes salts less sensitive to moisture, and sometimes favors deprotonation reactions: see Refs. 7d–f. For the synthesis of CuCl<sub>2</sub>-TMEDA, see: Handley, D. A.; Hitchcock, P. B.; Lee, T. H.; Leigh, G. J. *Inorg. Chem. Acta* **2001**, *316*, 59–64.
16. James, A. M.; Laxman, R. V.; Fronczek, F. R.; Maverick, A. W. *Inorg. Chem.* **1998**, *37*, 3785–3791.
17. The reaction of lithiated organics with molecular oxygen is well-documented: Wheatley, A. E. H. *Chem. Soc. Rev.* **2001**, *30*, 265–273. The process is thought to involve a radical chain decomposition in which a peroxide intermediate degrades to an organooxide product.
18. An organic radical was also observed ( $g = 2.005$ ,  $a_H = 1.096$  G, 2H), but was not identified:



19. TEMPO is similarly formed by preparing LiTMP in THF.
20. The spectral data are analogous to those obtained from a commercial sample.
21. The <sup>1</sup>H NMR data are analogous to those described: Azadi-Ardakani, M.; Wallace, T. W. *Tetrahedron* **1988**, *44*, 5939–5952.
22. The physical and spectral data are analogous to those previously described: Yuan, Y.; Bian, Y. *Appl. Organomet. Chem.* **2008**, *22*, 15–18.
23. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 0.78 (s, 6H), 1.23 (s, 6H), 1.49–1.64 (m, 6H), 3.75 (s, 3H), 6.82–6.88 (m, 2H), 7.13–7.19 (m, 1H), 7.29 (dd, 1H,  $J = 8.3$  and 1.9 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 18.6, 26.0, 31.5, 41.8, 54.4, 55.1, 111.0, 119.3, 126.4, 134.2, 136.0, 160.3.
24. The <sup>1</sup>H NMR data are analogous to those described: Ochiai, M.; Fujita, E.; Arimoto, M.; Yamaguchi, H. *Chem. Pharm. Bull.* **1982**, *30*, 3994–3999.
25. The <sup>1</sup>H NMR data are analogous to those described: Chen, X.; Yu, M.; Wang, M. *J. Chem. Res.* **2005**, 80–81.
26. The spectral data are analogous to those previously described: Waterlot, C.; Hasiak, B.; Couturier, D.; Rigo, B. *Tetrahedron* **2001**, *57*, 4889–4901.
27. The spectral data are analogous to those previously described: Rao, M. L. N.; Venkatesh, V.; Banerjee, D. *Tetrahedron* **2007**, *63*, 12917–12926.

28. The spectral data are analogous to those previously described: Inamoto, K.; Katsuno, M.; Yoshino, T.; Arai, Y.; Hiroya, K.; Sakamoto, T. *Tetrahedron* **2007**, *63*, 2695–2711.
29. The <sup>1</sup>H NMR data are analogous to those described: Güngör, T.; Marsais, F.; Quéguiner, G. *J. Organomet. Chem.* **1981**, *215*, 139–150.
30. The <sup>1</sup>H NMR data are analogous to those described: Trécourt, F.; Marsais, F.; Güngör, T.; Quéguiner, G. *J. Chem. Soc., Perkin Trans. 1* **1990**, 2409–2415.
31. Compound **11a**: yellow powder; mp 90 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.37 (ddd, 1H,  $J = 7.6, 4.9$ , and 1.9 Hz), 7.44–7.50 (m, 2H), 7.73–7.79 (m, 2H), 8.04 (ddd, 1H,  $J = 9.4, 7.5$ , and 2.0 Hz), 8.43 (ddd, 1H,  $J = 4.9, 2.1$ , and 1.2 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 121.3 (d,  $J = 30$  Hz), 121.9 (d,  $J = 4.5$  Hz), 129.2 (s, 2C), 131.1 (d, 2C,  $J = 1.2$  Hz), 135.0 (d,  $J = 0.9$  Hz), 140.7 (s), 142.0 (d,  $J = 3.3$  Hz), 150.9 (d,  $J = 15$  Hz), 160.1 (d,  $J = 243$  Hz), 190.7 (d,  $J = 4.9$  Hz).
32. Compound **13a**: beige powder; mp 153 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.31–7.36 (m, 2H), 7.88–7.96 (m, 2H), 8.30 (dd, 2H,  $J = 4.9$ , and 1.9 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 116.7 (m), 121.7 (m), 142.0 (t,  $J = 3.3$  Hz), 148.1 (m), 160.4 (d,  $J = 241$  Hz).
33. Mandeville, W. H.; Whitesides, G. M. *J. Org. Chem.* **1974**, *39*, 400–405.
34. Even if the basicity of pyrimidine nitrogens is low compared with that of pyridine, competitive quaternarization by reaction with allyl bromide and methyl iodide is not impossible under the conditions used.
35. Compound **15**: yellow oil; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 3.95 (s, 3H), 4.07 (s, 3H), 7.42–7.49 (m, 2H), 7.58 (tt, 1H,  $J = 7.3$  and 1.3 Hz), 7.74–7.78 (m, 2H), 8.46 (br s, 1H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 54.5, 55.5, 114.5, 128.5 (2C), 129.7 (2C), 133.3, 137.6, 161.2, 166.3, 169.3, 192.2. Compound **16**: yellow powder; mp 145–146 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.48 (s, 1H), 7.68–7.73 (m, 2H), 7.41–7.46 (m, 2H), 4.08 (s, 3H), 3.96 (s, 3H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 54.5, 55.5, 114.1, 128.8 (2C), 130.9 (2C), 136.0, 139.7, 161.3, 166.5, 169.1, 190.9. Compound **17**: yellow oil; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 3.22 (m, 2H), 3.96 (s, 3H), 3.97 (s, 3H), 5.00–5.08 (m, 2H), 5.82–5.97 (m, 1H), 7.98 (br s, 1H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 29.8, 54.0, 54.8, 113.6, 116.5, 135.2, 157.0, 164.4, 169.4. Compound **19**: red powder; mp 209 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 3.97 (s, 6H), 4.03 (s, 6H), 8.20 (s, 2H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 54.3 (2C), 55.0 (2C), 108.3 (2C), 158.8 (2C), 165.1 (2C), 168.7 (2C).
36. The spectral data are analogous to those previously described: Boudet, N.; Dubbaka, S. R.; Knochel, P. *Org. Lett.* **2008**, *10*, 1715–1718.
37. Typical procedure: To a stirred, cooled (0 °C) suspension of CuCl<sub>2</sub>-TMEDA (0.25 g, 1.0 mmol) in THF (5 mL) were successively added BuLi (about 1.6 M hexanes solution, 1.0 mmol) and, 15 min later, a solution of LiTMP prepared in THF (2 mL) at 0 °C from 2,2,6,6-tetramethylpiperidine (0.34 mL, 2.0 mmol) and BuLi (about 1.6 M hexanes solution, 2.0 mmol). The mixture was stirred for 15 min at this temperature before introduction of 2,4-dimethoxypyrimidine (125 μL, 1.0 mmol). After 2 h at room temperature, benzoyl chloride (0.24 mL, 2.0 mmol) was added at 0 °C. The mixture was stirred for 16 h at 60 °C before the addition of brine (5 mL) and extracted with Et<sub>2</sub>O (3 × 10 mL). The combined organic layers were washed with brine (10 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated under reduced pressure before purification by column chromatography on silica gel (eluent: heptane/EtOAc 8:2). Compound **15** (0.11 g, 45%) was isolated as a yellow oil.
38. Forbes, G. C.; Kennedy, A. R.; Mulvey, R. E.; Rodger, P. J. *J. Chem. Commun.* **2001**, 1400–1401; See also: Balloch, L.; Drummond, A. M.; García-Álvarez, P.; Graham, D. V.; Kennedy, A. R.; Klett, J.; Mulvey, R. E.; O'Hara, C. T.; Rodger, P. J. A.; Rushworth, I. D. *Inorg. Chem.* **2009**, *48*, 6934–6944.