



Corrigendum

Corrigendum to “Chrysanthemyl 2-acetoxy-3-methylbutanoate: the sex pheromone of the citrophilous mealybug, *Pseudococcus calceolariae*” [Tetrahedron Lett. 51 (2010) 1075]

Ashraf M. El-Sayed^{a,*}, C. Rikard Unelius^{a,b}, Andrew Twidle^a, Vanessa Mitchell^a, Lee-Anne Manning^a, Lyn Cole^c, David M. Suckling^a, M. Fernanda Flores^d, Tania Zaviezo^e, Jan Bergmann^d

^a The New Zealand Institute for Plant & Food Research Ltd, (PFR) PB 4704, Christchurch, New Zealand

^b School of Pure and Applied Natural Sciences, University of Kalmar, SE-391 82 Kalmar, Sweden

^c PFR, Hawke's Bay Research Centre, Havelock North, Hastings 4157, New Zealand

^d Instituto de Química, Pontificia Universidad Católica de Valparaíso, Avda. Brasil 2950, Valparaíso, Chile

^e Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile, Avda. Vicuña Mackenna 4860, Santiago, Chile

The legend within Figure 3 should read ‘Dose (μg)’ instead of ‘Dose (mg)’. This has been updated in the figure shown below. The authors regret the error.




Figure 3. Mean catch ± SE of male citrophilus mealybugs in red delta traps baited with various doses of racemic chrysanthemyl 2-acetoxy-3-methylbutanoate **3** (0.1, 1, 10, 100 and 1000 μg) loaded on red rubber septa. Letters on columns indicate significant differences (FPLS test, $P < 0.05$).

DOI of original article: 10.1016/j.tetlet.2009.12.106.

* Corresponding author. Tel./fax: +64 3 977 7358.

E-mail address: ael-sayed@plantandfood.co.nz (A.M. El-Sayed).