

Contents lists available at ScienceDirect

Tetrahedron Letters





Tetrahedron Letters Vol. 50, No. 17, 2009

Contents

Communications

A convenient synthesis of *ortho-ortho* disubstituted biphenyls containing an eight-membered lactam ring using radical chemistry

pp 1879-1881

C. H. Wang *, S. Alluri, A. K. Ganguly

Structure and reactivity of a chiral cyclononadienone

pp 1882-1885

Yue Zhang, Stephen D. Lotesta, Thomas J. Emge, Lawrence J. Williams *

Convenient preparation of unsymmetrical 2,5-disubstituted 1,3,4-oxadiazoles promoted by Dess-Martin reagent

pp 1886-1888

Cristian Dobrotă *, Codruța C. Paraschivescu, Ioana Dumitru, Mihaela Matache, Ion Baciu, Lavinia L. Ruță

$$R^1$$
 R^2
 R^1
 R^2
 R^2
 R^3
 R^2
 R^3
 R^2
 R^3
 R^3
 R^4
 R^5
 R^4
 R^5
 R^5
 R^5
 R^6
 R^6

2,5-Disubstituted 1,3,4-oxadiazoles have been conveniently prepared by oxidative cyclization of *N*-acyl-*N*-aryliden-hydrazines promoted by an excess of Dess-Martin reagent under mild conditions (23 examples, up to 92% isolated yields).

A practical method to access enantiopure β -perfluoroalkyl- β -amino acids: diastereoselective reduction of cyclic enamino-esters

pp 1889-1892

Yasuhiro Ishida *, Nobutaka Iwahashi, Nao Nishizono, Kazuhiko Saigo *


F₃C OMe F₃C F₃C F₃C OH Ph OH Ph (d.r. = 97:3)
$$(>99\% \text{ ee})$$



pp 1893-1896

Studies of multicomponent Kinugasa reactions in aqueous media

Craig S. McKay, David C. Kennedy, John Paul Pezacki



Studies of the micelle-promoted and copper-catalyzed multicomponent Kinugasa reaction in water are reported.



Fluorinated squaraine as near-IR label with improved properties for the labeling of oligonucleotides Brice-Loïc Renard, Yves Aubert, Ulysse Asseline *

pp 1897-1901

$$(CH_2)_6\text{-S-p} \underbrace{Oligonucleotides}_{\text{N}}$$

Fluorinated squaraine-oligonucleotide conjugates: detection beyond 670 nm and high quantum yields (0.27-0.39).



First phosphorous p-xylose-derived glycodendrimers

pp 1902-1905


Caroline Hadad, Jean-Pierre Majoral, Jacques Muzart, Anne-Marie Caminade, Sandrine Bouquillon *

pentoses + P-dendrimers

Heterocyclic columnar hexacatenar bisthiazoles

pp 1906-1910

Hui-Hsu Gavin Tsai, Lung-Chun Chou, Sheng-Chia Lin, Hwo-Shuenn Sheu, Chung K. Lai *





Synthesis of substituted 1-benzazepin-2-ones via ring-closing olefin metathesis

Scott B. Hoyt *, Clare London, Min Park

pp 1911-1913



Extending Pummerer reaction chemistry. Examination of the prospects for forming vicinal quaternary carbon centers

pp 1914-1916

Ken S. Feldman *, Ahmed Yimam Nuriye

$$\begin{array}{c|c} \text{PhMe}_2\text{SiO} & \text{O} \\ \text{O}^- & \text{R} & \text{Tf}_2\text{O} \\ \text{N} & \text{Ph} & 2,6\text{-lutidine} \end{array}$$

$A symmetric \ synthesis \ of \ 2-alkyl-4-hydroxycyclohex-2-en-1-ones \ by \ scandium (III) \ triflate-catalyzed \ fragmentation \ of \ 2-alkyl-3-iodo-1-oxocyclohexan-2,4-carbolactones$

pp 1917-1919

Jun-ichi Matsuo *, Mizuki Kawano, Kosuke Takeuchi, Hiroyuki Tanaka, Hiroyuki Ishibashi

Ring opening of aziridines with *ortho*-bromophenyl metal reagents: synthesis of 2-substituted indolines David J. Michaelis, Thomas A. Dineen *

pp 1920-1923

Triflic acid promoted synthesis of polycyclic aromatic compounds

pp 1924-1927

Ang Li, Daniel J. DeSchepper, Douglas A. Klumpp *

Conjugate addition of sodium methanesulfinate to vinyl pyridines and diazines for the synthesis of aliphatic sulfones

pp 1928-1933

Gregory M. Schaaf *, Sabuj Mukherjee, Alex G. Waterson

A closer insight into the mechanism operating in the trifluoroacetylation of pyrrole. New trifluoromethyl pyrroylmethane discovered

pp 1934-1938

W. J. Peláez, M. A. Burgos Paci, G. A. Argüello *



A thorough revision of the mechanism of trifluoroacetylation of pyrrole, identification, isolation, and characterization of a new pyrroylmethane probably missed in the synthesis of fluorinated porphyrins, is reported.



An efficient solution phase synthesis of triazadibenzoazulenones: 'designer isonitrile free' methodology enabled by microwaves

pp 1939-1942

Christopher Hulme *, Shashi Chappeta, Chris Griffith, Yeon-Sun Lee, Justin Dietrich

A novel two-step synthesis of triazadibenzoazulenones is described. The methodology represents the first example of a post-condensation Ugi modification that employs two 'internal nucleophiles'.

Highly enantioselective organocatalytic synthesis of piperidines. Formal synthesis of (-)-Paroxetine

pp 1943-1946

Guillem Valero, Jiri Schimer, Ivana Cisarova, Jan Vesely *, Albert Moyano *, Ramon Rios *

Difunctional additions to 1-cyclopropylallenes: an efficient and stereospecific method for the synthesis of 2,6-difunctional-1,3-hexadienes

pp 1947-1950

Bo Meng, Lei Yu, Xian Huang

$$R^{1} = C \longrightarrow R^{2} + E^{+} + Nu^{-} \longrightarrow R^{1}$$

$$E = I, Br, CI, PhSe$$

$$Nu = I, Br, CI, OR, OAc, NCS, NHCOPh$$

$$R^{1} \longrightarrow R^{2}$$

$$R^{1} \longrightarrow R^{2}$$

$$R^{1} \longrightarrow R^{2}$$

$$R^{1} \longrightarrow R^{2}$$


The difunctional additions of electrophiles and nucleophiles to 1-cyclopropylallenes were investigated. Two different functional groups were introduced at the same time to give 2,6-difunctional-1,3-hexadienes stereoselectively in good yields.



Dual signal (color change and fluorescence ON-OFF) ensemble system based on bis(Dpa-Cu^{II}) complex for detection of PPi in water

pp 1951-1953

Soon Young Kim, Jong-In Hong *





Formation of six- versus five-membered cyclic sulfones by C-H insertion

pp 1954-1957

Christian S. Jungong, Jinu P. John, Alexei V. Novikov *

Selectivity of six- versus five-membered ring formation in C-H insertion on alkylsulfonyl diazoacetates is sensitive to the substrate structure and catalyst used.



lodine-catalyzed one-pot three-component synthesis of homoallyl benzyl ethers from aldehydes

pp 1958-1960

Dolly Kataki, Prodeep Phukan *

A synthesis of esters, amides, and sulfones bearing a 1-cyclopentenyl group at the α -position from cyclobutanones with one-carbon ring-expansion

pp 1961-1964

Tsuyoshi Satoh *, Yu Awata, Shingo Ogata, Shimpei Sugiyama, Masami Tanaka, Motoo Tori

$$R^{1} \xrightarrow{\text{CI}} \text{LiCHX} \xrightarrow{\text{R}^{2}} \text{R}^{2} \xrightarrow{\text{R}^{3}\text{MgCI}} \text{R}^{3} \xrightarrow{\text{MgCI}} \text{MgCI} \xrightarrow{\text{insertion}} \text{R}^{1} \xrightarrow{\text{R}^{2}} \text{X}$$

$$X = \text{COO}^{t}\text{Bu, CON(CH}_{3})_{2}, \text{SO}_{2}\text{Ph}$$

An inexpensive and highly stable palladium(II) complex for room temperature Suzuki coupling reactions under ambient atmosphere

pp 1965-1968

Mengping Guo *, Qiaochu Zhang

An approach to the synthesis of dimeric resveratrol natural products via a palladium-catalyzed domino reaction Jenna L. Jeffrey, Richmond Sarpong *

pp 1969-1972

A route for the rapid assembly of the carbon framework of several resveratrol natural products is presented. A palladium-catalyzed domino reaction of bromostilbene derivative **6** and tolane **7**, involving two sequential Heck coupling reactions, provides access to the benzofulvene-based core of various resveratrol-derived natural products. The carbon skeleton of pallidol and its congeners is achieved by a Lewis acid-induced Nazarov-type oxidative cyclization of **9**.



1-Bromo-2-(diphenylphosphinoyl)ethyne and 1-bromo-2-(p-tolylsulfinyl)ethyne: versatile reagents eventually leading to benzocyclotrimers

pp 1973-1976

Pierluigi Padovan, Stefano Tartaggia, Silvia Lorenzon, Enrico Rosso, Cristiano Zonta, Ottorino De Lucchi, Fabrizio Fabris *

p-Phenylene sulfide oligomers and their properties. Ar–S couplings mediated by copper or by fluorine substitutions

pp 1977-1981

Olivier Goyot, Marc Gingras *

A series of monodisperse PPS oligomers of various length were synthesized by Cu-cat. Ar–S couplings or by fluorine aromatic substitutions with aryl thiolates. Fluorine chemistry brings new advantages such a greater solubility, reactivity and easier characterization by ¹⁹F/¹³C NMR. Crystallinity studies of a series of PPS oligomers were investigated.

New gem-difluoromethylene-containing isocyanide as a useful building block for the synthesis of difluorinated pseudopeptides via Ugi reaction

pp 1982-1985

Nianjin Liu, Song Cao *, Li Shen, Jingjing Wu, Jinlong Yu, Jian Zhang, Hui Li, Xuhong Qian *

A new and efficient method was developed for the synthesis of novel *gem*-difluoromethylene-containing isocyanide, which can be used as a building block for the synthesis of difluorinated pseudopeptides via Ugi reaction.



An improved amide coupling procedure for the synthesis of N-(pyridin-2-yl)amides

Allyn T. Londregan *, Gregory Storer, Ceshea Wooten, Xiaojing Yang, Joseph Warmus

pp 1986-1988



Novel and versatile protocol for the preparation of functionalized benzocyclotrimers

Erdin Dalkılıç, Murat Güney, Arif Daştan *, Nurullah Saracoglu, Ottorino De Lucchi, Fabrizio Fabris '

pp 1989-1991

Synthesis and mesomorphic behaviour of new mesogenic compounds possessing a cholesteryl ester moiety connected to a pyrimidine core

pp 1992-1995

K. C. Majumdar *, Shovan Mondal, Nilasish Pal, Randhir Kumar Sinha

$$\begin{array}{c} R \\ O \\ O \\ N \\ \end{array}$$

$$\begin{array}{c} O \\ O \\ \end{array}$$

A novel synthesis of N_1 -(substituted)-pyrimido[5,4-e]-1,2,4-triazine-5,7(1H,6H)-diones

pp 1996-1997

Anjanette J. Turbiak, H. D. Hollis Showalter *

Crystallization-based optical resolution of 1,1'-binaphthalene-2,2'-dicarboxylic acid via 1-phenylethylamides: control by the molecular structure and dielectric property of solvent

pp 1998-2002


Yuki Kato, Yuichi Kitamoto, Naoya Morohashi, Yosuke Kuruma, Shuichi Oi, Kenichi Sakai, Tetsutaro Hattori *

Control by the molecular structure of solvent toluene, AcOEt, 1-PrOH, 2-PrOH, EtOH acetone—hexane—1-PrOH
$$8.9 \le \varepsilon \le 10.2$$
 (RS_a, S)-1•solvent (solvent = acetone, CH₂Cl₂, or MeCN) (solvent = none or EtOH)

Oryzamutaic acid A, a novel yellow pigment from an Oryza sativa mutant with yellow endosperm

pp 2003-2005

Hiroshi Nakano *, Seiji Kosemura, Toshisada Suzuki, Katsutoshi Hirose, Ryota Kaji, Makoto Sakai




Oryzamutaic acid A, a novel yellow pigment, was isolated from the endosperm (polished rice) of an *Oryza sativa* mutant. The structure and absolute configuration of oryzamutaic acid A were elucidated on the basis of spectroscopic analysis, single-crystal X-ray diffraction analysis, and biogenetic reason.



A preliminary step toward molecular spring driven by cooperative guest binding

pp 2006-2009

Tomohiro Ikeda, Seiji Shinkai *, Kazuki Sada, Masayuki Takeuchi *





Fluorescent and colorimetric detection of acid vapors by using solid-supported rhodamine hydrazides

pp 2010-2012


Shincheol Kang, Sungwook Kim, Young-Keun Yang, Shinhyo Bae, Jinsung Tae



Metal ion induced FRET On-Off in naphthyl-pyrenyl pendent tetrahomodioxacalix[4]arene

pp 2013-2016

Ji Hee Jung, Min Hee Lee, Hyun Jung Kim, Hyo Sung Jung, Su Yeon Lee, Na Ri Shin, Kwanghyun No *, Jong Seung Kim *



Michael additions of primary and secondary amines to acrylonitrile catalyzed by lipases

pp 2017-2018

Rodrigo O. M. A. de Souza *, Lilian M. C. Matos, Karen M. Gonçalves, Ingrid C. R. Costa, Ivelize Babics, Selma G. F. Leite, E. G. Oestreicher, O. A. C. Antunes




Diterpenoids from Isodon pharicus

pp 2019-2023

Yong Zhao, Sheng-Xiong Huang, Wei-Lie Xiao, Li-Sheng Ding, Jian-Xin Pu, Xian Li, Li-Bin Yang, Han-Dong Sun *

A phytochemical investigation of *Isodon pharicus* led to the isolation of a novel asymmetric *ent*-kauranoid dimer, bispseurata F (1), and three new diterpenoids pharicinins **A–C** (**2–4**). Their structures were elucidated by extensive spectroscopic analysis. Compound **1** features a unique linkage pattern of C-17 with C-11' to connect the two monomers. A possible biogenetic pathway of **1** was also proposed. Compounds **3** and **4**





exhibited moderate inhibitory activity against NB4 and SH-SY5Y cell lines. OTHER CONTENTS

Erratum

p 2024

Corrigenda

pp 2025-2026

*Corresponding author

(**P** Supplementary data available via ScienceDirect

Abstracted/indexed in: AGRICOLA, Beilstein, BIOSIS Previews, CAB Abstracts, Chemical Abstracts, Chemical Engineering and Biotechnology Abstracts, Current Biotechnology Abstracts, Current Contents: Life Sciences, Current Contents: Physical, Chemical and Earth Sciences, Current Contents Search, Derwent Drug File, Ei Compendex, EMBASE/Excerpta Medica, Medline, PASCAL, Research Alert, Science Citation Index, SciSearch. Also covered in the abstract and citation database SCOPUS[®]. Full text available on ScienceDirect[®]



